INFORMATION TO USERS

This reproduction was made from a copy of a manuscript sent to us for publication and microfilming. While the most advanced technology has been used to photograph and reproduce this manuscript, the quality of the reproduction is heavily dependent upon the quality of the material submitted. Pages in any manuscript may have indistinct print. In all cases the best available copy has been filmed.

The following explanation of techniques is provided to help clarify notations which may appear on this reproduction.

1. Manuscripts may not always be complete. When it is not possible to obtain missing pages, a note appears to indicate this.
2. When copyrighted materials are removed from the manuscript, a note appears to indicate this.
3. Oversize materials (maps, drawings, and charts) are photographed by sectioning the original, beginning at the upper left hand corner and continuing from left to right in equal sections with small overlaps. Each oversize page is also filmed as one exposure and is available, for an additional charge, as a standard 35mm slide or in black and white paper format.*
4. Most photographs reproduce acceptably on positive microfilm or microfiche but lack clarity on xerographic copies made from the microfilm. For an additional charge, all photographs are available in black and white standard 35mm slide format.*

***For more information about black and white slides or enlarged paper reproductions, please contact the Dissertations Customer Services Department.**

U·M·I Dissertation
Information Service

University Microfilms International
A Bell & Howell Information Company
300 N. Zeeb Road, Ann Arbor, Michigan 48106

8618808

Levine, Wendy Beth

ISOLATION AND CHARACTERIZATION OF CADMIUM-RESISTANT MUTANTS
OF NEUROSPORA CRASSA

The Ohio State University

Ph.D. 1986

University
Microfilms
International 300 N. Zeeb Road, Ann Arbor, MI 48106

PLEASE NOTE:

In all cases this material has been filmed in the best possible way from the available copy. Problems encountered with this document have been identified here with a check mark ✓.

1. Glossy photographs or pages ✓
2. Colored illustrations, paper or print ✓
3. Photographs with dark background ✓
4. Illustrations are poor copy _____
5. Pages with black marks, not original copy _____
6. Print shows through as there is text on both sides of page _____
7. Indistinct, broken or small print on several pages ✓
8. Print exceeds margin requirements _____
9. Tightly bound copy with print lost in spine _____
10. Computer printout pages with indistinct print _____
11. Page(s) _____ lacking when material received, and not available from school or author.
12. Page(s) _____ seem to be missing in numbering only as text follows.
13. Two pages numbered _____. Text follows.
14. Curling and wrinkled pages _____
15. Dissertation contains pages with print at a slant, filmed as received _____
16. Other _____

University
Microfilms
International

ISOLATION AND CHARACTERIZATION OF CADMIUM-RESISTANT
MUTANTS OF NEUROSPORA CRASSA

DISSERTATION

Presented in partial fulfillment of the requirements for
the Degree Doctor of Philosophy in the Graduate
School of The Ohio State University

By

Wendy Beth Levine, B.S.

* * * * *

The Ohio State University

1986

Reading Committee:

Dr. G. A. Marzluf

Dr. T. J. Byers

Dr. L. F. Johnson

Dr. P. S. Perlman

Approved by

Advisor
Molecular, Cellular, and
Developmental Biology
Program

To my mother and father

ACKNOWLEDGEMENTS

I would like to acknowledge the Molecular, Cellular, and Developmental Biology Program and the Department of Biochemistry for providing me with the opportunity to pursue this research, and my advisor, Dr. George A. Marzluf, for his patience and guidance during my graduate training. I would also like to thank the other members of my advisory committee, Drs. Thomas J. Byers, Lee F. Johnson, and Philip S. Perlman, for their suggestions and support.

I am grateful to Susan Maxim and Erika Karlak for their valuable assistance in the preparation of this manuscript.

I extend special thanks to all of my friends who have made my tenure as a graduate student more enjoyable.

Finally, I would like to thank my parents, for without their emotional and financial support, this dream might not have been fulfilled.

VITA

1957.	Born - Dayton, Ohio
1979.	B.S., Biological Sciences Indiana University, Bloomington, Indiana
1979-1985	Graduate Teaching Associate, The Ohio State University, Columbus, Ohio
1979-1986	Graduate Research Associate, Molecular, Cellular, and Developmental Biology Program, The Ohio State University, Columbus, Ohio

PUBLICATION

Calcott, P. H., C. Zaborowski, W. B. Levine, and N. H. Troung. 1979. Drug-resistance plasmid (pPL1) mediated changes in the susceptibility of Pseudomonas aeruginosa to stress. FEMS Microbiol. Lett. 6:75-80.

TABLE OF CONTENTS

	Page
DEDICATION.	ii
ACKNOWLEDGEMENTS.	iii
VITA.	iv
LIST OF TABLES.	vii
LIST OF FIGURES	viii
LIST OF PLATES.	x
INTRODUCTION.	1
MATERIALS AND METHODS	33
Strains.	33
UV Mutagenesis	33
Cadmium-induced Mutagenesis.	35
Training Flasks.	36
Growth Rate Plates	37
Death <u>vs.</u> Inhibition	38
Crosses.	38
Cross Analysis Plates.	39
Growth Rate Flasks	39
Cross-resistance	43
Heat Shock/Cadmium Shock	43
Transport Studies.	44
Cadmium/Hemoglobin Assay	46
Radioactive Labelling of Proteins <u>In Vivo</u>	47
Protein Quantification	48
Electrophoresis.	48
Fluorography	49
RESULTS	50
Isolation of Cadmium-resistant Mutants	50
Genetic Analysis of Mutants.	64

	Page
Characterization of Mutants Under Various Growth Conditions.	74
Cadmium Transport.	91
Cadmium-binding Potential.	120
Heat Shock/Cadmium Shock	125
Effect of Cadmium on Cytoplasmic Protein Profile.	126
DISCUSSION.	138
LIST OF REFERENCES.	164

LIST OF TABLES

Table	Page
1 <u>Neurospora</u> Strains	34
2 Cadmium-induced mutation frequency on various concentrations of CdCl_2	56
3 Growth rates of UV-induced Cd-resistant mutants	61
4 Growth rates of Cd-induced Cd-resistant mutants	62
5 Growth of Specific Cd-resistant mutants in liquid selective medium.	63
6 Percentage of Cd^R and Cd^S progeny	65
7 Assignment of linkage group for Cd^R mutants 9A2R5, 15-10R22, 4A2R1, and TFR38 .	
8 Assignment of map location for Cd^R mutants 15-10R22 and TFR38.	70
9 Assignment of map location for Cd^R mutant 4A2R1.	71
10 Assignment of map location for Cd^R mutant 9A2R5.	72
11 Evaluation of progeny from crosses between Cd-resistant mutants.	75
12 Mycelial levels of cadmium binding capacity for wild-type and the cadmium-resistant mutants in the absence and presence of cadmium	124

LIST OF FIGURES

Figure	Page
1 Growth Rate of <u>N. crassa</u> wild-type strains on various concentrations of CdCl ₂	51
2 UV Mutagenesis Survival Curve	53
3 Growth of wild-type strain on various concentrations of CdCl ₂	79
4 Percent of control growth rate for the wild- type and the Cd ^R mutants TFR38, 9A2R5, and 15-10R22 on various concentrations of CdCl ₂	81
5 Growth of Cd ^R mutant TFR38 on various concentrations of CdCl ₂	83
6 Growth of Cd ^R mutant 9A2R5 on various concentrations of CdCl ₂	85
7 Growth of Cd ^R mutant 15-10 on various concentrations of CdCl ₂	87
8 Percent of wild-type growth for the Cd ^R mutants TFR38, 9A2R5, and 15-10R22 on various concentrations of CdCl ₂	89
9 Transport of cadmium into wild-type conidia .	92
10 Transport of cadmium into wild-type mycelia .	94
11 Transport of cadmium into wild-type mycelia for various concentrations of CdCl ₂	97
12 Effect of cadmium concentration on the rate of cadmium transport into mycelia	99
13 Effect of azide on transport of cadmium by wild-type mycelia	102

	Page
14 Transport of cadmium into wild-type mycelia in the presence of magnesium, manganese, or zinc	104
15 Transport of cadmium into wild-type mycelia in the presence of cobalt, copper, iron, or nickel.	106
16 Transport of cadmium into wild-type mycelia in the presence of copper and manganese . . .	109
17 The effect of cycloheximide (CHX) on the transport of cadmium into wild-type mycelia in the presence of copper	112
18 Transport of leucine into wild-type mycelia and the effect of CHX on leucine incorporation into proteins	114
19 Transport of cadmium into wild-type and Cd ^R mutant mycelia.	116
20 Effect of azide on cadmium transport into wild- type and Cd ^R mutant mycelia	118
21 Efflux of cadmium in the presence of various concentrations of metals.	121
22 Electrophoretic pattern of Coomassie-stained [³⁵ S]methionine-labelled heat shock and cadmium shock proteins.	127
23 Fluorogram of [³⁵ S]methionine-labelled proteins for the same gel electrophoretic pattern in Figure 22.	129
24 Electrophoretic pattern of Coomassie-stained [³⁵ S]O ₄ -labelled proteins from wild- type and Cd ^R mycelia grown in the absence and presence of cadmium	132
25 Fluorogram of [³⁵ S]O ₄ proteins for the same gel electrophoretic pattern in Figure 24	134

LIST OF PLATES

Plate		Page
I	Cadmium-sensitive and cadmium-resistant isolates grown in the absence and presence of cadmium	40
II	Cadmium-sensitive and cadmium-resistant isolates grown in the presence of 15, 25, or 30 mM CdCl ₂	42
III	Comparison between the wild-type strain and Training Flask mutant grown in the presence of 13 mM CdCl ₂	59

INTRODUCTION

Cadmium is most commonly found as cadmium sulfide in ores of copper, lead, and zinc. Sulfide ore mines and refineries are prevalent in Japan, with a large number in the Toyama Prefecture area (Friberg, 1974). Much of the arable land in Japan is used for rice crops, and often, unfortunately, the polluted run-off water from the smelting operations is used to irrigate the fields. In the 1940's and 1950's a physician in the vicinity of Toyama Prefecture diagnosed a vast number of cases of patients suffering from "Itai-itai", or "ouch-ouch" disease, exhibiting symptoms of kidney trouble, rheumatic disease, and the demineralization of bones. At the same time, the farmers in that area experienced serious damage to their rice fields and blamed the local mining company for polluting the river water used for irrigation. Subsequent investigation of the cadmium content of rice from that area revealed that it was greater than ten times that found in rice samples from other areas of Japan, which substantiated the farmers' claim that contaminating metals in the river water were responsible for the crop damage. In 1961 Japanese physicians postulated that

cadmium played a causative role in Itai-itai disease and the issues of cadmium pollution and toxicity are now of great importance in Japan. Since that time, cadmium has also been implicated as a cause of lung cancer by the National Institute for Occupational Safety and Health and is now regarded as a potential carcinogen (NIOSH, 1984).

Ecosystems are easily disturbed by small changes in the concentration of pollutants in the environment and the outcome can include changes in a species' physiology or reproduction, or altered trophic interactions among species. Considering man's great role in the introduction of high levels of contaminating heavy metals into the environment from industrial processes and the serious effects on man as a consumer of aquatic life forms, many field and laboratory studies exist on cadmium toxicity and tolerance in an array of organisms from protozoans to fungi, plants, and animals.

Summers and Silver (1978) reviewed the currently accepted interactions among the environment, the biota, and heavy metals from the standpoint of microbial transformations of heavy metals, especially mercury and cadmium. They discussed tolerant strains of Pseudomonas and Staphylococcus aureus from both biochemical and genetic aspects. Chroococcus paris, a blue-green alga and important member of the trophic food chain, can concentrate metals from the environment, including

cadmium, zinc, and copper, up to 4000-fold (Les and Walker, 1984). This factor definitely presents a hazardous situation to those organisms in higher trophic levels. Aquatic hyphomycetes are less sensitive to cadmium in hard stream water than in soft stream water, probably due to the higher calcium levels competing for cadmium binding sites (Abel and Bärlocher, 1984). Minnows taken from metal-contaminated flyash ponds exhibited significant metal tolerance when compared to those raised in hatchery ponds (Benson and Birge, 1985). After the fish were transferred from a metal-contaminated pond to contaminant-free water, their metal tolerance decreased significantly, whereas the hatchery minnows exhibited tolerance induction when maintained on sub-lethal concentrations of cadmium. The induced tolerance was lost upon transfer back to contaminant-free water.

A vast amount of information involving laboratory studies on the effects of cadmium on an array of different organisms and cultured cells is available. Cadmium induces a number of responses in Drosophila melanogaster including retardation of development and an increase in the time required for eclosion. There is also a significant delay in the appearance and disappearance of developmentally-expressed proteins (Jacobson et al., 1981). Cell culture systems, including Chinese hamster

kidney cells (De Ruiter et al., 1985) and primary cultures of rat hepatocytes (Santone et al., 1982) have been used to demonstrate cadmium's inhibitory effect on cell proliferation, cell viability, and the metabolic parameter, glucose consumption. At the microscopic level, rat liver cells cultured in the presence of cadmium show a series of morphological effects in the following order: irreversible chromatin condensation, a decrease in the amount of granular endoplasmic reticulum, an increase in vesicularization, and finally, cell membrane leakage and cell death. A potential involvement of a disturbance in ribosomal RNA synthesis has been hypothesized based on previous results (Morselt et al., 1983).

Cameron and Foster (1963; Foster and Cameron, 1963) performed an extensive investigation into the histological effects of sub-cutaneous injections of CdCl_2 on rabbit kidney and testis. One of the most noticeable effects was an increase in the number of glycoprotein-containing "stainable droplets". These droplets were similar to the protein inclusions seen after injection with foreign proteins, which appeared to enter the cells via pinocytosis; these foreign proteins first bound to sites in the membrane-associated mucopolysaccharide layer and then were segregated into membrane-bound droplets in the pinocytotic channels. Cadmium treatment apparently increased the cells' permeability to plasma proteins,

which were segregated into vesicles in the cytoplasm in a manner similar to that described above. Another visible effect of cadmium injection was the disruption of nuclear division in spermatogenic cells. Cadmium was presumed to interact with sulfhydryls, which have been reported as being abundant in the region of the mitotic spindle. Finally, cadmium also had a carcinogenic effect: one of the treated rabbits developed a small testicular neoplasm. Cadmium is considered to exert its mutagenic or carcinogenic effect by decreasing the fidelity of DNA synthesis (Hoffman and Niyogi, 1977). Cadmium affects the properties of zinc-containing enzymes involved in nucleic acid synthesis and complexes with nucleotides, altering the structure of nucleic acids (Jacobson and Turner, 1980).

Studies similar to those involving the acquisition of metal tolerance in minnows were performed on mammalian cells in culture. HeLa cells and cultured Chinese hamster cells can adapt to a toxic concentration of cadmium in the medium if they have been previously treated with a low non-toxic dose of cadmium (Fischer, 1985). This tolerance is not due to cadmium exclusion, but rather by an intracellular detoxification mechanism. Yoshikawa and Ohta (1982) reported that in rat liver, the acute toxicity of contaminant metals (Cd, Hb, Pb) was relieved by pretreatment with essential metals (Cu, Zn, Mn), via some

biological response other than the induction of heavy metal-binding proteins.

While searching for a biological role for the element cadmium, Margoshes and Vallee (1957) discovered the protein metallothionein (MT). When cadmium enters an organism it often combines with specific low molecular weight proteins which possess a high concentration of cysteine. It is thought that these proteins, termed metallothioneins, play a central role in metal metabolism, principally functioning in zinc metabolism and homeostasis and cadmium detoxification. The term metallothionein refers strictly to the metal-containing form of these low molecular weight proteins, the apoprotein being referred to as thionein. The designation Cd-thionein signifies that the protein contains only Cd. If other metals are also bound to the MT, they are listed in order of abundance, for example, (Cd, Cu, Zn)-MT.

The injection into laboratory animals of sub-toxic doses of cadmium salts causes an induction of both Cd-thionein and Zn-thionein, whose accumulation is predominantly in the liver and kidneys (Winge et al., 1975; Brady et al., 1979). Upon binding to MT, Cd becomes less available for excretion and for this reason, MT is thought to serve as a less-toxic storage form of the metal (Shaikh, 1982). In addition, cultured cells selected for Cd-resistance produce much higher amounts of MT (Beach and

Palmiter, 1981; Hildebrand et al., 1979) and take up much larger amounts of Cd than sensitive cells (Bakka and Rugstad, 1981; Rugstad and Norseth, 1978). As mentioned earlier, pretreatment of organisms with sub-lethal doses of heavy metals induces a tolerance to subsequent exposures; this tolerant response is accompanied by an increased concentration of cellular MT (Yoshikawa and Ohta, 1982; Kito et al., 1982b; Benson and Birge, 1985; Endreson et al., 1983). However, since heavy metals are not frequently encountered at high levels and because the basal level of MT expression prior to Cd exposure is relatively high, it seems unlikely that detoxification is the primary function of MT. Exogenously-administered Cd causes a disturbance in the endogenous Zn pool whereby the intracellular free Zn concentration increases, probably due to its displacement from MT by Cd (Brady et al., 1979). Brady has demonstrated in vitro the donation of Zn or Cu from MT to metal-requiring apoenzymes and hypothesizes that MT probably plays a significant role in tissues undergoing rapid growth and development (e.g. testes) by supplying Zn or Cu as cofactors for apoenzymes which take part in nucleic acid metabolism or protein synthesis. The peptide hormone thymulin, synthesized in the thymus gland, is located predominantly in the cytoplasmic vacuoles of the thymic epithelial cells. The

biological activity of thymulin requires the presence of equimolar amounts of zinc and, as expected, zinc is prevalent in the vacuoles of thymic epithelium. P.C. Huang (Danielson et al., 1982), using an immunostaining technique on histological sections of rat thymus epithelium, saw a pattern of MT distribution in these specialized cells, supporting Brady's proposed storage and transport role for MT (Savino et al., 1984). It is, therefore, generally accepted that MT plays an important role in heavy metal homeostasis, storage, and at higher metal concentrations, detoxification.

Metallothioneins have very specific characteristics which form the basis for the generally accepted criteria for their identification. MTs exhibit a very high metal content (approximately 9%), usually contain Cd, Zn, Hg, or Cu, and are of low molecular weight (6-11,000 daltons [dal]). Their cysteine content is approximately 30%, the highest of any known protein. Two unusual properties are the lack of absorption near 280 nm due to an absence of aromatic amino acids, and a strong absorption at 250nm resulting from cadmium-mercaptide bonds. Typically three to four cysteine residues bind one atom of cadmium, with 7 g atoms of cadmium bound per molecule of MT (Kagi et al., 1974). Boulanger has constructed a two-domain molecular model for mammalian MT based on data from ^{113}Cd NMR and has demonstrated that the 7 bound cadmium ions are

arranged in two separate metal clusters (Boulanger et al., 1983).

In vertebrates, 2 different isometallothioneins exist, MT-I and MT-II, which differ in up to 15 of the 61 amino acid positions. Equine MT-I exists in 2 variant forms, MT-I_A and MT-I_B. Most vertebrate MT amino acid sequences show a fixed distribution of cysteinyl residues, -Cys-X-Cys- (X is an amino acid other than cysteine) with adjacent serines, which appear to be the primary metal-binding sites; these occur in identical locations in the two variants. Most of the amino acid substitutions in MT are due to single base changes outside of the -Cys-X-Cys- regions (Nordberg & Kojima, 1979).

The human and rat liver MTs display the typical characteristics of mammalian MT: low molecular weight, no aromatic amino acids or histidine, a high A₂₅₀ indicative of Cd-sulfur interactions, and heat stability; however, a number of variations from this basic pattern do occur. The MT of newborn rat liver contains one phenylalanine residue which is not present in the adult form of MT (Waalkes et al., 1984c; Wong and Klassen, 1979). Rat testes contain three different classes of metal-binding proteins, TMBP-1, -2, and -3. TMBP-1 is comprised of the high molecular weight metal-binding proteins. TMBP-2 is the major heat-stable metal-binding protein with a

molecular weight of 25,000 dal. TMBP-3 appears to be a breakdown product of TMBP-2, resulting from either oxidation or proteolysis. TMBP-3 can be separated into two major forms, each containing very few cysteine residues and significant amounts of aromatic amino acids and histidine. Although the two forms of TMBP-3 have molecular weights of 8000 dal, they are very distinct from mammalian MT (Waalkes et al., 1984a; Waalkes et al., 1984b). The (Cd, Cu, Zn)-MT from carp has been purified and appears to have the same molecular weight and amino acid content as mammalian MT, including the presence of two isoforms of MT (Kito et al., 1982a). Upon exposure to Cd, marine invertebrates display an abundant amount of MT in the digestive gland called the hepatopancreas. The edible crab has three MT's, each with a different metal content but whose size and amino acid content are similar to that of mammals (Overnell, 1982). The mud crab has the two common isoforms of MT (Olafson et al., 1979) as does Drosophila (Debec et al., 1985). In contrast, oyster MT is very different from mammalian MT. It has a low molecular weight, but contains aromatic amino acids, a large number of dicarboxylic amino acids, and only 8% cysteine residues (Ridlington and Fowler, 1979).

There are two forms of Cd-binding proteins in Schizosaccharomyces pombe, Cd-BP1 and -BP2, and both are made up of a unit peptide, cadystin. Cd-BP1 has a

molecular weight of 4000 dal, while Cd-BP2 has a molecular weight of 1800 dal and they consist of 4 and 2 molecules of cadystin, respectively (Kondo et al., 1985; Murasugi et al., 1981). Cadystin actually exists in two forms, cadystin A and B. Cadystin A is a heptapeptide whereas cadystin B is a pentapeptide. Their amino acid sequences are (Glu-Cys-Glu-Cys-Glu-Cys-Gly) and (Glu-Cys-Glu-Cys-Gly). Cd-BP1 and -BP2 are remarkable in that, in spite of their very small size, they have a high capacity to sequester Cd.

The Cd-binding proteins (Cd-BP) from maize and the grass, Agrostis gigantea, are also smaller than mammalian MT, with molecular weights of 3100 and 3700 dal, respectively. The grass Cd-BP contains 29% cysteine and is heat-stable, whereas the maize Cd-BP contains 40% cysteine and is not heat-stable. Both contain relatively high concentrations of glutamic acid, not common in mammalian MT's (Rauser, 1984a; Rauser and Glover, 1984). The Cd-BP from the green alga Chlorella ellipsoidea has a molecular weight of 9000 dal and contains only 11% cysteine and greater than 50% acidic amino acids (Nayano et al., 1984).

The protist Tetrahymena synthesizes a single Cd-BP when grown in the presence of cadmium (Nakamura et al., 1981). Although it contains no aromatic amino acids and

has a high A_{250} due to Cd-sulfur coordination, it appears to be twice the size of mammalian MT and may have arisen from a duplication of the MT gene.

Hildebrand and Enger (Hildebrand et al., 1982) have done extensive work on the molecular biology of regulation of MT expression as it pertains to Cd detoxification. Using several Chinese Hamster Ovary (CHO) cell lines, including some selected on very high concentrations of Cd, they measured the amounts of MT, MT messenger RNA (mRNA), and the number of copies of Cd-responsive genes in response to Cd exposure. In the presence of Cd, Cd-resistant (Cd^R) CHO cells show an increase in the rate of MT synthesis and the more-resistant cell lines exhibit a more rapid response. Cd also induces the synthesis of 4 non-MT proteins ranging in size from 25,000 to 100,000 dal in all of the various Cd^R CHO cell lines. Actinomycin D completely inhibits the increased MT synthesis without a severe effect on general protein synthesis (Hildebrand and Enger, 1980); moreover, translatable MT mRNA appears slightly before MT and follows identical kinetics, suggesting that the transcription of MT mRNA is required for the induction of MT. Using a presumptive MT cDNA probe, a 2,000-fold increase in induction-specific RNA sequences was observed upon exposure of a Cd^R cell line to Cd.

Palmiter's and Brinster's laboratories have recently defined a 12-base-pair (bp) conserved sequence in the mouse MT-I gene promoter which, when placed in the upstream regulatory region, confers metal regulation upon a heterologous gene (Stuart et al., 1984). This metal regulatory element may bind a regulatory protein that is instrumental in the mediation of transcription of the downstream gene. Hamer, defining similar sequences, found that a distal 5' sequence plays a more general role in transcription of the MT gene whereas a proximal sequence is more necessary for metal-induced expression (Carter et al., 1984). Utilizing the mouse MT-I regulatory region as part of a fusion gene, he used competition assays to define MT-specific cellular factors which positively regulate MT-gene expression in the presence of Cd. These factors appear to interact with the DNA at the conserved regulatory elements (Séguin et al., 1984).

Glucocorticoids (GC), responsible for the synthesis of several stress-related proteins, also regulate the expression of MT. Dexamethasone, a synthetic GC, increases the transcription of the mouse MT-I gene approximately 90-fold (Hager and Palmiter, 1981). The 5' upstream sequences required to confer dexamethasone induction of transcription are in the region of approximately -245 to -265 and this site appears to be a

DNA binding site for the GC hormone-receptor complex (Karin et al., 1984).

It is interesting to note that stressful stimuli appear to have a universal effect on protein synthesis in all organisms ranging from bacteria to mammals and even plants. The response involves a diminution of general cytoplasmic protein synthesis with a concomitant increase in the synthesis of a few specific proteins termed "heat shock proteins" (HSP), or stress proteins. In Drosophila melanogaster, the 7 major HSPs vary in size from 22,000 to 83,000 dal (Asnburn and Bonner, 1979). Immunoassays using antibodies to 3 chicken embryo fibroblast HSPs demonstrated that many other organisms, including Drosophila, Xenopus, mouse, and yeast, produce cross-reactive proteins of similar molecular weights upon exposure to heat stress. It appears that the universality of the response to heat stress includes an evolutionary conservation of the HSPs (Kelly and Schlesinger, 1982). Stressful stimuli other than heat have a similar characteristic effect on the protein synthesis of cells. It is speculated that treatments inducing this response cause the accumulation of denatured or damaged proteins within the cell (Munro and Pelham, 1985; Ananthan et al., 1986). Arsenite, Cd^{++} , and Zn^{++} cause the enhanced synthesis of major stress proteins in avian red blood cells, some with molecular weights similar to those of the

larger HSPs (Atkinson and Dean, 1985); Cu^{++} , Cd^{++} , Zn^{++} , Hg^{++} , arsenite, chelating drugs, heat shock, and amino acid analogues all induce similar stress-related proteins in chick embryo fibroblasts (Levinson et al., 1980).

Cultured rat epithelial cells exposed to lead exhibit a specific induction of proteins, particularly those involved in the intranuclear inclusion bodies, which may protect the cells from harmful levels of the lethal metal (Shelton et al., 1986). Neurospora crassa conidia respond to heat shock in a manner similar to Drosophila, synthesizing major HSPs of 98,000, 83,000, and 67,000 dal (Plesofsky-Vig and Brambl, 1985); however, it is not known if Neurospora cells will synthesize similar "stress proteins" in response to the toxic metal, cadmium.

The selective amplification of the number of copies of a particular gene is a common adaptive mechanism shown to increase the cell's resistance to a number of toxic agents. It is doubtful that the phenomenon of gene amplification is caused by the toxic compounds. Those cells in which gene amplification has taken place, usually obtained after stepwise increases in the concentration of the toxic substance in the medium, simply survive and are selected for over those which have not undergone gene amplification or acquired another means of resistance.

Schimke has done extensive research into the mechanisms of methotrexate-(MTX-) resistance in murine cell lines. He has found both stable and unstable MTX-resistant cell lines which exhibit increased cellular levels of dihydrofolate reductase (DHFR) and DHFR mRNA, accompanied by proportional increases in the number of DHFR gene copies (as many as 200 times the number in MTX-sensitive cells) (Alt et al., 1978). The unstable resistant cell lines show the presence of double-minute (DM) chromosomes (small, self-replicating, extrachromosomal pieces of DNA lacking centromeres). Since they are lost in parallel with the loss of MTX resistance, the DMs are suggested to contain the amplified DHFR genes (Brown et al., 1981). In the stably amplified cell lines, the DHFR genes are integrated into the genome (Kaufman et al., 1981).

Palmiter has shown that gene amplification can contribute to MT overproduction (Beach and Palmiter, 1981). Cd^R Friend leukemia cells resistant to 12 times the concentration of Cd toxic to Cd^S cells accumulated 14 times more MT-I mRNA than Cd^S cells and have 6 times more MT-I genes. Note that this resistant cell line is approximately tetraploid and contains some very small chromosomes (not identified as DMs). CHO cells, resistant to a 120-fold greater Cd concentration and 4-fold Zn concentration than wild type cells, accumulated 200 times

more MT-1, had more than twice the MT-I mRNA induction capacity, were tetraploid, and showed 40- to 50-fold amplification of sequences homologous to MT-I cDNA (Gick and McCarty, 1981).

An example of gene amplification in plants as an adaptive mechanism even exists; one of the genes for glutamine synthetase (GS) exhibits an increase in copy number (Donn et al., 1984). GS, an enzyme essential for ammonia assimilation and the regulation of nitrogen metabolism, is inhibited by L-phosphinothricin (L-PPT); an L-PPT-resistant alfalfa cell line produces approximately 6 times more GS, 8 times more GS mRNA, and has a 5- to 11-fold increase in GS gene copy number.

An exciting aspect of the phenomenon of gene amplification as a method to overcome toxicity is the occurrence of co-amplification of a tightly-linked gene. N-(phosphonacetyl)-L-aspartate- (PALA)-resistant syrian hamster cells exhibit amplification of the CAD gene at the resident locus, adjacent to a nucleolus organizer. In two of the PALA-resistant mutants analyzed, tightly linked rDNA genes are co-amplified with the CAD genes (Wahl et al., 1983). A model has been proposed for the amplification of the CAD genes by unequal sister chromatid exchange facilitated by two flanking rDNA genes.

Copper-binding proteins (Cu-BP) have been studied in rats (Winge et al., 1975), grass (Rauser, 1984b), spinach (Tukendorf et al., 1984), yeast (Karin et al., 1984b), and Neurospora crassa (Lerch, 1980). The Cu-BP in rat liver is induced in vivo after Cu administration and is under transcriptional control as is the case with Cd-MT (Bremner and Davies, 1974; Premakumar et al., 1975); however it is distinct from Cd-MT and is therefore called copper chelatin. This metal-binding protein contains only 15% cysteine, has several aromatic amino acids, and has a different molecular weight than the Cd-MT. Rauser (1984b) characterized a Cu-BP in the grass Agrostis gigantea which is rapidly produced in plants exposed to copper. Two Cu-BPs with molecular weights of 12,500 and 9500 dal exist in spinach root, and a third with a molecular weight of 8500 dal is present in the leaves. These proteins appear to alleviate the toxic effects that Cu^{++} has on photosynthesis in plants grown in soil containing high levels of copper. The yeast Cu-thionein protein is very well characterized. In 1951, Ashida studied Cu-resistance in Saccharomyces ellipsoideus. The Cu^R cells appeared as brown colonies in the presence of Cu and Ashida proposed that an "effective principle", exhibiting the characteristics of RNA, diffused from Cu^R cells and enhanced the survival of Cu^S cells grown on copper. A two-step training process was required to obtain these

mutants (Minagawa et al., 1951). A sensitive strain was mated to a resistant strain and each ascus scored yielded 2 sensitive and 2 resistant progeny. In crosses homozygous for either trait, only progeny resembling the parental trait were obtained (Brenes-Pomales et al., 1955). Thus, the character conferring Cu-resistance is due to a single gene, known as the CUP1 locus. Later, a Cu-thionein, molecular weight of 9500 dal, was isolated from a brown Cu-resistant strain of S. cerevesiae. This thionein has a high cysteine content, no aromatic amino acids, but 2 to 3 times more acidic amino acids than mammalian Cd-MT (Prinz and Weser, 1975; Naiki and Yamagata, 1976). Weser purified this Cu-thionein from yeast grown under anaerobic conditions. He suggested the possibility that these Cu-sulfur proteins may serve some alternate function in the electron transport system. Similarly, Porter (1974) purified an insoluble Cu-thionein from newborn mammalian liver (neonatal hepatic mitochondriocuprein) which had characteristics very similar to equine Cd-MT; it was located in the heavy mitochondria/lysosome fraction, and was suggested to serve as a Cu storage reservoir for hepatic cytochrome oxidase during the neonatal developmental period.

Recently, this inducible Cu-thionein, also called copper chelatin, was analyzed at the molecular level. The gene encodes a protein of 61 amino acids, 20% of which are

cysteine (compared to 32% in mammalian Cd-MT). The protein contains more acidic amino acids than mammalian MT, but the -Cys-X-Cys- regions, and therefore the metal-thiolate binding clusters, along with a common 6-nucleotide sequence, are conserved. Unusually, this thionein contains two aromatic amino acid residues and one histidine residue (Butt et al., 1984). The induction of this Cu-BP appears to also occur at the level of transcription (Premakumar et al., 1974).

Welch and Fogel were first to isolate the CUP1^R gene (Fogel and Welch, 1982) and discovered that the CUP1^R locus is actually a tandemly iterated sequence containing 10-15 repeat units of the "sensitive allele" (Karin et al., 1984b). Upon examining several commercial yeast strains originally isolated from nature, they discovered that the different strains exhibited resistance to Cu over a 20-fold concentration range, and strains with higher degrees of Cu-resistance possessed a proportionally higher gene copy number. Some of the strains even exhibited different sizes of repeat units or different numbers of iterated sequences in the genome (Welch et al., 1983).

Hamer analyzed the physiological role of this yeast Cu-thionein. Using the one-step gene replacement method, he substituted a plasmid-borne URA3 gene flanked by CUP1 sequences for the chromosomal CUP1 gene via homologous

recombination, resulting in a cup1^Δ yeast strain lacking the CUP1 gene. In comparison, cup1^Δ, cup1^S, and CUP1^R all showed normal growth on standard medium, but the cup1^Δ was hypersensitive to Cu--it could not even grow on levels of Cu that the cup1^S strain could tolerate. Since its growth rate was normal and it could exist in the diploid state yielding viable progeny, the Cu-thionein protein must not be essential for normal growth and development. These experiments show that the Cu-thionein serves to protect yeast cells against Cu toxicity and that other cellular factors must be responsible for Cu transport, storage, and enzyme activation in its absence (Hamer et al., 1985).

A Cu-binding protein from Neurospora crassa has been completely characterized, rendering some remarkable parallels with the mammalian Cd-MT. Wild type cells grown on a very low concentration of Cu produce a heat-stable low molecular weight protein capable of binding copper. This protein contains only 7 different amino acids in a total of 25, of which 28% are cysteine; it has no aromatic amino acids. Upon comparison with equine kidney, human liver, and mouse liver (Cd, Zn)-MT, all 7 of the N. crassa cysteinyl residues occur in the same position as the first 7 cysteinyl residues at the N-terminal end of the vertebrate MT, as -Cys-X-Cys- clusters. The protein binds 6 g atoms of Cu per mole and has a molecular weight of

2600, including the Cu. This striking similarity must indicate an evolutionary relationship between these metallothioneins (Lerch, 1980). In reconstitution studies, the in vitro reconstitution of Neurospora tyrosinase is faster and more complete in the presence of Cu-MT than Cu^{++} alone, suggesting that Cu-MT may function in the transfer of metals to apometalloproteins (Beltramini and Lerch, 1982).

Studies involving the selection of mutants resistant to various chemical agents are very useful in that they not only provide additional genetic markers, but they also can provide insight into the genes responsible for enzymes of metabolic pathways as well as their regulatory genes.

N. crassa mutants resistant to the antibiotic actidione (cycloheximide, or CHX), have been selected (Hsu, 1962; Hsu, 1963; Howe and Terry, 1962). These mutants map to at least 4 independent loci and each involves the mutation of a single Mendelian factor. Two of the well-characterized mutants act-1 and act-2, are dominant over their corresponding wild-type alleles in heterokaryons. The double mutant, act-1;act-2 grows very slowly on minimal medium, whereas the act-1+act-2 heterokaryon grows normally. This signifies that at least one wild-type allele at either locus (in a single nucleus) is necessary for normal growth on minimal medium. The double mutant, however, is resistant to as much as ten

times the CHX concentration as either single mutant (Hsu, 1963). Later investigations showed that CHX affects protein synthesis on 80S ribosomes. Using N. crassa cell-free hybrid translation systems consisting of various combinations of CHX-resistant and CHX-sensitive ribosomes and supernatant fractions, inhibition of translation by CHX was found to depend only upon the origin of the ribosomal fraction. Three genes, act-1, act-2, and act-5 each determine ribosomal elements that can be modified by mutation to produce CHX resistance (Vomvayani, 1974).

Incubation of ultraviolet (UV) light-treated wild-type conidia without CHX for 13 hours yielded a 100-fold increase in the recovery of resistant mutants selected on CHX (Neuhaüser et al., 1970). Since CHX-resistant ribosomes in a cell-free translation system demonstrate a similar lag period before they overcome CHX inhibition, this lag may be necessary to synthesize the resistant ribosomes (Pongratz and Klingmüller, 1973). It is not yet known whether these mutations affect the rRNA or r-proteins or even whether the mutations are in the structural or regulatory genes.

CHX-resistant S. pombe mutants at different loci exhibit resistance to different levels of the antibiotic: some mutants only have low levels of resistance and others have high levels of resistance (Ibrahim and Coddington,

1976). Two of the low-level CHX-resistant mutants are also cross-resistant to anisomycin and trichodermin, both of which affect translation on 80S ribosomes. These mutations may affect permeability of the antibiotic, hence the low level of resistance and cross-resistances. Cold-sensitive mutants occur at each of the CHX-resistant loci, which may specify defective r-proteins.

The elucidation of the regulation of nitrogen catabolic pathways in N. crassa and Aspergillus nidulans was greatly facilitated by the study of mutants. While ammonia, glutamic acid and glutamine are the favored nitrogen sources for these fungi, secondary sources including nitrate and nitrite are also acceptable. In order to utilize such secondary sources, nitrogen catabolite repression must be lifted and the synthesis of the enzymes specific to the particular catabolite must be induced.

Mutants with the inability to use nitrate as a nitrogen source fall into several categories: mutations within (1) the structural genes for either nitrate reductase or nitrite reductase, the enzymes which convert nitrate to nitrite and nitrite to the preferred source, ammonia, (2) the genes which are responsible for the synthesis and assembly of the molybdenum-containing cofactor of nitrate reductase, (3) the positive-acting regulatory gene for nitrate-specific induction, and (4)

the positive-acting regulatory gene which is responsible for lifting nitrogen catabolite repression.

The selection of chlorate-resistant isolates of N. crassa and A. nidulans allowed for the accumulation of mutants with altered nitrate assimilation (Tomsett and Cove, 1979; Tomsett and Garrett, 1980). It was initially believed that in A. nidulans, chlorate was converted to the toxic form, chlorite, via nitrate reductase in chlorate-sensitive strains (Cove, 1976a; Cove, 1976b). However, current investigations revealed that not all nitrate reductase mutants are chlorate-resistant; the catalytic activity of nitrate reductase is not involved in chlorate-resistance. An explanation for the toxicity of chlorate might be that chlorate mimics nitrate and inhibits or represses some nitrogen catabolism via both nitrate reductase and the induction-specific regulatory gene product such that nitrogen deficiency results.

In a similar fashion, mutants with altered sulfate assimilation were selected on the toxic analogue, chromate (Marzluf, 1970). In order for sulfate to be utilized as a sulfur source, both induction and de-repression of a family of related sulfatases and permeases must take place. The chromate-resistant N. crassa mutants had a reduced capacity for sulfate transport and involved mutations at two different loci: one governing the

induction of the genes necessary for sulfate assimilation and the other encoding the structural gene for the conidial sulfate permease. In the absence of the sulfate permease the toxic chromate was excluded, hence the resistant phenotype.

More fungal mutants with resistance to various other toxic compounds have been isolated, yet their resistance mechanisms are probably less well understood. Ethionine-resistant mutants of the basidiomycete Coprinus lagopus have been isolated and mapped near two genes responsible for methionine synthesis (Lewis, 1963). Ethionine, a toxic analogue of methionine, may not exert its toxic effect solely via its incorporation into proteins. One ethionine-resistant mutant does not incorporate ethionine into protein and its methionine-activating enzyme does not activate ethionine. However, another ethionine-resistant mutant incorporates almost twice as much ethionine into protein as the ethionine-sensitive strain does. It is presumed that the mutation conferring ethionine-resistance lies in the gene for the methionine-activating enzyme; this enzyme may serve the dual function of activating methionine and transmethylation of important methylated compounds. Ethionine toxicity would then result from the ethylation of precursors of methylated compounds, and a mutation which blocks transethylation via the methionine-activating enzyme could thus be responsible for reducing

the toxic effects of ethionine.

Fungi resistant to antibacterial and antifungal agents including acriflavin (Roper and Käfer, 1957; Hsu, 1965; Howe and Terry, 1962), teoquil, malachite green, iodoacetate, p-fluorophenylalanine (Warr and Roper, 1965), isoniazid, sodium benzoate (Howe and Terry, 1962), and others have been studied in less detail, and only a few of the mutants have been genetically mapped.

Another interesting category of resistant mutants in N. crassa is that of the os mutants. The os mutants are inhibited by high osmolalities of growth medium and are classified by their sensitivity to the addition of 4% sodium chloride to the medium (Mays, 1969). However these, and the similar smco mutants, are also resistant to both dicarboximide and aromatic hydrocarbon fungicides. The osmotic sensitivity could be due to alterations in the cell wall/plasma membrane complex which would not allow the cells to adapt to the mechanical stress caused by changes in the solute concentration of the fungal cytoplasm or external environment. The cell walls of os mutants are thinner and more irregular than those of wild type, causing the os mutants to look like collapsed or twisted tubes as opposed to regular cylinders (Trevithick and Metzenberg, 1966). In wild-type, cell wall pore sizes appear to vary over a continuous range from small to large

pores; os mutants appear to have many more large pores than wild type and this trait, along with the thinner cell wall, may affect the kinetic sieving of macromolecules. Other cell wall-specific changes in os mutants include a drop in the amount of glucose in the alkali-insoluble portion of the cell wall and a change in the amount of galactosamine (Livingston, 1969).

Mutants have been selected both on the basis of their osmotic sensitivity and their fungicide resistance. Many os mutants are resistant to fungicides, and vin mutants (resistant to the dicarboximide fungicide vinclozolin, VIN) are sensitive to high osmotic pressure. VIN-resistance appears to be controlled by several genes, one of which is either allelic with or tightly linked to the os-1 locus. The fungicide resistance may be caused by a disturbance in the normal balance between cell wall lysis and the synthesis of new cell wall material or by an inability to maintain turgor pressure due to an observed alteration in carbohydrate metabolism (Grindle, 1984; Grindle and Temple, 1982; Grindle and Temple, 1983).

At this point, several mechanisms for the relief of metal toxicity must be discussed. The most prevalent mechanism is the sequestering of cadmium by metallothionein in such a fashion that the cytoplasmic environment is protected from the presence of the cadmium ions. The transcription of the MT gene is induced by the

presence of Cd so that the MT is produced upon entrance of Cd into the cell. In several cases, cells can tolerate very large concentrations of Cd because of amplification of the MT gene. This process usually occurs when cells are subjected to stepwise increases in Cd concentration.

In the case of several strains of Staphylococcus aureus, neither of these mechanisms appears to be responsible for Cd-resistance. Many strains of S. aureus bearing antibiotic-resistant plasmids also exhibit Cd-resistance. It was discovered that the resistant strains manifest their Cd tolerance by taking up approximately 20-fold less Cd than the sensitive strains (Tynecka et al., 1981). The Cd-sensitive strains take up Cd from the environment via a high affinity, energy-dependent manganese (Mn) transport system meant for the uptake of physiological cations. At low Cd concentrations, the resistant strains take up negligible amounts of Cd via this Mn-transport system. At higher concentrations, what little Cd enters the cells is extruded through what is hypothesized to be an energy-dependent Cd^{++} efflux system which exchanges cellular Cd^{++} for protons. This Cd^{++} /proton antiporter appears to be encoded by the antibiotic-resistance plasmid harbored in the Cd-resistant S. aureus strains.

Other resistance mechanisms may include exclusion of Cd from the cell via alterations in the cell wall/plasma membrane complex as seen in the os mutants of N. crassa, or perhaps by vesicularization, compartmentation, or precipitation of the cadmium ions, protecting the cytoplasmic components from their presence.

In the yeast Cryptococcus albidus, cells exposed to the fungistatic compound silver nitrate showed two distinct ultrastructural differences when compared to cells grown in the absence of silver nitrate (Brown and Smith, 1976). First, electron micrographs revealed the presence of electron-opaque deposits within the vacuole, as was confirmed by X-ray microanalysis. Second, ring- or disc-shaped electron-opaque areas were detected in the cell wall of treated cells. The silver ions may bind to the cell wall, resulting in the disc-shaped granules, and then enter the cytoplasm after which they are packaged into the vacoules.

The research reported in this dissertation involved the isolation and characterization of cadmium-resistant mutants of Neurospora crassa. One of the main reasons that Neurospora is an ideal organism for scientific investigation is that it has been studied for many years and a considerable amount of information about it is available. Beginning with Beadle and Tatum's first nutritional mutants (Tatum and Beadle, 1942), Neurospora

has proven itself advantageous for genetics and molecular biology.

Neurospora has a rapid life cycle and is easily grown on a simple, chemically-defined medium. It has several well-defined developmental stages, including vegetative and aerial hyphae, the asexual macroconidia and microconidia, and the sexual ascospores. The mycelial organization is coenocytic, such that the hyphae are compartmentalized by porous septa through which nuclei can pass, and each compartment typically contains several nuclei. All vegetative cells exist in the haploid state but heterokaryons can be formed to study the interaction of homologous genes. Crosses are easily made and its genetic behavior is similar to that of higher plants and animals, which makes Neurospora an attractive eukaryotic system. One valuable characteristic of Neurospora is that the progeny of a genetic cross can be analyzed either as random meiotic products or as ordered tetrads, which provide evidence of the events that happened during meiosis. There is a large number of mapped and well-characterized mutants for each of Neurospora's seven chromosomes, thus making the system suitable for genetic studies.

The investigation reported in this dissertation involved the isolation and characterization of cadmium-resistant mutants of N. crassa. The goal of this study

was to identify a set of genes which were able to confer a Cd^R phenotype, with the possibility of obtaining amplification of a Cd-metallothionein gene.

Cadmium-resistant mutants were obtained by several techniques and four Cd^R mutants, as well as a naturally-occurring Cd^R isolate from Japan, reside at three different genetic loci: one on Linkage Group (LG) II, one on LG VI, and a third at an unmapped but distinct locus. The growth of these mutants under different conditions was compared to that of wild-type: the mutants apparently have altered morphologies, possibly involving the cell wall or cell membrane. Transport of cadmium into Neurospora cells involves more than a single uptake system, but it is the same for the Cd^S wild-type and the Cd^R mutants; therefore these mutations do not appear to alter membrane transport of cadmium. No increase in cadmium-binding potential was evident for either Cd^R mutant, and resistance is probably not conferred by an increase in heat-stable cadmium-binding proteins. My work demonstrated that cadmium does not induce the heat shock/stress response in Neurospora, but it does produce a definite alteration in the protein profile of both the wild-type and the Cd^R mutants (this alteration may be a new cadmium-induced stress response). It is unclear whether these changes are actually the property that confers the resistance phenotype or are simply a result of growth in the presence of a toxic metal ion.

MATERIALS AND METHODS

Strains

The cadmium-sensitive Neurospora crassa wild-type strain used in this work was 74-OR23-1A and the cadmium-resistant strains were derived from this. All other strains were obtained from the Fungal Genetics Stock Center, University of Kansas Medical School, Kansas City, Kansas, and are listed in Table 1. Cultures were routinely grown on Fries' Minimal Medium (Ryan et al., 1943) containing 1.5-2.0% sucrose, biotin, and any necessary supplements. (This medium will hereafter be referred to as FMM).

UV Mutagenesis

In a dimly-lit room, a 300 ml conidial suspension containing 10^9 conidia in a 1 l beaker was irradiated with a germicidal short wave UV light for 40 min. During the irradiation the suspension was gently stirred. This duration of UV light exposure yielded approximately a 90% kill. To test the percentage of surviving conidia, aliquots of the conidial suspension were removed at various time points, diluted, and plated on FMM containing

Table 1 Neurospora Strains

Genotype	Mating Type	FGSC ^a
74-OR23-1	A	987
74-OR23-1	a	988
<u>N. sitophila</u> , Obama, Kyushu	A	1764
<u>N. sitophila</u> , Obama, Kyushu	a	1765
<u>N. sitophila</u> 173.25	a	2538
<u>N. sitophila</u> NRRL 2884	a	3299
<u>N. intermedia</u> Unzen, Kyushu	A	1768
WT Lein 7A	A	847
WT Lindegren 1A	A	354
WT Costa Rica	A	851
WT Groveland-1c	a	1945
WT Houma-1n	A	2220
WT Liberia	A	961
WT Mauriceville-1c	A	2225
WT Puerto Rico 18	a	429
WT Roanoke-1m	A	2227
WT Tatum	a	-- ^b
WT Emerson 5297	a	352
WT St. Lawrence STA	A	262
WT <u>N. Sitophila</u>	A	2216
<u>alcoy</u>	A	997
<u>alcoy</u>	a	998
<u>arg5</u>	A	3589
<u>arg12</u>	A	3595
<u>aur;pe</u>	A	1203
<u>fluffy</u>	A	1838
<u>fluffy</u>	a	1690
<u>nic3,met7</u>	A	152

^aFungal Genetics Stock Center Number.^bNot obtained from Fungal Genetics Stock Center.

1.2% sorbose and 0.1% sucrose as colonizing factors. The plates were incubated at 30°C for at least 48 hours in the dark to prevent photo repair of genetic lesions, then for 24-48 hours in the light, after which the surviving colonies were counted. For the isolation of mutants, aliquots of the irradiated conidia were spread at a density of approximately 0.5×10^7 per plate (15 x 100 mm) onto FMM containing 1.2% sorbose, 0.1% sucrose, and 10, 15, or 20 mM CdCl₂ (filter-sterile CdCl₂ was added to autoclaved medium). After 2 days of growth in the dark and 6 days growth in the light at 30°C, Cd-resistant colonies were picked and transferred to growth rate plates.

Cadmium-induced Mutagenesis

Utilizing the mutagenic effects of cadmium, 1×10^7 and 0.5×10^7 wild-type conidia were spread, without any additional treatment, onto plates containing FMM, 1.2% sorbose, 0.1% sucrose, and 4-10 mM CdCl₂. Three plates per concentration of CdCl₂ per plating density were incubated at 30°C for 8 days, at which time small foci of dense resistant colonies appeared on a background of light mycelial growth. These foci were picked and transferred to growth rate plates.

Training Flasks

Training flasks were begun by inoculating Erlenmeyer flasks with wild-type conidia onto solid medium containing agar or into standing liquid medium. All cultures were serially transferred by way of a small inoculum to fresh medium every 10 days. The five series of training flasks employed were as follows: (1) maintained on 5 mM CdCl_2 , (2) initially inoculated on 5 mM CdCl_2 , then transferred to and maintained on 10 mM CdCl_2 , (3) initially inoculated on 5 mM CdCl_2 , transferred to 10 mM CdCl_2 , then transferred to 20 mM CdCl_2 after 6 weeks, (4) maintained on 10 mM CdCl_2 , and (5) initially inoculated on 10 mM CdCl_2 , then transferred to and maintained on 20 mM CdCl_2 . After 4 months of growth on the maintenance levels of CdCl_2 , all concentrations were increased by 2 mM and then by another 1 mM CdCl_2 after 4 additional months of growth. When a trained culture began to show an adaptive response in the presence of cadmium (i.e. aerial hyphae and conidia, similar to the growth of wild-type in the absence of cadmium), a small inoculum was transferred to a flask containing a higher concentration of CdCl_2 in attempts to increase the level of cadmium-resistance.

Growth Rate Plates

Sterile pasteur pipettes were used to remove "plugs" from individual Neurospora colonies growing in Petri plates containing FMM. The plug inoculum was transferred to a "growth rate plate" containing FMM and CdCl_2 ranging from 0-20 mM (no sorbose). The mycelial plug was placed at the edge of the agar medium in the Petri dish and the concentric rings of mycelial growth were marked directly on the dish every 24 hours. The average growth in cm/day was calculated for five plates at each concentration of CdCl_2 . It was determined that 15 mM CdCl_2 was the minimum level needed to restrict wild-type growth; thus, the amount of CdCl_2 used in plating experiments was 15 mM or higher. Only those mutants which exhibited a growth rate on cadmium medium greater than 2 times that of wild-type were selected for further investigation.

As an additional test, Cd^R mutants that revealed high growth rates on the growth rate plates were tested in liquid medium containing CdCl_2 . This test involved inoculating 20 conidia (determined by absorbance at 660 nm) into a test tube containing 1 ml of sterile FMM plus 6, 8, or 10 mM CdCl_2 , 6mM CdCl_2 being the minimal level required to restrict wild-type growth in liquid medium.

Death vs. Inhibition

Sterile pieces of dialysis tubing cut to the size of a Petri dish were layered on top of agar medium consisting of FMM, 1.2% sorbose, 0.1% sucrose, and 5, 10, 15, or 20 mM CdCl_2 . An inoculum of 5×10^6 conidia was carefully spread on top of the dialysis tubing. After 11 days of growth at 30 C, the dialysis tubing was transferred to plates without CdCl_2 and the extent of subsequent growth was recorded.

Crosses

Crosses were made by inoculating slants of Westergaard's medium (Westergaard and Mitchell, 1947) with either a small conidial or mycelial inoculum of each parent at opposite ends of the slant. Occasionally crosses required that conidia of the male parent be spread over the surface of a growing female mycelial culture. The slants were incubated in the dark at 25°C and transferred to the light after the appearance of perithecia. The ascospores were heat shocked at 60°C for 45-60 minutes and spread on plates containing FMM, 1.2% sorbose, and 0.1% sucrose.

The mating types of the progeny were determined by inoculating a conidial suspension of each isolate onto a plate containing Westergaard's medium, covered with a "lawn" of fluffy (fl) mycelia (both mating types, A and a,

were tested). Fluffy is a macroconidia-less mutant used routinely for determining mating type. The plates were incubated at 25°C in the dark until perithecia were evident on either of the two plates and mating type was scored.

Cross Analysis Plates

Pasteur pipette plugs of the germinated ascospore colonies were transferred to plates containing three different cadmium concentrations: FMM plus 0.8% sorbose, 0.4% sucrose, and 15, 25, or 30 mM CdCl₂. Resistant progeny were usually able to grow on all three concentrations of CdCl₂, whereas wild-type growth was inhibited by all three levels of cadmium. Plates I and II show examples of cross analysis plates, demonstrating both cadmium-sensitive and -resistant isolates grown in the absence and presences of cadmium.

Growth Rate Flasks

To determine the effects that cadmium has on N. crassa in liquid medium, Erlenmeyer flasks (125 ml) containing 20 ml of FMM and 0, 0.5, 2.0, or 10 mM CdCl₂ were inoculated with conidia to a final absorbance at 420 nm of 0.35. The cultures were grown at 30°C, shaking at 200 RPM. The mycelia were harvested by suction filtration

Plate I. Cadmium-sensitive and cadmium-resistant isolates grown in the absence and presence of cadmium. An example of the pasteur pipette plug method for determining cadmium-resistance.

Plate I.a. Cadmium-sensitive and cadmium-resistant isolates grown in the absence of cadmium.

Plate I.b. Cadmium-sensitive and cadmium-resistant isolates grown in the presence of 25 mM CdCl_2 .

B

A

Plate I

Plate II. Cadmium-sensitive and cadmium-resistant isolates grown in the presence of 15, 25, or 30 mM CdCl_2 .

on tared Whatman No. 1 filter paper, washed well with distilled water, dried in a 45°C oven for at least 20 hours, and weighed. For comparative purposes, growth rate flasks were inoculated with cadmium-sensitive or cadmium-resistant conidia and harvested at various time points during a 45-50 hour period. Duplicate or triplicate flasks were inoculated for each different time point during that period and treated as above.

Cross-Resistance

Cross-resistance to other heavy metals was tested using the same plating methods as for the cross analysis plates. The medium contained 35 and 40 mM ZnCl_2 , 35 and 40mM $\text{Co}(\text{NO}_3)_2$, and 20 and 25 mM CuSO_4 . These levels were determined to be the minimum inhibitory concentrations for each metal.

Heat Shock/Cadmium Shock

Using the method of Plesofsky-Vig (Plesofsky-Vig and Brambl, 1985), wild type conidia were suspended in 30 ml of FMM plus 2% sucrose to a final density of 1 mg/ml; 25 μCi [^{35}S]methionine (1132 Ci/mole; ICN) were added and the flasks were shaken in a water bath at 180 RPM at either 30 or 45°C. After 30 minutes the flasks were removed, 0.5 μmoles of non-radioactive methionine were added, and the flasks were chilled on ice to stop [^{35}S]methionine

incorporation. The conidia were harvested by filtration on a 1.2um Millipore filter and suspended in 10 ml of homogenization buffer consisting of 62.5 mM Tris-HCl (pH 6.7), 2% sodium dodecyl sulfate (SDS), 0.75 mM phenylmethylsulfonyl fluoride (PMSF), and 7.5% 2-mercaptoethanol. The suspension was homogenized in a Braun homogenizer with 25 g of glass beads for two 30 second intervals. The homogenized extracts were heated in a boiling water bath for 3 minutes and centrifuged at 12,000 x g for 10 minutes. For Cd shock, the conidia were treated the same way except for the addition of either 1 mM or 10 mM CdCl₂ at 30°C. The extracts were electrophoresed on a 10% polyacrylamide gel with 1 x 10⁵ cpm of each extract loaded per lane.

Transport Studies

The transport of cadmium was determined by incubating either conidia or 300 ug of mycelia in FMM containing 1.5% sucrose and CdCl₂. For conidial transport, the conidial suspension was filtered and inoculated to a final OD at 420 nm of 1.0. For mycelial transport, a filtered conidial suspension was inoculated into FMM plus 1.5% sucrose to a final OD at 420 nm of 0.1 and grown, shaking at 200 RPM, for 13 hours at 25°C. At zero time ¹⁰⁹CdCl₂ (NEN) was added to 0.9 ml aliquots of the 13 hour mycelial

suspension to a final concentration of 0.5 mM (1 mCi/mmol) unless otherwise noted; the 1 ml samples were incubated in a 25°C shaking water bath at 120 RPM. At various time intervals the samples were collected on glass fiber filters, washed well with a solution of non-radioactive Cd, dried, and counted in a Beckman LS 7000 Liquid Scintillation Spectrophotometer using Formula-963 scintillation cocktail (NEN). For assays to determine the energy-dependence of Cd transport, mycelia were pre-incubated with 10^{-4} M sodium azide 15 minutes prior to the addition of $^{109}\text{CdCl}_2$. Competitive metal ions were added at equimolar concentrations of Cd unless otherwise noted and all were the divalent chloride salt form.

Leucine transport was assayed using conditions as described above, with 0.1 mM [^3H]leucine (50 mCi/mmol; ICN). The amount of leucine incorporated into protein was determined by stopping identical samples with 4 ml of 12.5% trichloroacetic acid (TCA), heating the samples for 30 minutes at 80°C, and collecting the precipitate on glass fiber filters followed by thorough washing with 5% TCA. The filters were dried and counted as above. To assay transport under conditions in which protein biosynthesis was inhibited, 5×10^{-4} M cycloheximide was added to the mycelia 5 minutes prior to the addition of Cd or leucine.

For Cd efflux studies, ^{109}Cd was added to 10 ml of 13 hour mycelia and transport was allowed to continue for 90 minutes. After the cells were pre-loaded with ^{109}Cd , the mycelia were harvested under low pressure onto Whatman No. 1 filter paper, washed, and resuspended in 10 ml of various media without ^{109}Cd . At timed intervals 1 ml samples were removed, filtered, washed, and counted as described above.

Cadmium/Hemoglobin Assay

Preparation of homogenates. Lyophilized mycelia of various strains, grown in the presence or absence of 1 mM CdCl_2 , were ground by mortar and pestle with glass beads (4 times the dry weight of mycelia). The powdered mycelial samples were suspended in 20 volumes of 10 mM Tris-HCl buffer, pH 7.4, and vortexed for 1 minute. Each homogenate was centrifuged at $10,000 \times g$ for 10 minutes and heated in a boiling water bath for 2 minutes, after which the centrifugation step was repeated.

Assay (Eaton and Toal, 1982). A ^{109}Cd solution was made by combining carrier-free $^{109}\text{CdCl}_2$ with CdCl_2 in 10 mM Tris-HCl buffer, pH 7.4, to yield a final concentration of 2 $\mu\text{g Cd/ml}$ with a specific radioactivity of 1 $\mu\text{Ci/ml}$. A 200 μl aliquot of each homogenate was incubated with 200 μl of the ^{109}Cd solution for 10 minutes. After the incubation, 100 μl of a 2% bovine hemoglobin solution was

added to each sample and mixed. The samples were heated in a boiling water bath for 2 minutes and another 100 μ l of 2% hemoglobin was added. The samples were cooled and centrifuged at 15,600 $\times$ g in an Eppendorf centrifuge for 5 minutes, and 500 μ l of the supernatant was removed; 20 μ l of each supernatant fluid was counted as described above.

Radioactive Labelling of Proteins In Vivo

Erlenmeyer flasks (250 ml) containing 50 ml of FMM and 250 μ Ci [35 S]sulfur as Na_2SO_4 were inoculated with conidia to an absorbance at 420 nm of 0.35. The mycelia were grown, shaking at 200 RPM, for 25 hours at 30°C. Labelling was performed either in the presence or absence of 1 mM CdCl_2 . Mycelia were harvested on Miracloth, frozen in liquid nitrogen, lyophilized, and stored at -70°C.

Portions of the dried mycelia were ground by mortar and pestle with glass beads (4 times the dry weight of mycelia) and suspended by extensive vortexing in 20 volumes of 20 mM sodium phosphate (pH 7.2), 5 mM 2-mercaptoethanol, and 0.5 mM benzamidinium HCl. The homogenate was centrifuged at 13,000 RPM for 20 minutes at 4°C. The supernatant was removed and spun in a Beckman L8-70 Ultracentrifuge at 35,000 RPM for 1 hour at 4°C. The supernatant was filtered through a Millex-GV 0.22 μ m Filter Unit and stored at -20°C.

Protein Quantification

Protein was measured using the Bio-Rad Protein Assay kit with bovine gamma globulin as the standard. A standard curve was performed with each assay.

Electrophoresis

Heat shock/cadmium shock proteins were separated on a discontinuous 10% polyacrylamide gel using the electrophoresis system of Laemmli (Laemmli, 1970) as modified by O'Farrell (O'Farrell, 1975). Briefly, the stacking gel consisted of 4.5% acrylamide, 0.125 M Tris-HCl (pH 6.8) and 0.1% SDS. The resolving gel consisted of 10% acrylamide, 0.375 M Tris-HCl (pH 8.8), and 0.1% SDS. The reservoir buffer contained 0.0125 M Tris-HCl, 0.1 M glycine, and 0.05% SDS. The samples were mixed in a 1:1 ratio with sample buffer, consisting of 0.0625 M Tris-HCl, 2% SDS, 10% glycerol, and 5% 2-mercaptoethanol. Electrophoresis was performed at 30 mA, constant current, until the dye front reached the bottom of the gel. The gel was stained in a solution containing 0.1% Coomassie Brilliant Blue (R-250) dissolved in a 5:1:5 mixture of methanol, acetic acid, and water and destained in the same solution lacking the dye.

Crude protein homogenates from in vivo labelling experiments were separated on a 5-20% acrylamide gradient

gel using a modification of the Shapiro system (Shapiro, 1967). The gel consisted of 0.1% sodium phosphate (pH 7.2), 6 M urea, and 0.1% SDS. The reservoir buffer contained 0.1 M sodium phosphate (pH 7.2), 0.1% SDS, and 1 mM EDTA (to prevent electroplating of the cathode). The homogenates were mixed in a 1:1 ratio with sample buffer, consisting of 0.01 M sodium phosphate (pH 7.2), 2% SDS, 6 M urea, 2% 2-mercaptoethanol, 30% glycerol, and 0.02% bromophenol blue. The gel was run at 50 mA, constant current, and treated as described above.

Molecular weights were estimated by the coelectrophoresis of the molecular weight standards (Sigma Chemical Company) ovalbumin, 45,000, and lysozyme, 14,300. The [^{14}C]-labelled protein molecular weight standards (Bethesda Research Laboratories) included: bovine trypsin inhibitor, 6,200; lysozyme, 14,300; β -lactoglobulin, 18,400; α -chymotrypsinogen, 25,700; ovalbumin, 43,000.

Fluorography

Gels were soaked in 5 times the volume of Enlightning (NEN) for 15-30 minutes to introduce fluors into the gel. The gels were then dried with a Bio-Rad Model 224 Slab Gel Dryer, placed next to a piece of Kodak X-Omat AR film, and stored at -70°C .

RESULTS

Isolation of Cadmium-resistant Mutants

To determine the minimum level of CdCl_2 required to inhibit the growth of the N. crassa wild-type strain, inocula were placed at the perimeter of agar medium containing 0-20 mM CdCl_2 in Petri dishes. Daily linear growth rates were recorded by marking the concentric rings of growth away from the point of inoculation. The daily growth was determined for five plates for each concentration of CdCl_2 and used to calculate an average daily growth rate (in cm/day) for a 6-day growth period (Figure 1). From this figure, 15 mM CdCl_2 was determined to be the level of CdCl_2 required to restrict wild-type growth.

Three different types of mutagenic techniques were employed to obtain Cd^R mutants: UV light irradiation, single step selection by Cd mutagenesis, and multiple step selection by Cd mutagenesis using training flasks.

Selection of mutants resistant to CdCl_2 required that a large number of conidia be irradiated by UV light. Figure 2 shows the relationship between length of UV light

Figure 1. Growth rate of N. crassa wild-type strain on various concentrations of CdCl_2 . "Plug" inocula were placed at the perimeter of agar medium in Petri dishes containing 0-20 mM CdCl_2 . Daily growth rates were calculated for five plates for each concentration and used to calculate an average daily growth rate (in cm/day) for a 6-day growth period.

Figure 1

Figure 2. UV mutagenesis survival curve. A 300 ml suspension containing 10^9 wild-type conidia was irradiated with UV light for 60 min. At various time points, an aliquot of the conidial suspension was removed and plated in triplicate on FMM containing 1.2% sorbose and 0.1% sucrose as colonizing factors. The plates were incubated for 48 hr in the dark, then 24-48 hr in the light (30°C), after which time the surviving colonies were counted. Values are presented as the log of the number of surviving colonies.

Figure 2

exposure and the number of surviving conidia when 1×10^9 spores were used. A length of UV light exposure was used that killed approximately 90% of the spores, in order to obtain a high yield of genetic lesions in the survivors. For UV mutagenesis experiments, 1×10^9 conidia were exposed to UV light for 40 minutes, and Cd^R mutants were selected on 10, 15, and 20 mM CdCl_2 . The 43 resistant mutants which appeared after 8 days were picked and transferred to growth rate plates (see Materials and Methods section) containing 18 mM CdCl_2 . Those mutants which showed growth rates greater than 2 times that of wild-type on solid medium containing cadmium were tested for growth in liquid medium containing inhibitory levels of CdCl_2 .

To see if Cd^R mutants could be generated solely by the mutagenic effect of CdCl_2 , 1×10^7 and 0.5×10^7 wild-type conidia were inoculated onto plates containing various concentrations of cadmium. After 8 days, small foci of resistant growth appeared upon a background of very light growth. The foci were counted and a Cd-induced mutation frequency was calculated (Table 2). The foci were picked, transferred to growth rate plates, and manipulated in the same manner as the UV light-induced Cd^R mutants.

Training flasks are a classical method used for the development of cell lines resistant to toxic agents. By

Table 2 Cadmium-induced Mutation Frequency on Various Concentrations of CdCl₂

Concentration CdCl ₂ (mM)	Spontaneous Mutation Frequency	
	Experiment 1	Experiment 2
4	8.20×10^{-6}	5.27×10^{-6}
5	3.53×10^{-6}	2.70×10^{-6}
6	5.40×10^{-6}	2.13×10^{-6}
7	2.67×10^{-6}	2.37×10^{-6}
8	3.73×10^{-6}	1.70×10^{-6}
9	2.33×10^{-6}	1.43×10^{-6}
10	0.73×10^{-6}	1.13×10^{-6}

Note. 0.5×10^7 and 1.0×10^7 conidia were plated in Experiment 1 and Experiment 2, respectively. Experimental values represent an average of three plates per concentration of CdCl₂ per plating density.

culturing cells in a medium containing a suitable concentration of the toxic agent, a certain percentage of the cells either have or can acquire the ability to adapt to the environment, and this adaptive response can be continued on higher levels of the toxicant. This "training" can take place in a natural situation as well as in the laboratory. Resistant strains obtained by training, however, may or may not maintain their resistance when cultured without the toxic agent used for training.

Several series of training flasks were used in this study, each cultured on increasingly higher concentrations of CdCl_2 . At one point in the stepwise increases of CdCl_2 concentration, a particular training flask culture began to conidiate. This isolate had been initially inoculated on 5 mM CdCl_2 , and then was transferred to and maintained on 10 mM CdCl_2 . Approximately 4 months after the cadmium concentration was increased from 10 mM to 12 mM, the growth and conidiation of this training flask mutant in the presence of CdCl_2 resembled that of the wild-type strain grown in the absence of cadmium. At this time, all training flasks were recultured on medium containing an amount of CdCl_2 that was 1 mM higher than that previously used.

One month after the conidiating training flask culture had adjusted to 13 mM CdCl_2 , it was cultured on

medium lacking CdCl_2 . This culture was used for subsequent growth tests and genetic crosses. One interesting feature of this training flask culture (in the absence of cadmium) was that the aerial hyphae were "seepy": rather than appearing as dry hyphal filaments covered with conidia, there were dark, wet trickles of material oozing down the hyphal surface. A flask of this mutant growing in the presence of 13 mM CdCl_2 was compared to a culture of wild-type growing in the presence and absence of 13 mM CdCl_2 (Plate III). The wild-type appearance of this training flask mutant was short-lived, and eventually it took on the appearance of a smooth layer of mycelia on the surface of the agar.

An experiment was performed to understand the effect that cadmium has on the Cd-sensitive cells: whether growth is merely inhibited in the presence of cadmium or if cell viability is actually lost. Wild-type Cd^S conidia were plated on media containing low, medium, or high levels of CdCl_2 . They were, however, plated on top of a piece of dialysis tubing to allow transfer to medium without cadmium. After nine days of incubation on CdCl_2 , all plates showed colonies of resistant growth; the plates with higher concentrations of cadmium had fewer resistant colonies. After two more days of growth, the dialysis tubing was transferred to plates containing medium lacking cadmium. One day following the transfer the resistant

Plate III. Comparison between the wild-type strain
and Training Flask mutant grown in the presence of
13 mM CdCl_2 .

colonies grew larger and after two additional days, the colonies grew even larger and only a few additional colonies appeared. These additional colonies were probably just too small to score initially. Given several more days of growth, colony size increased and conidiation took place. Upon microscopic examination, it was obvious that no background growth had occurred, even after transfer to medium without the presence of the toxic metal ions. These results show that the resistant colonies appeared on a background of inviable cells. Apparently the Cd^S conidia were unable to germinate or grow in the presence of cadmium and death ensued.

Comparative growth rates of wild-type and the Cd^R mutants, including UV-induced mutants, single step Cd -induced mutants, and the training flask mutant, were used to identify the most resistant isolates. These growth rates are listed in Tables 3 and 4. The mutants with growth rates significantly higher than wild-type were then tested in liquid medium containing $CdCl_2$ and these results are listed in Table 5. The results from these two tests were the basis for selecting certain mutants for further analysis. The 10 mutants in Table 5 which were able to grow in liquid medium containing inhibitory levels of $CdCl_2$ were chosen for additional study.

Table 3 Growth Rates of UV-induced Cd-resistant Mutants

Strain	mM CdCl ₂ ^a	Growth Rate (cm/day)
74-OR23-1	--b	0.180
10-1	10	0.137
10-2	10	0.242
10-3	10	0.139
10-4	10	0.220
15-1	15	0.181
15-2	15	0.243
15-3	15	0.146
15-4	15	0.402
15-6	15	0.231
15-7	15	0.199
15-8	15	0.323
15-10	15	0.275
20-1	20	0.210
20-2	20	0.180
20-3	20	0.185
20-4	20	0.245
20-5	20	0.230
20-6	20	0.246
20-7	20	0.276
20-9	20	0.215
20-10	20	0.371
20-12	20	0.411
20-13	20	0.257
20-14	20	0.198
20-16	20	0.321
20-17	20	0.233
20-19	20	0.283
20-20	20	0.212

^aConcentration of CdCl₂ used in selection of mutants.^bCd-sensitive control.

Table 4 Growth Rates of Cd-induced Cd-resistant Mutants

Strain	mM CdCl ₂ ^a	Growth Rate (cm/day)
74-OR23-1	- ^b	0.180
4A2	4	0.473
4E2	4	0.264
5C1	5	0.224
5C2	5	0.215
5D3	5	0.169
5D4	5	0.243
5E2	5	0.233
6B2	6	0.269
6C2	6	0.248
6D2	6	0.301
6D3	6	0.244
6E1	6	0.260
7A2	7	0.270
8B2	8	0.237
8C2	8	0.392
8D2	8	0.264
8D6	8	0.298
8E1	8	0.263
8E2	8	0.300
9A2	9	0.306
9D1	9	0.249
9E1	9	0.283
9E4	9	0.280
10A2	10	0.257
10C1	10	0.242
10D4	10	0.263
TF	13	0.464

^aConcentration of CdCl₂ used in selection of mutants.^bCd-sensitive control.

Table 5 Growth of Specific Cd-resistant Mutants in Liquid Selective Medium

Strain	Mutagen	mM CdCl ₂ ^a	Growth ^b
74-QR23-1	-- ^c	--	-
10-2	UV	10	+/-
15-2	UV	15	+
15-4	UV	15	+
15-10	UV	15	++
20-7	UV	20	+
20-10	UV	20	-
20-12	UV	20	-
20-16	UV	20	-
20-19	UV	20	+
4A2	Cd	4	N.T. ^d
4E2	Cd	4	+/-
6D2	Cd	6	+
8C2	Cd	8	+
8D2	Cd	8	+/-
8E2	Cd	8	+
9A2	Cd	9	+
TF	Cd	13	++

^aConcentration of CdCl₂ used in selection of mutants.

^bVisible growth was ranked as "-", +/-, +, or ++, with "-" signifying no visible growth and "++" signifying profuse growth.

^cCd-sensitive control.

^dNot tested.

Genetic Analysis of Mutants

The next step of the analysis was to determine which mutants exhibited a resistance trait that was transmitted through meiosis. Each Cd^R mutant was crossed to the sensitive strain 74-OR23-1A; the resulting ascospore progeny were germinated on FMM and the resistance of each isolate was tested on medium containing CdCl_2 . The results of these crosses are listed in Table 6. Of the ten Cd^R strains examined, only five transmitted their resistant character through meiosis. These isolates are the UV mutants 20-19 and 15-10, the Cd-induced mutants 9A2 and 4A2, and the training flask mutant. At this point it was determined that the UV mutant 20-19 did not show a clear difference from wild-type in resistance to CdCl_2 ; since it was very slow-growing and not significantly resistant, it was dropped from the study.

When crossed to the wild-type sensitive strain, the TF mutant not only segregated Cd^R and Cd^S progeny, but on sorbose medium it also segregated large and small colonies. The small colonies were all Cd^S and resembled wild-type in size. The large colonies were both Cd^R and Cd^S . Although these large colonies varied in size on medium containing sorbose, they were all larger than wild-type and displayed very diffuse growth, rather than exhibiting a discrete colonial morphology. This mutant may contain some membrane alterations that cause the loss

Table 6 Percentage of Cd^R and Cd^S Progeny

65

Cd ^R Parent	Progeny		
	Number Examined ^a	%Cd ^R	%Cd ^S
15-4		0	100
20-7		0	100
6D2		0	100
8C2		0	100
8E2		0	100
15-10	26	4	96
20-19	36	28	72
4A2	62	53	47
9A2	26	38	62
TF	33	38	62

Note. Crosses involved each Cd^R mutant crossed to the Cd^S wild-type strain.

^aNumber Examined in blank columns was greater than 25.

of the well-defined colonial appearance on sorbose medium.

The mutant 4A2 also segregated large and small colonies when crossed to the wild-type sensitive strain; however in this case the large colonies were all Cd^R . They were much larger than the wild-type colony size on sorbose medium and were very distinctly colonial rather than showing the diffuse colonial morphology of the TF mutant. The small colonies were both Cd^R and Cd^S . The Cd^R colonies varied in size, but were generally larger than wild-type colonies, whereas the Cd^S colonies very closely resembled wild-type colonies. These morphological differences will be discussed in more detail later.

Once the reliable Cd^R mutants were selected for study, it was necessary to determine the number of different complementation groups that were represented and the relative map location of each. A convenient genetic manipulation exists for N. crassa whereby a mutant of unknown location is crossed to alcoy, a multiple translocation tester strain which contains markers for six of the seven linkage groups (LG). Each of the three translocations is represented by an easily scored morphological marker at or close to the interchange point: albino conidia (al- 1; LG I and II), temperature-sensitive colonial growth (cot; LG IV and V), and yellow conidia (ylo; LG III and VI). Linkage of the Cd^R trait to any of

these markers would suggest that it is located on either of the two linkage groups involved in the translocation. Otherwise, the resistance-conferring gene is presumably associated with the seventh linkage group. Upon determination of the linkage group(s) involved, crosses to several follow-up marker strains can be used to localize the gene.

One resistant reisolate of each Cd^R mutant was crossed to alcoy. Information regarding these crosses is found in Table 7. Mutant Cd^R 9A2R5 a segregates 50:50 with the cot and ylo markers, but a strong linkage to the al marker was indicated by its low (10%) recombination frequency. A follow-up cross to mutants with markers on LG I and II was required to further localize this Cd^R mutant.

The results of the cross of the Cd^R mutant TFR38 A to alcoy a show that this mutant is not linked to the ylo marker, and because the recombination frequencies for the cot and al markers are both near 30%, a cross to a marker on LG VII might more clearly show a definite linkage.

Linkage groups for the other two Cd^R mutants were not as clearly defined. When crossed to alcoy, the Cd^R mutant 15-10R22 yielded certain classes of progeny that appeared to have very low viability. For example, of the 99 isolates tested, 70 were Cd^S and only 29 were Cd^R. In spite of this problem linkage is not indicated to the cot

Table 7 Assignment of Linkage Group for Cd^R Mutants 9A2R5,
15-10R22, 4A2R1, and TFR38

Cross	Progeny Phenotype					R.F. ^d
	m ^a (LG) ^b	Cd ⁺ m ⁺ ^c	Cd ⁺ m ⁻	Cd ⁻ m ⁺	Cd ⁻ m ⁻	
9A2R5 <u>a</u> x <u>Alcoy A</u>	<u>cot</u> (IV,V)	33	24	18	22	57
	<u>al</u> (I,II)	0	57	30	10	10
	<u>ylo</u> (III,VI)	0	0	14	16	53
15-10R22 <u>a</u> x <u>Alcoy A</u>	<u>cot</u> (IV,V)	38	32	24	5	43
	<u>al</u> (I,II)	18	52	27	2	20
	<u>ylo</u> (III,VI)	5	13	20	7	27
4A2R1 <u>A</u> x <u>Alcoy a</u>	<u>cot</u> (IV,V)	17	21	16	27	54
	<u>al</u> (I,II)	6	32	30	13	23
	<u>ylo</u> (III,VI)	- ^e	-	-	-	-
TFR38 <u>A</u> x <u>Alcoy a</u>	<u>cot</u> (IV,V)	10	49	11	17	31
	<u>al</u> (I,II)	13	46	15	13	30
	<u>ylo</u> (III,VI)	6	7	4	11	61

^aMarker evaluated.

^dRecombination Frequency.

^bLinkage Group.

^eNot determinable.

^cCd⁺ = Cd^S; Cd⁻ = Cd^R.

marker, nor did the differences in viability seem to affect the segregation of the ylo marker. The two parental classes, $Cd^S\text{al}^-$ and $Cd^R\text{al}^+$, are present in nearly equal numbers, but of the recombinant classes, there are 18 $Cd^S\text{al}^+$ progeny and only 2 $Cd^R\text{al}^-$ progeny. In this instance, the Cd^R gene could be located on any linkage group other than LG IV and V (represented by the cot marker).

Similar mapping difficulties were experienced with the Cd^R mutant 4A2R1 A. This mutant did not conidiate very well and therefore the distinction between orange and yellow conidia was not easily made. In this case, linkage was not indicated to the cot marker (LG IV and V), but linkage to the al marker was difficult to determine. Again, follow-up crosses had to be performed for each linkage group other than LG IV and V.

After the required additional crosses were completed for each Cd^R mutant, a definite linkage was finally reached for each of them. Results from the successful follow-up crosses are listed in Tables 8-10. Again, certain classes of recombinants appear to have poor viability, therefore the recombination frequencies should be regarded as approximate. Recombination between the Cd^R gene and the nic3 marker is roughly 10-13% for 15-10R22 and TFR38, and slightly higher for 4A2R1 (18%). The

Table 8 Assignment of Map Location for Cd^R Mutants 15-10R22 and TFR38

Cross	Progeny Phenotype									R.F. ^c
	m ₁	m ₂	m ₃ ^a	m ₁ ⁺ m ₂ ⁺ ^b	m ₁ ⁺ m ₂ ⁻	m ₁ ⁻ m ₂ ⁺	m ₁ ⁻ m ₂ ⁻	m ₁ ⁺ m ₂ ⁺ m ₃ ⁺	m ₁ ⁻ m ₂ ⁻ m ₃ ⁻	
15-10R22 <u>a</u>	Cd	<u>nic3</u>		8	27	28	0			13
x										
<u>nic3,met7</u> <u>A</u>	Cd	<u>met7</u>		11	24	28	0			17
	<u>nic3</u>	<u>met7</u>		32	4	7	20			17
	<u>nic3</u>	Cd	<u>met7</u>					4	0	30
TFR38 <u>A</u>	Cd	<u>nic3</u>		6	25	31	0			10
x										
<u>nic3,met7</u> <u>a</u>	Cd	<u>met7</u>		14	17	31	0			23
	<u>nic3</u>	<u>met7</u>		34	3	11	14			23
	<u>nic3</u>	Cd	<u>met7</u>					2	0	29

^aMarker evaluated.

^cRecombination Frequency.

^bCd⁺ = Cd^S; Cd⁻ = Cd^R.

Table 9 Assignment of Map Location for Cd^R Mutant 4A2R1

Cross	Progeny Phenotype									R.F. ^c
	m ₁	m ₂	m ₃ ^a	m ₁ ⁺ m ₂ ⁺ ^b	m ₁ ⁺ m ₂ ⁻	m ₁ ⁻ m ₂ ⁺	m ₁ ⁻ m ₂ ⁻	m ₁ ⁺ m ₂ ⁺ m ₃ ⁺	m ₁ ⁻ m ₂ ⁻ m ₃ ⁻	
4A2R1 <u>Δ</u> x <u>nic3,met7 a</u>	Cd	<u>nic3</u>		10	23	22	0			18
	Cd	<u>met7</u>		8	25	22	0			15
	<u>nic3</u>	<u>met7</u>		22	10	8	15			33
	<u>nic3</u>	Cd	<u>met7</u>					0	0	33

^aMarker evaluated.

^bCd⁺ = Cd^S ; Cd⁻ = Cd^R.

^cRecombination Frequency.

Table 10 Assignment of Map Location for Cd^R Mutant 9A2R5

Cross	m ₁	m ₂ ^a (LG) ^b	Progeny Phenotype				R.F. ^d
			m ₁ ⁺ m ₂ ⁺ ^c	m ₁ ⁺ m ₂ ⁻	m ₁ ⁻ m ₂ ⁺	m ₁ ⁻ m ₂ ⁻	
9A2R5 <u>a</u>	Cd	<u>aur</u> (I)	9	10	21	22	50
x							
<u>aur;pe</u> <u>A</u>	Cd	<u>pe</u> (II)	0	9	19	2	7
9A2R5 <u>a</u>	Cd	<u>arg</u> (II)	9	32	49	8	17
x							
<u>arg5</u> <u>A</u>							
9A2R5 <u>a</u>	Cd	<u>arg</u> (II)	23	29	43	3	35 ^e
x							
<u>arg12</u> <u>A</u>							

^aMarker evaluated.^bLinkage Group.^cCd⁺ = Cd^S ; Cd⁻ = Cd^R.^dRecombination Frequency.^eDetermined using arg⁺ progeny, only.

recombination frequency between the Cd^R gene and the met7 marker ranged from 15-23% for these mutants, but taking into consideration any double recombinational events, all distances add up reasonably well: The mutant gene for the three independent Cd^R isolates designated 15-10R22, TFR38, and 4A2R1, all appear to fall approximately midway between the nic3 and met7 markers on LG VII (Table 9).

The Cd^R mutation 9A2 is on LG II, as shown by the cross to aur;pe A (Table 10). Because there are very few useful mutants located on LG II, it was not possible to carry out a three point cross to more accurately establish the mutation's position. Instead, mutant 9A2 was crossed separately to two different arg mutants: arg5, located near the centromere, and arg12, on the right arm of LG II. The germination of arg⁻ progeny from the cross of 9A2 to arg12 was very low (32 arg⁻ to 66 arg⁺), and therefore the recombination frequency was calculated using the arg⁺ progeny only. Taking the arg5 and arg12 data together with the known map distance between these two loci (approximately 15 map units), the Cd locus must be to the left of the arg5 marker. It therefore appears that there are at least two loci which are able to confer Cd -resistance upon N. Crassa: one locus on the right arm of LG II and one locus on the left arm of LG VII.

To support this assignment, two of the mutants which had been localized to L.G. VII, TFR38 and 15-10R22, were

crossed to each other; as expected, the progeny were 100% Cd^R (Table 11). Because both parental mutations mapped to the same location, recombination would not be expected to occur between the two mutant genes and no sensitive progeny should be produced. In contrast, when the mutant from L.G. II, 9A2R5, was crossed to either of the mutants on L.G. VII, the progeny segregated in a $Cd^S:Cd^R$ ratio of 1:3. In this case, one parent had the sensitive allele for the LG II gene and the resistant allele for the LG VII gene, whereas the other parent had the reverse situation. Thus, the outcome would yield 75% of the progeny carrying at least one of the resistant alleles (thus conferring a resistant phenotype) and 25% of the progeny carrying both sensitive alleles. These results therefore strongly support the above-mentioned conclusion of two distinct loci for cadmium resistance.

Analysis of Mutants Under Various Growth Conditions

It is well known that in some cases, resistance to an inhibitor is lost when the cells are not cultured on the selective agent. This can occur by a recombinational loss of the gene(s) which confers the resistance, or a segregational loss of extrachromosomal elements carrying those genes. To assess the stability of my Cd^R strains, each was maintained on non-selective medium

Table 11 Evaluation of Progeny from Crosses Between Cd-resistant Mutants

Parents (L.G.) ^a	Progeny	
	Number Evaluated	Cd ^S :Cd ^R
TFR38 <u>A</u> (VII) x 15-10R22 <u>a</u> (VII)	54	0:4
TFR40 <u>a</u> (VII) x 9A2R4 <u>A</u> (II)	54	1:3
15-10R22 <u>a</u> (VII) x 9A2R4 <u>A</u> (II)	60	1:3

^aLinkage Group.

lacking cadmium for six serial transfers after which Cd-resistance was again tested. All mutants exhibited the same resistance to the high levels of cadmium which inhibit wild-type.

Fourteen different wild-type Neurospora strains, both laboratory strains and others isolated from around the world, were tested for Cd-resistance in liquid medium. None of the 14 tested was able to grow in the presence of the heavy metal. Due to the abundance of cadmium in Japan's environment, it was interesting to see whether any wild-type strains isolated from different areas of Japan had developed Cd-resistance from growing in a cadmium-rich environment. The N. sitophila strains from Obama were isolated from a small nonindustrial resort town near the ocean--probably not polluted. The N. intermedia strain from Unzen was isolated from a national park in an area full of volcanic outcroppings, hot springs, and sulfurous vents. There was no industrial pollution in the area, but the area may be naturally rich in heavy metals (D. Perkins, personal communication). This N. intermedia strain was actually resistant to high levels of cadmium. When crossed to the Cd^S wild-type strain 74-OR23-1a, the progeny were 63% Cd^S and 37% Cd^R (number examined was 59). Crosses of N. crassa to N. intermedia are very fertile, but so few viable ascospores are produced that genetic analyses are difficult. However, this naturally-resistant

wild-type strain did segregate Cd^S progeny when crossed to the Cd^R mutants 9A2R5 and 15-10R22. Apparently there is a distinct gene in this wild-type N. sitophila strain, unlinked to either of my two mapped mutations, which confers the Cd-resistance phenotype.

In addition to resistance to cadmium, the mutants were tested for resistance to zinc, copper, and cobalt. Since metallothionein has the potential to alleviate the toxicity of heavy metals by binding to them and because it can be induced by metals other than cadmium (Zn, Cu, Ag, and Au), the cross-resistance of each of the Cd^R mutants was tested on concentrations of $ZnCl_2$, $CuSO_4$, and $Co(NO_3)_2$ that were inhibitory to wild-type growth. None of the Cd^R mutants was able to grow in the presence of Zn, Cu, or Co. Likewise, the Japanese wild-types were tested for cross-resistance to these other heavy metals and they, too, were unable to grow.

Because there was a possibility that any of these Cd^R mutants might be due to a phenotype similar to the osmotic mutants, each one was tested for growth in liquid medium containing 4 M NaCl. Each of the mutants was able to grow in the presence of high osmotic pressure, and therefore an os⁻ phenotype is not involved.

The growth rates of wild-type and three of the Cd^R mutants were compared over a 45-50 hour period in liquid

medium containing cadmium. Figure 3 shows the growth of wild-type on 0, 0.5, 2.0, and 10 mM CdCl_2 . Cadmium appears to have two noticeable effects on wild-type: the most noticeable effect is that the total mass of mycelia produced is reduced in the presence of cadmium (Figure 4). As the concentration of cadmium increases, the total growth mass decreases. The other effect of cadmium is the appearance of a lag period before growth occurs, especially visible with 10 mM CdCl_2 .

It is interesting to note that cadmium appears to have a slightly different effect on each of the individual mutants (Figures 5-8). First of all, none is able to grow as well as wild-type in shaking liquid medium containing cadmium; they were all selected as surpassing wild-type growth on solid medium. This must have some relationship to the ability of the aerial hyphae to grow away from the substratum containing the cadmium source (possible in both selection techniques: agar medium and standing liquid culture), whereas in liquid medium the mycelia are constantly immersed in the cadmium source. The TF mutant is able to grow better than the other Cd^R mutants (Figure 8), especially at the higher concentrations of cadmium. It was surprising that the TF mutant should have such a different behavior than mutant 15-10 in the presence of 10 mM CdCl_2 , since both involve a mutation at the same locus on LG VII. The mutant 9A2 is

Figure 3. Growth of wild-type strain on various concentrations of CdCl_2 . Wild-type conidia were added to a series of identical flasks containing FMM and 0, 0.5, 2.0, or 10 mM CdCl_2 . Triplicate cultures were grown at 30°C , shaking at 200 RPM, and removed at various time points. Mycelia were harvested, washed, and dried in a 45°C oven. Values are presented as the dry weight of mycelia from each flask at each point.

Figure 3

Figure 4. Percent of control growth for the wild-type and the Cd^R mutants TFR38, 9A2R5, and 15-10R22 on various concentrations of CdCl₂. Triplicate flasks containing FMM and 0, 0.5, 2.0, or 10 mM CdCl₂ were inoculated to the same density with conidia of each strain and these cultures were grown for 20 hr. Mycelia were harvested, washed, dried, and weighed. Values are presented as the amount of mycelial growth at the different concentrations of cadmium as compared to the growth of each isolate on 0 mM CdCl₂. At each concentration, the isolates are (from left to right): wild-type, TFR38, 9A2R5, and 15-10R22.

Figure 4

Figure 5. Growth of Cd^R mutant TFR38 on various concentrations of CdCl₂. Triplicate flasks containing FMM and 0, 0.5, 2.0, or 10 mM CdCl₂ were inoculated and the cultures were grown for various time periods. Mycelia were harvested, washed, dried, and weighed. Values are presented as the dry weight of mycelia from each flask at each time point.

Figure 5

Figure 6. Growth of Cd^R mutant 9A2R5 on various concentrations of CdCl_2 . Duplicate flasks containing FMM and 0, 0.5, 2.0, or 10 mM CdCl_2 were inoculated and the cultures were grown for various time periods. Mycelia were harvested, washed, dried, and weighed. Values are presented as the dry weight of mycelia from each flask at each time point.

Figure 6

Figure 7. Growth of Cd^R mutant 15-10R22 on various concentrations of CdCl₂. Triplicate flasks containing FMM and 0, 0.5, 2.0, or 10 mM CdCl₂ were inoculated and the cultures were grown for various time periods. Mycelia were harvested, washed, dried, and weighed. Values are presented as the dry weight of mycelia from each flask at each time point.

Figure 7

Figure 8. Comparison of of wild-type and the Cd^R mutants' growth on various concentrations of CdCl₂. Duplicate or triplicate flasks containing FMM and 0, 0.5, 2.0, or 10 mM CdCl₂ were inoculated to the same density with conidia of each strain and the cultures were grown for 20 hr. Mycelia were harvested, washed, dried, and weighed. Values are presented as the percentage of growth of each isolate (at each concentration of CdCl₂) compared to the growth of wild-type at that same concentration of cadmium, with the amount of wild-type growth equaling 100%. At each concentration, the isolates are (from left to right): wild-type, TFR38, 9A2R5, and 15-10R22.

Figure 9

unable to tolerate high concentrations of CdCl_2 , but at the lower cadmium concentration, 0.5 mM, it grows as well as the other two mutants. In comparing the amount of dry weight produced with time, mutant 9A2 grows to the same extent in 2 mM CdCl_2 as it does in 0.5 mM CdCl_2 (in spite of an increased initial lag period), whereas the other two mutants, both at the other locus, show a marked decrease in mass at the higher concentration of CdCl_2 .

Cadmium Transport

The transport of cadmium into Neurospora cells was investigated, first comparing conidial transport to mycelial transport, and then examining various parameters of mycelial transport, including a comparison of cadmium transport between wild-type and the Cd^R mutants.

In the same period of time, mycelia transport almost 40 times more cadmium (per mg wet weight) into the cells than do conidia (Figures 9 and 10). At the end of a two hour transport period, conidia had transported 8.125×10^{-5} umoles Cd/mg wet weight, whereas mycelia had transported 3×10^{-3} umoles Cd/mg wet weight. This result seems reasonable because the conidia are in a metabolically inactive state as compared with mycelia. Conidia are in a semidormant state, whereas mycelia are metabolically active, with growth proceeding from the

Figure 9. Transport of cadmium into wild-type conidia. Wild-type conidia were added to FMM containing 0.1 mM CdCl_2 (1 mCi $^{109}\text{Cd}/\text{ml}$). Samples were removed at various time points, filtered, washed well, dried, and counted. Values represent the umoles of Cd (calculated from the counts/min) transported.

Figure 8

Figure 10. Transport of cadmium into wild-type mycelia. A 13 hr mycelial suspension was added to FMM containing 0.1 mM CdCl_2 (1 mCi ^{109}Cd /mmole). Samples were removed at various time points, filtered, washed well, dried, and counted. Values represent the umoles of Cd (calculated from the counts/min) transported.

Figure 10

hyphal tip.

To determine whether a previous exposure to cadmium induces a transport system for this metal ion, a mycelial suspension was incubated with 0.1 mM CdCl_2 for 20 minutes prior to the addition of ^{109}Cd ; the uptake of ^{109}Cd was compared to the control flask which received no preincubation with CdCl_2 . Over a three hour period, there was no difference in ^{109}Cd uptake between the cells which received a cadmium pretreatment and those which did not (data not shown). Cadmium therefore does not have an ability to induce its own uptake.

The uptake of ^{109}Cd was followed for five different concentrations of CdCl_2 over a 100-fold range, from 0.01 mM to 1.0 mM (Figure 11). Uptake was slow for concentrations ranging from 0.01 to 0.1 mM, and rose rapidly for 0.5 and 1.0 mM CdCl_2 . It appears that cadmium transport does not involve a single uptake system, since there is not a saturation effect upon increasing the concentration of the substrate (cadmium) when the amount of mycelia is held constant (Figure 12). A single straight line cannot be drawn between the points on the double reciprocal plot. The internal transport of cadmium might involve a multicomponent system, and free diffusion may even be involved.

In an effort to determine whether cadmium transport was an energy-dependent process, azide was used to

Figure 11. Transport of cadmium into wild-type mycelia for various concentrations of CdCl_2 . A 13 hr mycelial suspension was added to FMM containing 0.01, 0.05, 0.1, 0.5, or 1.0 mM CdCl_2 (1 mCi ^{109}Cd /mmole). Samples were removed at various time points, filtered, washed well, dried, and counted. Values represent the umoles of Cd (calculated from the counts/min) transported.

Figure 11

Figure 12. Effect of cadmium concentration on the rate of cadmium transport into mycelia. Data from Figure 11 were used to calculate the rate of cadmium transport into mycelia for various concentrations of CdCl_2 (Left) and for the double-reciprocal plot (Right).

Figure 12

inhibit ATP synthesis occurring during oxidative phosphorylation. Mycelial suspensions were incubated for 15 minutes with or without azide before the addition of ^{109}Cd (Figure 13). The cells which were treated with azide exhibited a 75% reduction in cadmium transport, suggesting that 75% of the cadmium transported is via energy-dependent processes and the remaining 25% does not require an energy input. If cells are allowed to accumulate ^{109}Cd for 40 or 60 minutes before azide is added, the amount of cadmium transported during the next 15 minutes (in the presence of azide) shows a similar sharp reduction (Figure 13).

It seemed possible that cadmium transport in Neurospora occurs by an uptake system which also transports other ions, for example there are instances whereby cadmium shares a transport system with manganese, calcium, or potassium. Cadmium transport was investigated in the presence of a variety of different potential competitive metal ions to determine the possible antagonistic or synergistic effects of each (Figure 14, 15). These include the Group II alkaline earth metals magnesium and calcium, and zinc, a Group IIb metal whose chemistry is similar to that of cadmium, as well as several metals from the first transition series, manganese, iron, cobalt, nickel, and copper. None of the

Figure 13. Effect of azide on transport of cadmium by wild-type mycelia. Closed circles: a 13 hr mycelial suspension was incubated in FMM containing 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole). Samples were removed at various time points, filtered, washed well, dried, and counted. Open circles: a 13 hr mycelial suspension was incubated in FMM containing 10^{-4} M sodium azide for 15 min. prior to the addition of the cadmium solution. At zero time, 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole) was added and transport was followed for the times indicated. Open circles from solid line; a 13 hr mycelial suspension was incubated in FMM containing 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole) for 40 or 60 min. Azide was added (final concentration 10^{-4} M) and cells were allowed to continue transport for 15 min. Cells were then harvested and treated as above.

Figure 13

Figure 14. Transport of cadmium into wild-type mycelia in the presence of magnesium, manganese, or zinc. A 13 hr mycelial suspension was added to FMM containing 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole) and 0.5 mM of each metal ion as indicated. Samples were removed at various time points, filtered, washed well, dried, and counted. Values represent the umoles of cadmium transported. Symbols: ●, CdCl_2 , only; □, + ZnCl_2 ; △, + MgCl_2 ; ○, + MnCl_2 .

Figure 14

Figure 15. Transport of cadmium into wild-type mycelia in the presence of cobalt, copper, iron, or nickel. A 13 hr mycelial suspension was added to FMM containing 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole) and 0.5 mM of each metal ion as indicated. Samples were removed at various time points, filtered, washed well, dried, and counted. Values represent the umoles of cadmium transported. Symbols: ●, CdCl_2 , only; □, + CoCl_2 ; △, + NiCl_2 ; ○, + FeCl_2 ; X, + CuCl_2 .

Figure 15

metals appear to have any competitive effects at equimolar concentrations, and when the concentration of manganese was increased 10-fold over the concentration of cadmium, the drop in the amount of cadmium transported is probably not significant (Figure 16). In the case of S. cerevisiae, the inhibition of cobalt or cadmium uptake was most severe by cations with similar ionic radii (Norris & Kelly). The ionic radius of calcium ($0.99 \text{ \AA}$) is the only one similar to cadmium ($0.97 \text{ \AA}$), with the other radii ranging from $0.72 \text{ \AA}$ (cobalt and copper) to $0.80 \text{ \AA}$ (manganese). However, even calcium did not compete with the transport of cadmium (data not shown). Copper, on the other hand, seemed to have a positive effect upon the uptake of cadmium, increasing its uptake approximately 11-fold (Figure 16). In the presence of an equimolar amount of copper, 220×10^{-5} umoles of ^{109}Cd were transported into the cells and in the absence of copper, only 20×10^{-5} umoles were transported. When the ratio of copper to cadmium was reduced from an equimolar ratio to 0.5 mM, the amount of ^{109}Cd transported returned to the control level.

It is not clear how copper enhances cadmium uptake, however the possibility exists that it may trigger the de novo synthesis of additional transporter molecules. To test this hypothesis, the transport of cadmium with and without the presence of an equimolar amount of copper was compared to (1) the transport of cadmium in the presence

Figure 16. Transport of cadmium into wild-type mycelia in the presence of copper or manganese. A 13 hr mycelial suspension was added to FMM containing 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole), ●, including 0.5 mM CuCl_2 , X, 0.05 mM CuCl_2 , ▲, or 5 mM MnCl_2 , ■. Samples were removed at various time points, filtered, washed well, dried, and counted. Values represent the umoles of cadmium transported.

Figure 16

of copper after a five minute preincubation with cycloheximide (CHX), and (2), the transport of cadmium after an initial five minute preincubation with CHX and then a 15 minute preincubation with copper (Figure 17). CHX has no effect on the increase of cadmium transport due to the presence of copper; therefore copper does not stimulate the de novo synthesis of any proteins involved in cadmium transport. To confirm that CHX did indeed inhibit any further synthesis of proteins, its effect on the incorporation of [^3H]leucine into proteins was examined (Figure 18). CHX had little effect on the amount of [^3H]leucine transported into the cells, but a five minute CHX preincubation caused a 96% inhibition of [^3H]leucine incorporation into TCA precipitable material.

Next, cadmium transport by wild-type was compared to one isolate of each representative Cd^{R} mutant: 15-10R22 and 9A2R5. Cadmium uptake (micromoles of cadmium transported per gram dry weight) occurred at comparable rates for wild-type and the two Cd^{R} mutants (Figure 19). In Figure 20, the wild-type and Cd^{R} mutants were compared with and without a preincubation with azide. Similar to wild-type, as shown in Figure 13, a portion of both of the Cd^{R} mutants' cadmium transport was inhibited in the presence of azide. The different systems responsible for cadmium transport must be very similar in the wild-type

Figure 17. The effect of cycloheximide (CHX) on the transport of cadmium into wild-type mycelia in the presence of copper. A 13 hr mycelial suspension was added to FMM containing 0.5 mM CdCl_2 (1 mCi ^{109}Cd /mmole), ●, or 0.5 mM CdCl_2 (same specific activity) plus 0.5 mM CuCl_2 , ○. Another portion of the 13 hr mycelial suspension was added to FMM containing 5×10^{-4} M CHX and incubated for 5 min, after which the $^{109}\text{CdCl}_2$ and CuCl_2 were added (same concentrations and specific activity as above), ▲. A sample of the 13 hr mycelial suspension was added to FMM containing 5×10^{-4} M CHX and incubated for 5 min. CuCl_2 was added to a final concentration of 0.5 mM and the mycelia incubated for another 15 min before the addition of the radioactive cadmium solution, X. Samples were removed at various time points and treated as described in the legend of Figure 10.

Figure 17

Figure 18. Transport of leucine into wild-type mycelia and the effect of CHX on leucine incorporation into proteins. A 13 hr mycelial suspension was added to FMM containing 0.1 mM leucine (120 Ci [^3H]leucine/mmol). At various time points, samples were removed, filtered, washed, dried, and counted for transport, ●. Parallel samples were heated in TCA, filtered, washed, dried, and counted for incorporation, ○. To determine the effects of CHX on transport and incorporation, mycelia were preincubated with 5×10^{-4} CHX for 5 min prior to the addition of [^3H]leucine and treated as above (■, no CHX; □, + CHX).

Figure 18

Figure 19. Transport of cadmium into wild-type and Cd^R mutant mycelia. A 13 hr mycelial suspension of either wild-type, Cd^R 9A2R5, or 15-10R22 was added to FMM containing 0.5 mM CdCl₂ (1 mCi ¹⁰⁹Cd/mmole) and treated as in the legend to Figure 10. Values represent the umoles of cadmium transported per gram of dry weight. Symbols: ●, wild-type; ▲, 9A2R5; ■, 15-10R22.

Figure 19

Figure 20. Effect of azide on cadmium transport into wild-type and Cd^R mutant mycelia. Transport was carried out as in Figure 13. Closed symbols represent transport in the absence of azide (●, wild-type; ▲, 9A2R5; ■, 15-10R22) and open symbols represent transport after a 15 min preincubation with 10⁻⁴ azide (○, wild-type; △, 9A2R5; □, 15-10R22).

Figure 20

(sensitive) and cadmium-resistant strains.

Cadmium efflux from cells was studied by allowing mycelia to accumulate ^{109}Cd for 90 minutes and then monitoring the cellular content present at various times after transfer into different types of media. The amount of ^{109}Cd efflux from the cells was the same after their transfer to medium lacking cadmium, medium containing the same amount of cadmium as used for transport, or medium containing 10 times the concentration as used for transport (Figure 21). Manganese had no effect on cadmium efflux, nor did the presence of azide (data not shown). Therefore, neither the energy state of the cell nor the extracellular metal concentration had any effect on the amount of cadmium that left the cell.

Cadmium-binding Potential

It was of interest to determine whether any of the Cd^R strains possessed larger amounts of Cd-binding proteins than present in the sensitive strain. Because many documented cases of metal ion-resistant strains (derived from initially-sensitive strains) are due to an increase in the amount of cytosolic metallothionein, the Cd/hemoglobin affinity assay of Eaton and Toal was used to quantify the levels of low-molecular-weight metal-binding proteins in the mycelia of wild-type and the Cd^R mutants. Although this method measures the cadmium-binding capacity

Figure 21. Efflux of cadmium in the presence of various concentrations of metals. A solution of CdCl_2 (final concentration 0.5 mM; 1 mCi ^{109}Cd /mmole) was added to a 13 hr myclial suspension and incubated for 90 min. An aliquot was removed, the cells were washed free of ^{109}Cd , and then transferred to either FMM, ●, 0.5 mM CdCl_2 , ■, 0.5 mM CdCl_2 and 0.5 mM MnCl_2 , X, or 0.5 mM CdCl_2 and 5 mM MnCl_2 , ▲. Samples were removed at various time points and treated as in Figure 10.

Figure 21

rather than the true amount of protein, the values from this assay, when applied to rat liver homogenates, were in close agreement with those obtained from ion exchange chromatography as measured by atomic absorption spectroscopy. The basis of this assay involves the interaction of the heat stable proteins with ^{109}Cd . Any ^{109}Cd which remains free in solution is bound by the added hemoglobin and subsequently removed from the solution by the heat denaturation of the hemoglobin.

The linearity of this assay for the *N. crassa* system was determined using several dilutions of wild-type mycelial homogenate. The results from mycelia which had not been exposed to CdCl_2 deviated from linearity to some extent over a 5-fold range of concentrations. Mycelia which had been exposed to 1 mM CdCl_2 did not yield a linear response over the same concentration range. In spite of the lack of linearity, the Cd/hemoglobin affinity assay was performed on mycelial homogenates from wild-type and two Cd^R mutants. Homogenates were prepared from mycelia which had been exposed to either 0 or 1 mM CdCl_2 . As shown in Table 12, both the wild-type and the two Cd^R mutant extracts show almost a 50% decrease in the Cd-binding potential when the cells had been grown in the presence of CdCl_2 . It is possible that the Cd-binding sites of the proteins not previously exposed to cadmium

Table 12 Mycelial Levels of Cadmium Binding Capacity for Wild-type and the Cadmium-resistant Mutants in the Absence and Presence of Cadmium

Isolate	Growth Conditions (mM CdCl ₂)	<u>nmoles Cd Bound</u> ml Homogenate	<u>mg Protein</u> ml Homogenate	<u>nmoles Cd Bound</u> mg Protein	Ratio ^a
Wild-type	0	2.1	0.5	4.6	0.4
Wild-type	1	1.6	0.9	1.8	
9A2R5	0	1.4	0.2	6.7	0.3
9A2R5	1	1.6	0.8	2.1	
15-10R22	0	1.4	0.3	4.4	0.5
15-10R22	1	1.8	0.9	2.1	

^aRatio of nmoles Cd Bound/mg Protein from isolate grown on 1 mM CdCl₂ to Cd Bound/mg Protein from isolate grown on 0 mM CdCl₂. These values represent the data from 1 of 2 experiments.

were able to react with the added ^{109}Cd , whereas the Cd-binding sites of the proteins from the mycelia grown in the presence of 1 mM CdCl_2 were already partially filled with non-radioactive cadmium and not available for rapid exchange. In any case, there is no substantial difference in the Cd-binding potential between extracts of the wild-type and either Cd^R mutant. Apparently in these strains cadmium-resistance is not due to an increase in cytosolic cadmium-binding potential.

In an attempt to visualize the various Cd-binding proteins, an aliquot of each homogenate (containing the bound ^{109}Cd) was electrophoresed on a nondenaturing acrylamide gel and the ^{109}Cd was visualized by autoradiography. Although a sufficient amount of radioactivity was loaded per lane, no bands were visible. The ^{109}Cd must not have been bound tightly enough to proteins to remain during the electrophoretic process.

Heat Shock/Cadmium Shock

Upon exposure of Neurospora conidiospores to supraoptimal temperatures, the translation of many normal mRNAs into proteins ceases and the synthesis of a large amount of a few characteristic proteins, known as the HSPs, becomes prominent. In some organisms, heavy metals induce a similar stress response. To determine whether cadmium induces a response similar to that produced by

elevated temperatures in Neurospora, conidia were incubated at either 30°C or 45°C and compared to conidia incubated at 30°C in the presence of 1 mM or 10 mM CdCl₂. Proteins synthesized in these cells were radioactively labelled by adding [³⁵S]methionine; proteins in the cellular extracts were separated by SDS polyacrylamide gel electrophoresis and visualized by fluorography. Figure 22 shows that those proteins stored in the conidia (stainable by Coomassie blue) are identical under each condition. A difference is only evident in the ³⁵S profile of the newly-synthesized proteins (Figure 23). It is apparent that the presence of toxic cadmium ions, whether at a low concentration (1 mM CdCl₂) which does not severely inhibit the growth of mycelia, or at a high concentration (10 mM CdCl₂), does not elicit the same response as elevated temperature. Even though the HSPs are evident in the control lane and the two Cd-treated lanes, only the 45 C lane shows the true heat shock response: enhanced appearance of the HSPs and a reduction in the amount of the non-HSP proteins.

Effect of Cadmium on Cytoplasmic Protein Profile

Finally, a broad-spectrum characterization of the cytosolic proteins synthesized by both the wild-type strain and two representative Cd^R mutant strains was

Figure 22. Electrophoretic pattern of Coomassie-stained [^{35}S]methionine-labelled heat shock and cadmium shock proteins. Proteins were extracted from conidiospores that were incubated at 30°C (lane 1), at 30°C in 1 mM CdCl_2 (lane 2) or 10 mM CdCl_2 (lane 3), or at 45°C (lane 4). The conidia were radiolabelled for 30 min after culture inoculation. The same number of counts was applied to each lane.

Figure 22

Figure 23. Fluorogram of [^{35}S]methionine-labelled heat shock and cadmium shock proteins for the same gel electrophoretic pattern in Figure 22. The arrows designate the three major heat shock proteins.

Figure 23

performed. In the event that the presence of increased amounts of metallothionein was missed by the Cd/hemoglobin assay, or that any other prominent changes in major proteins took place, a comparison of proteins in crude homogenates was made for the Cd^S wild-type strain and the Cd^R mutants 9A2R5 and 15-10R22. Homogenates were prepared from mycelia which had been grown in medium containing either 0 or 1 mM CdCl₂ in the presence of Na₂³⁵S₄ and then subjected to electrophoresis. By viewing the Coomassie-stained protein electrophoretic pattern (Figure 24) and the corresponding ³⁵S fluorogram (Figure 25), a series of comparisons were made. Because the cells were grown the entire time in the presence of the ³⁵S label, these two patterns should be similar, if not identical, and therefore will be discussed together. If a protein were particularly rich in sulfur, it would be expected to label more intensely than other proteins present at the same level. Also, any major proteins simply stored in the conidia might be detected in the Coomassie-stained profile but absent in the fluorograph, although any such stored proteins would be expected to be largely diluted out during the 25 hour growth period.

In comparing the wild-type and two mutant homogenates prepared from cells grown in the absence of cadmium (Lanes 1, 3, and 5), the profiles of proteins detected by staining and fluorography are similar. The only two

Figure 24. Electrophoretic pattern of Coomassie-stained $[^{35}\text{S}]\text{O}_4$ -labelled proteins from wild-type and Cd^{R} mycelia grown in the absence or presence of cadmium. Mycelia were grown in either 0 or 1 mM CdCl_2 and labelled during the entire 25 hr growth period. Lanes 1 and 2: wild-type grown in 0 and 1 mM CdCl_2 , respectively. Lanes 3 and 4: Cd^{R} 9A2R5 grown in 0 and 1 mM CdCl_2 , respectively. Lanes 5 and 6: Cd^{R} 15-10R22 grown in 0 and 1 mM CdCl_2 , respectively. Approximately the same amount of TCA-precipitable radioactive protein was applied to each lane. The protein molecular weight standards were ovalbumin (45,000) and lysozyme (14,300).

Figure 24

Figure 25. Fluorogram of [^{35}S]O₄-labelled proteins for the same gel electrophoretic pattern in Figure 25. The [^{14}C]-labelled protein molecular weight standards were (from the slowest-migrating to the fastest-migrating): ovalbumin (43,000), α -chymotrypsinogen (25,700), β -lactoglobulin (18,400), lysozyme (14,300), and bovine trypsin inhibitor (6,200). Circles mark protein bands that either increased or decreased in amount in the presence of cadmium. Triangles mark the protein bands that either appeared or disappeared in the presence of cadmium.

Figure 25

differences are between wild-type (lane 1) and mutant 9A2 (Lane 3): mutant 9A2 exhibits the reduction in amount of a protein band at 39,000 and the lack of a band at 34,000. Otherwise, each strain accumulates an analogous set of proteins during growth on medium without cadmium.

The wild-type protein profiles from cells grown in the absence and presence of Cd are very similar (Lanes 1 and 2). The wild-type (and also each mutant) lacks a major protein band at 72,500 in the presence of Cd. A Coomassie-stained band at 40,000 and an ^{35}S -labelled band at 39,000 are increased in intensity in the presence of Cd.

Whereas cadmium has a minor effect on the wild-type strain causing an increase in amount of two proteins and the loss of one protein, it has a profound effect on each of the mutants. The altered protein pattern is generally the same for both of the mutants, but some of the changes induced by cadmium are unique for a particular mutant.

There are prominent differences between the mutant 9A2 protein profiles of mycelia grown in the absence and presence of Cd (Lanes 3 and 4). Most of these differences are also found with mutant 15-10 (Lanes 5 and 6). A major protein band at 72,500 and three minor bands (49,000, 45,700, 42,100) which lie between major bands at 52,000 and 39,800 are missing in the presence of Cd (Coomassie

profile and fluorograph). Major protein bands at 52,000 and 39,800 are present in larger amounts, as are minor bands at 21,400 and 19,500. The band at 19,500 is especially prominent on the fluorograph (Lane 4) for both mutants grown in the presence of Cd but is only faintly visible in the wild-type grown with Cd present. In addition, in the mutants in the presence of cadmium, new, but very faint protein bands appear at 33,100 and 30,200 (these proteins are both Coomassie-stained and radioactively-labelled).

Although both mutants exhibit almost identical profiles in the presence of Cd, a protein band at 35,500 is much more intense in mutant 15-10 than in 9A2.

DISCUSSION

Cadmium toxicity and the resistance to the deleterious effects of this toxic metal have been studied in many organisms, including common experimental systems and less-characterized systems, entire organisms and cell cultures. Because of the importance in understanding metal metabolism and also the focus on environmental contamination and occupational metal contact, biochemical research in the area of enzymatic and non-enzymatic metalloproteins is of current interest.

N. crassa is an ideal organism for metal toxicity studies, based on its ease of manipulation, its haploid nature, and its well-characterized genetic structure. No successful attempts to characterize cadmium-resistant mutants in *Neurospora* have been reported; however, a copper thionein from wild-type *N. crassa* grown in the presence of copper has been purified and sequenced. Surprisingly, all seven cysteinyl residues of this thionein occupy the same positions as the first seven cysteines in the amino-terminal region of mammalian MT, which is three times larger than the *Neurospora* protein. This protein does not confer cadmium-resistance; its

synthesis is only induced by copper and it only binds cuprous ions.

The isolation of mutants resistant to toxic concentrations of cadmium would provide Neurospora with a new selectable marker and also a new genetic locus for mapping purposes. Few N. crassa genes have been isolated to date, and the structure and function of upstream regulatory regions is not well understood. The mammalian metallothionein gene has 5' regulatory regions conferring transcriptional regulation on the gene by both heavy metals and glucocorticoids. It would be interesting to study the control of a cadmium-binding protein in Neurospora; it would provide a useful case with which to study.

Another interesting question which could result from the isolation of a fungal cadmium-binding protein and its gene would be the possibility of coamplification of adjacent genes. In both yeast and Syrian hamster cells, when the selection of resistant cells on higher concentrations of a toxic compound is a result of tandem gene amplification (e.g. the metallothionein gene), a neighboring gene in the repetitive unit is coamplified. Coamplification could be an ideal technique to use to obtain large amounts of a protein (e.g. a regulatory protein) which would normally only be present at very low levels in a cell.

Several methods for the selection of Cd^R mutants in N. crassa were used, each with the potential to create different types of mutations. Irradiation of cells with UV light causes base pair alterations and occasionally deletions or chromosomal rearrangements. These mutants would probably each only have mutations in a single gene that uniquely confers resistance upon the organism. Since cadmium interferes with nucleic acid synthesis, Cd^R mutants selected by a single plating of untreated cells on medium containing cadmium might also contain a mutation in a single gene (most chemical mutagens induce point mutations). Training flasks, however, due to their adaptive nature of culturing, allow for the accumulation of several mutations which might work in concert to confer a resistant phenotype. Gene amplification also may occur in training flasks.

After subjecting the cells to the various mutagenic techniques, it was difficult to distinguish between the sensitive isolates and those which were definitely cadmium-resistant. The traditional methods were to either place a chunk of agar containing some mycelia onto medium containing the selective agent or to add a "light" conidial inoculum to a test tube containing liquid medium and the selective agent. However, such tests for cadmium resistance were not reliable and gave variable results, depending on the size of the inoculum. It appeared that

with a larger inoculum, the cells were able to overcome the toxic effects of cadmium. It was possible that the cells were depleting the ions from the medium and thus creating a less toxic environment. Regardless of the exact mechanism, it was clear that toxicity was definitely a function of the number of cells in the assay.

To maintain uniform inocula, a pasteur pipette plug method for removing a mycelial inoculum was devised and used for each step in the selection: (1) determining inhibitory levels of cadmium in the medium, (2) preparing growth rate plates for the selection of Cd^R mutants, and (3) analyzing progeny from a genetic cross. For analysis in liquid medium, the inoculum was held at 20 conidia per milliliter of medium (obtained by serial dilution of a conidial suspension). With these two techniques, I overcame the difficulty in scoring the Cd^R phenotype and the distinction between cadmium-resistance and cadmium-sensitivity was clear-cut.

Cadmium-resistant mutants were obtained by all three techniques, and one naturally- Cd^R wild-type isolate from Japan was also obtained. Each was crossed to the standard Cd^S wild-type strain to obtain reisolates with the Cd^R trait that was transmissible through meiosis; then each mutant was genetically mapped using standard nutritional and morphological mutants. One Cd^R mutant obtained by

single step cadmium-mutagenesis, 9A2R5, mapped to the left arm of LG II, whereas the other single-step mutant (4A2R1) as well as the training flask mutant (TFR38) and the UV-induced mutant (15-10R22) all mapped to the left arm of LG VII. Interestingly, the Japanese Cd^R mutant mapped to a location outside of these two regions, as it segregated Cd^S progeny when crossed to any of these Cd^R mutants.

The three Cd^R mutants that mapped to LG VII exhibited altered morphologies. The training flask mutant grown on a solid surface in the absence of cadmium appeared to be osmotically unstable. The aerial hyphae either leaked their cytoplasmic contents or secreted an extracellular fluid, and in addition, this mutant did not conidiate well. When grown on solid medium in the presence of a colonizing factor, the colonies were larger and more diffuse than wild-type. When crossed to a Cd^S isolate, the large size was transmitted 100% with the Cd^R trait. Thus, this mutant definitely possesses cell wall/plasma membrane alterations. These characteristics may actually confer the resistance trait, or could be a result of the Cd^R phenotype.

The Cd^R mutant 4A2 also did not conidiate well. When grown on a solid surface containing a colonizing factor it, too, produced very large colonies. In this mutant, however, the large size was not transmitted with cadmium-resistance and the colonies were distinct rather than

diffuse. The cell wall/plasma membrane alterations in this mutant may be different than those in the training flask mutant. One final morphological alteration was that of the Cd^R mutant 15-10. Its conidial masses appeared bulbous or beaded, rather than filamentous or thread-like. In general, all of the cadmium resistant mutants had characteristics that suggested they had an altered cell membrane or cell wall. It seems virtually certain that these morphological alterations are directly related to the cadmium resistance and may be responsible for the resistant phenotype.

When the Cd^R mutant 9A2 (LGII) was crossed to the Cd^R mutant 15-10 (LG VII), the progeny segregated $Cd^S:Cd^R$ as expected for mutations on two different chromosomes (1:3). Interestingly, there was a definite difference in resistance with the pasteur pipette plug assay among the Cd^R progeny: some (less than half) appeared to be much more resistant to cadmium than the others and grew noticeably further away from the point of inoculation. Those progeny exhibiting the higher degree of cadmium-resistance may have received a "double dose" of resistance-conferring factors: both the mutant 9A2 gene and the mutant 15-10 gene. Progeny with a lower degree of resistance would presumably be those with only a single resistant trait. A similar situation was noted in

actidione-resistant strains of N. crassa (Hsu, 1963). Double mutants, containing mutant genes at two independent actidione loci, were more resistant to higher concentrations of actidione than single mutants. The double mutants were also slower-growing than the single mutants or the wild-type in the absence of actidione, signifying the requirement of at least one wild-type allele for normal growth.

Another interesting comparison is between wild-type and the Cd^R mutants grown in liquid medium. These mutants were all initially selected because they surpassed wild-type growth first on solid medium and then in liquid, standing culture. When the growth rates were all compared in liquid, shaking medium, none of the Cd^R mutants were able to grow as well in the presence of cadmium as they did in its absence, nor were they able to grow as well as wild-type, either in the presence or absence of cadmium.

The inability of the Cd^R mutants to grow as well as wild-type in this experiment must have been related to the nature of the growth conditions. When grown on agar medium or in standing liquid culture, the mycelia were either growing on top of the cadmium-containing source or able to escape it as aerial hyphae, whereas in shaking liquid culture the mycelia were constantly immersed in the cadmium source. These Cd^R mutants clearly have some cell wall/plasma membrane alterations and these may affect

their ability to grow under such conditions. When grown on the surface of a cadmium source, the small amount of mycelia that comes in contact with the metal must be able to support the growth of the additional mass (by resisting the toxic effects of the metal and transporting nutrients from the medium) and it might even have a slightly altered morphology than the cells not in contact with the cadmium.

Additional evidence of a morphological difference was the texture of the mycelial mat. The method used to harvest the mycelia from shaking liquid culture was suction filtration. The mycelia collected on the piece of filter paper and formed a thick, fibrous mat or pad. Even though the wild-type did not produce as much mycelial mass in the presence of cadmium, the mycelial mat still maintained a fibrous nature, whereas the mutant mats took on a crumbly or almost powdery texture in the presence of cadmium.

At higher concentrations of cadmium, the training flask mutant was able to tolerate the toxic metal ions better than mutant 9A2 or 15-10 and in the latter two mutants the appearance of a lag in growth was especially evident at the higher concentrations of cadmium. The lag might be the time required for the cells to develop any biochemical or morphological alterations required for adaptation to the toxic conditions (especially in the case

of mutant 9A2 and 15-10 in 2 mM CdCl_2), or it might simply represent the inability of most cells to grow at that high cadmium concentration. The "Death vs. Inhibition" experiment showed that in wild-type, the cells which were unable to grow in the presence of cadmium were not just growth-inhibited, but actually dead. The training flask procedure did not give rise to mutants with levels of cadmium-resistance much higher than those produced by UV-irradiation. It was hoped that the repeated culture of Neurospora in the presence of increasingly higher concentrations of cadmium would yield much higher resistances due to a gradual amplification of a gene encoding a cadmium-binding protein. However, no such strains with high cadmium resistance were obtained. Increases in gene copy number may require that more than a single copy of the gene be present (as in yeast) and may not be easily accomplished in Neurospora without the introduction of additional (plasmid-borne?) gene copies.

Cadmium-resistance in the N. crassa mutants could be due to a decrease in cadmium uptake whereby the entrance of cadmium into the cell was blocked by an alteration in the cell surface receptors or other membrane components. Thus, transport of cadmium into wild-type and the Cd^R mutants was investigated.

A copper-resistant mutant of S. cerevisiae took up less copper than the copper-sensitive strain, and by using

protoplasts, it was determined that this decrease in copper-permeability was due to an alteration in membrane transport properties (Gadd et al., 1984). In S. aureus cells, extracellular cadmium actually entered the cells via the manganese transporter, having a higher affinity for this transport system than even did manganese. This accumulation was irreversible in Cd^S cells and its inhibition of respiration caused cell death. Cadmium accumulation in Cd^R cells, on the other hand, was prevented by a plasmid-encoded Cd⁺⁺/proton antiporter which expelled cadmium from the cell, thereby creating a less-toxic intracellular environment (Tynecka et al., 1981). Resistance can therefore be conferred by a decrease in permeability or an increase in efflux.

Cadmium uptake in N. crassa mycelia does not appear to occur by a single transport system. At concentrations of 0.1 mM CdCl₂ and lower, cadmium uptake was slow and gradual over a three hour time period; the rate of uptake increased sharply at concentrations of 0.5 mM and above. S. cerevesiae exhibits similar cadmium uptake: at lower concentrations of cadmium (0.005 mM and 0.05 mM) cadmium uptake was also slow and gradual and a similar sharp increase was noted upon a ten-fold increase in concentration to 0.5 mM (Gadd and Mowll, 1983). The major difference between the Neurospora and yeast cadmium

transport is the saturability of the yeast system.

Neurospora did not exhibit saturation of the transport system, as would be the situation if only a single mode of transport existed.

That multiple transport systems probably exist was also apparent when the energy-dependence of cadmium transport was investigated. Seventy-five percent of cadmium transport was energy-requiring, whereas the other twenty-five percent was independent of the energy-state of the cells. Cadmium seemed to enter the cells via both active transport and an energy-unrelated means, which could include carrier-mediated mechanisms or free diffusion.

In many cases the transport of cadmium into the cell is facilitated by a transporter molecule that normally transports a physiological divalent cation, often of similar ionic radius. Yeast cobalt and cadmium uptake was competitively inhibited by cations of similar size (Norris and Kelly, 1977), and thallium uptake is via an energy-dependent potassium ion uptake system (Norris, et al., 1976).

None of the divalent cations, whether similar to the size of cadmium or not, had any inhibitory effects on cadmium uptake in Neurospora. Apparently cadmium enters the cells via a novel system not shared by any of the other transition metals tested. Also, cadmium had no

effect on its own transport, because a preincubation with nonradioactive cadmium did not alter ^{109}Cd uptake.

The most interesting effect of divalent cations on cadmium transport was the enhancement of cadmium transport by copper. When copper was present at an equimolar concentration, an 11-fold increase in the amount of cadmium transported occurred. This increase was not due to the de novo synthesis of a cadmium transport system in the presence of copper, because inhibition of protein syntheses either prior to the simultaneous addition of cadmium and copper, or prior to a preincubation with copper had no effect on cadmium transport. It is possible that the binding of copper to a transport system somehow facilitated additional binding of cadmium to the transporter, thereby increasing the amount of cadmium transported into the cell. Another possibility is that copper entered the cell via an antiporter molecule, and when the opposite ion was extruded from the cell, multiple translocations of cadmium into the cell occurred.

Once cadmium transport in the wild-type Cd^{S} strain was characterized, transport was compared between the wild-type and one isolate of each representative Cd^{R} mutant. There was no obvious difference in the $\mu\text{moles Cd/g dry weight}$ accumulating over a 90 minute period among these three isolates. (Referring back to the mycelial mat

texture of the Cd^{R} mutants grown in the presence of cadmium, it is possible that over an extended period of growth in the presence of cadmium, the mutants do take up more cadmium than the wild-type and hence the powdery texture.) When assayed in the presence of an energy-formation-inhibiting compound there was again no difference between the Cd^{R} and Cd^{S} isolates. There seemed to be no alteration in either the energy-dependent transport mechanism or the energy-independent process. Apparently some mechanism other than the inhibition of transport of cadmium into the Cd^{R} cell is conferring the resistant phenotype.

The cadmium which entered the cell was not tightly bound, as evidenced by the rapid efflux of cadmium out of the wild-type cell. Over a 90 minute period, approximately two thirds of the cadmium which had entered the cell during the accumulation period then left the cell, and the loss was independent of the extracellular metal type or concentration. It is possible that the Cd^{R} mutants have an altered cadmium efflux such that more cadmium exits from the cell than in the case of the wild-type. However, because the amount of radioactive cadmium remaining in the cells after a 90 minute incubation was the same for all strains, it is doubtful that an efflux system is involved in cadmium-resistance.

The synthesis of the heavy metal-binding protein metallothionein is one of the most common and well-characterized mechanisms known to confer the cadmium-resistant phenotype on an organism. Upon the entrance of cadmium into the cell and the rise in its intracellular concentration, transcription of the metallothionein gene is stimulated with the eventual rise in MT concentration. The MT protein has the capacity to bind at least seven atoms of metal per molecule. Metallothionein is presumed to play a role in the regulation of the cellular concentration of zinc; it is naturally present at a basal level in organisms not exposed to toxic metals and it is prevalent in specialized secretory, absorptive and nutrient cells. Cultured Friend leukemia cells selected for high levels of cadmium-resistance accumulate at least 30 times more MT than the cadmium-sensitive cell line--91% of the heat stable proteins are composed of MT (Beach and Palmiter, 1981).

The results of the cadmium/hemoglobin affinity assay showed that there is not an increase in cadmium-binding potential when N. crassa cells are exposed to cadmium nor do the cadmium-resistant mutants have an elevated cadmium binding potential. With this data it is impossible to rule out either the possibility that N. crassa can be induced to synthesize MT upon cadmium exposure or that the mutants are capable of a more significant MT induction

than the wild type strain. The cadmium/hemoglobin assay is based on two important assumptions which, although they are valid for most the organisms whose MT is well-characterized to date, may not be the case in N. crassa. The first assumption is that MT is a heat-stable protein. In mammalian systems, as well as several marine organisms, Drosophila, and algae, the MT protein is able to withstand temperatures of at least 60-70°C. The grass cadmium-binding protein is a heat-stable protein, but the cadmium-binding protein of maize is not heat-stable. Attempts to isolate maize cadmium-binding protein using the heat denaturation step also used for grass were unsuccessful and were subsequently omitted. It is possible that a major cytoplasmic protein functioning in sequestering cadmium in N. crassa is heat labile and therefore discarded in the preparation of the homogenate for the cadmium-hemoglobin assay. The other assumption is that either there are cadmium-binding sites of the MT molecule which are unoccupied and therefore available to bind with the added ^{109}Cd or else that the cadmium atoms are bound in such a manner that they are able to readily exchange with the free ^{109}Cd in solution. When exposed to cadmium, N. crassa displayed a 50% drop in the heat-stable cadmium-binding potential. It would be surprising if the cells were actually synthesizing less cadmium-binding protein in

the presence of the metal than in its absence. It is possible, however, that when the cells are grown in the presence of cadmium, the majority of the protein's cadmium-binding sites interact with the intracellular metal and thus are not able to bind the free ^{109}Cd . In this situation, it would appear that the cells not exposed to cadmium would have a higher cadmium-binding potential than those exposed to cadmium, as I observed with Neurospora. Since cadmium exits from the cells, as seen in the efflux experiments, and since it does not remain tightly bound to protein during electrophoresis, it is difficult to determine if this is the case.

A strong argument against metal induction of MT is the absence of a strong low molecular weight protein band in the ^{35}S in vivo labelling protein profile. One would presume, that since the MT protein is composed of 30% cysteine residues, a low molecular weight band would stand out as being heavily labelled with sulfur, particularly after induction by cadmium (it would probably be more noticeable by fluorography than by Coomassie staining). The only possible candidate for this protein would be the 19,500 band synthesized by both Cd^{R} mutants. This band is synthesized to a greater extent in the presence of cadmium than in its absence, compared to other protein bands of the same intensity, it is more highly labelled with ^{35}S .

Cross-resistance of a Cd^R mutant to related divalent cations can be due to an increase in MT content; often the synthesis of increased amounts of the MT protein can be induced by metals other than cadmium. It is not surprising that the Cd^R mutants which I have studied are sensitive to toxic levels of zinc, cobalt, and copper, since cadmium, itself, does not appear to induce MT synthesis. If, however, a metallothionein is synthesized in the presence of cadmium, this Cd-thionein may not be induced by, or able to bind to, other metals (as in the case of the yeast copper-binding protein). It is possible that these metal ions interact with the cell wall/plasma membrane complex in a different manner than cadmium ions; my results showed that none of these three metals compete with cadmium for intracellular transport. Thus, these other metal ions may have a different mode of cytoplasmic accumulation.

Many of the data presented in this study point to the possibility that cell wall/plasma membrane alterations may be the means by which the Cd^R mutants of N. crassa described herein obtain their resistance. All of the mutants show similar amounts of cadmium transport and neither Cd^R mutant 9A2 nor 15-10 exhibit an appreciable increase in a high-sulfur-content heavy metal-binding protein. For this reason, there must be a difference between the wild-type and the Cd^R mutant in the way the

cadmium is treated once it is associated with the cells.

The three Cd^R mutants mapping to LG VII all exhibited morphological alterations to various degrees. Yeast cells grown in silver nitrate contain silver deposits in vacuoles and in the cell wall (Brown and Smith, 1976). Another type of yeast manifests its copper resistance by the precipitation of copper sulfide deposits in the cell wall (Ashida et al., 1963). At low toxic cadmium concentrations, Dictyostelium exhibits several morphological alterations (Cyr and Bernstein, 1984). Cadmium blocked the endocytic event of phagocytosis and also created a shift in internal vacuole size so that more but smaller vacuoles appeared. These changes caused a shift in total membrane from internal membrane to the plasmalemma, and these cadmium-treated cells were larger than untreated cells. Because cadmium binds to membrane components and increases membrane stability, the effect on membrane fluidity can alter membrane utilization and compartmentalization. Vinclozolin-resistant mutants of N. crassa exhibited poor radial growth and sporulation (Grindle, 1984); they were very sensitive to media of high osmotic pressure, probably due to lesions or alterations in the cell wall/plasma membrane complex which reduced tolerance to mechanical stress.

In addition to synthesizing three cadmium-binding proteins with some resemblance to MT, a cadmium-resistant

strain of Pseudomonas putida showed three changes similar to that of Neurospora when grown in the presence of cadmium: (1) changes in cellular morphology, (2) a long lag phase before growth, and (3) a prolonged phase of exponential growth followed by a rapid decline in cell number due to cell lysis. Much of the cytoplasmic cadmium was associated with polyphosphate granules (Higham et al., 1984). It is therefore possible that the Cd^R trait of my N. crassa mutants is conferred by an alteration in the cell wall/plasma membrane complex that allows the cells to tolerate a higher extracellular and intracellular cadmium concentration than can the sensitive parental strain.

When Neurospora conidiospores are germinated at elevated temperatures, the translation of normal RNA stops and the synthesis of a large amount of the heat shock proteins (HSPs) is all that is evident (Plesofsky-Vig and Brambl, 1985). The three major HSPs produced have molecular weights of 98,000, 83,000, and 67,000. The synthesis of these HSPs has been shown to protect the cells against the lethal effects of exposure to even higher temperatures, and to be required for the cells to recover from heat shock and to resume synthesis of the normal proteins. In many organisms, elevated temperature is not the only stress that induces the heat shock response: heavy metals, amino acid analogues, and

superoxide ions are among some of the other treatments. All of the conditions that induce this response are capable of causing protein denaturation, therefore making the intracellular environment less physiological. Very recently it was demonstrated that denatured proteins (unrelated to the HSPs or their regulation) were able to stimulate the expression of a heat shock gene, whereas the same proteins in their native state did not (Ananthan et al., 1986). The size of the pool of proteins designated for degradation and proteolysis is thought to be one regulatory factor involved in the heat shock response. By binding to sulfhydryl groups, cadmium and other heavy metals can cause conformatational changes in intracellular proteins and thus trigger this stress response.

When exposed to either low non-toxic levels of cadmium or higher, toxic levels, N. crassa conidiospores did not exhibit the traditional heat shock response. The heat shock bands were visible in the control and cadmium-treated lanes, possibly because the initial transfer of the conidia from an inactive state to conditions favoring growth was a stress, but only upon temperature elevation was the synthesis of the HSPs elevated and the synthesis of the rest of the proteins depressed.

Several reasons may explain why cadmium does not cause the heat shock response in N. crassa. First of all, cadmium may not readily enter the cell. It is possible

that most of the cadmium is associated with the cell wall, in a precipitated form, or associated with the plasma membrane. Cadmium within the cell may not be free in the cytoplasm: it may be sequestered in vacuoles or complexed with proteins such that its presence is not sensed. Alternatively, even when free in the cytoplasm, cadmium may not activate the triggering mechanism that signals the synthesis of the genes in the heat shock circuit.

Chappell et al. (1986) have recently identified HSP 70 as being indistinguishable from the mammalian uncoating ATPase. This protein is involved in the uncoating of clathrin-coated vesicles; following heat shock, this protein is localized in the nucleus, more specifically, the nucleolus, and has been shown to have RNA-binding properties. After further investigation it was discovered that each member of the family of HSPs contains a conserved domain with an ATP-binding site. It is unclear how proteins with ATP binding sites, and therefore potential ATPase activity, could function in the protective role as do HSPs and it is even more vague why an uncoating ATPase would localize in the nucleolus as a stress response.

The most profound effect of cadmium on N. crassa cells and the most noticeable difference between wild-type and the Cd^R mutant was displayed in the electrophoretic

pattern of the cytoplasmic proteins of each strain grown in the absence and presence of CdCl_2 . In the absence of cadmium, the wild-type and the Cd^R mutant synthesize essentially the same proteins, with only a decrease in one protein and the absence of another protein in the mutant 9A2. Cadmium has only a minor effect on the wild-type strain: two bands are increased in intensity and one band at 72,500 is absent. This major protein at 72,500 is lacking in the wild-type and each Cd^R mutant when grown in the presence of cadmium; it is difficult to say whether its synthesis is actually turned off in the presence of the metal or if the protein becomes unstable and is turned over more rapidly.

In contrast, cadmium, had a profound effect on each of the mutants. In several cases the effects were the same for both of the mutants: four proteins were missing (including the major one at 72,500), four proteins were present in larger amounts, and two new proteins appeared. One of the proteins which was increased in the presence of cadmium has a molecular weight of 19,500 and it labeled more intensely in the presence of ^{35}S . This protein could in fact be a metallothionein-related metal binding protein.

Overall, the greatest effect of cadmium observed was its ability to alter the cytoplasmic protein pattern, both in the Cd^S and the Cd^R strains. Cadmium ions do not

appear to evoke the heat shock response, but in the wild-type and especially in the mutants, it causes a definitive alteration in the cytoplasmic protein profile. It is possible that these changes are simply a result of the stress placed on these cells in the presence of cadmium; alternatively the changes may constitute the mechanism that allows the cells to compensate for or resist the stress imposed by toxic metal ions. The three HSPs synthesized upon an immediate temperature stress are distinct from those proteins synthesized after a prolonged cadmium stress. These protein differences could comprise a different set of stress proteins and represent a change in the cytoplasm that permits certain key enzymes to function even in the presence of cadmium.

At this point, several experiments can be suggested for further investigation of the effects of cadmium on N. crassa and the biochemical and morphological alterations induced by the cadmium-resistance-conferring mutations. The subcellular compartmentalization of cadmium should be analyzed to determine if it is associated with the cell wall, the plasma membrane, or any organelles including the mitochondria or lysosomes, or whether it is localized in the cytoplasmic fraction. The surface properties of the plasma membrane could be examined and compared between the wild-type and the Cd^R mutants as done for the os⁻ mutants,

as well as certain constituents of the cell wall. Many biochemical lesions can cause morphological changes in fungi, including alterations in apical growth rates, branching patterns, and the size and shape of individual cellular components. A more thorough classification of these Cd^R mutants at the microscopic level might provide additional information about their morphological alterations and their relationship to cadmium resistance.

It might be possible to select for Cd^R mutants in liquid medium following UV irradiation to bias the selection in the favor of cytoplasmic changes rather than morphological changes. This might enhance the chances of obtaining higher levels of resistance, particularly to isolate metallothionein overproducers. It may be necessary to purify a cadmium-specific MT using a non-classical approach; the putative N. crassa cadmium-binding protein may not be heat stable, may have an unusual cysteine content or an atypical molecular weight.

Other aspects of cadmium transport could be investigated, including the rather puzzling ability of copper to enhance cadmium uptake. Cadmium's entrance into the cell may be coupled to potassium release or even to the expulsion of copper. Preloading the cells with copper and then allowing the cells to transport cadmium in the absence of copper may show whether cadmium entrance is coupled to copper release. It is also possible that

copper could enhance the uptake of other heavy metals, of potassium or sulfate, or even the metabolic compounds such as amino acids or glucose. Finally, cadmium efflux should be performed in the presence of copper and should also be compared between the wild-type and the cadmium-resistant mutants.

Investigations at the nucleic acid level may be the most fruitful of all, determining whether Neurospora synthesizes a cadmium-binding protein with any homology to the yeast or mammalian metal thioneins. Munger et al. (1985) have just isolated and sequenced the Neurospora copper metallothionein gene and found it to be strikingly homologous to the amino-terminal part of mammalian metallothionein. Using a synthetic primer (derived from the known amino acid sequence of the protein) as the basis for primer extension in the presence of enriched Neurospora metallothionein mRNA, they obtained the gene sequence, from which they created a synthetic hybridization probe. With this probe they screened a genomic Neurospora library and obtained the Neurospora copper MT gene. Using this probe in low stringency hybridization might allow detection and isolation of a similar, related thionein gene for cadmium or zinc. Neurospora may synthesize a separate cadmium-binding protein with homology to its copper-binding protein, or to

the yeast copper-binding protein (which is different from the mammalian MT), or even with homology to one of the less-related members of the thionein family. Research in these various directions is essential if the entire picture of metal homeostasis and detoxification is to be understood in Neurospora.

The research I have presented here provides a basis for further investigation aimed at understanding heavy metal resistance in Neurospora. Genetic mutants have been isolated and mapped, and a basic characterization of them has been completed. It will be interesting to understand the relationship between their morphological alterations and the cadmium-resistant phenotype. It will also be interesting to know the total number of genes conferring heavy metal-resistance and their mode of action. Ultimately, it will be important to ascertain the number of different genes encoding metal thioneins, the relationship between the different thioneins, and the mechanisms of regulation involved in the expression of these genes.

LIST OF REFERENCES

- Abel, T. H. and F. Barlocher. 1984. Effects of cadmium on aquatic hyphomycetes. *Appl. Environ. Microbiol.* 48:245-251.
- Alt, F. W., R. E. Kellems, J. R. Bertino, and R. T. Schimke. 1978. Selective multiplication of dihydrofolate reductase genes in methotrexate-resistant variants of cultured murine cells. *J. Biol. Chem.* 253:1357-1370.
- Ananthan, J., A. L. Goldberg, and R. Voellmy. 1956. Abnormal proteins serve as eukaryotic stress signals and trigger the activation of heat shock genes. *Science* 232:522-524.
- Ashburn, M. and J. J. Bonner. 1979. The induction of gene activity in Drosophila by heat shock. *Cell* 17:241-254.
- Ashida, J., N. Higashi, and T. Kikuchi. 1963. An electron microscopic study on copper precipitation by copper-resistant yeast cells. *Protoplasma* 57:26-32.
- Atkinson, B. G. and R. L. Dean. 1985. Effects of stress on the gene expression of amphibian, avian, and mammalian blood cells. IN: Changes in Eukaryotic Gene Expression in Response to Environmental Stress. B. G. Atkinson and D. B. Walden, eds. Academic Press, Orlando, FL.
- Bakka, A. and H. E. Rugstad. 1981. Uptake and egress of cadmium in cultures of cadmium-resistant and the corresponding "wild-type" cells. *Acta Pharmacol. Toxicol.* 48:81-86.
- Beach, L. R. and R. D. Palmiter. 1981. Amplification of the MT-1 gene in cadmium-resistant mouse cells. *Proc. Natl. Acad. Sci. USA* 78:2110-2114.
- Beltramini, M. and K. Lerch. 1982. Copper transfer between Neurospora copper metallothionein and type 3 copper apoproteins. *FEBS Lett.* 142:219-222.

- Benson, W. H. and W. J. Birge. 1985. Heavy metal tolerance and metallothionein induction in fathead minnows: Results from field and laboratory investigations. *Environ. Toxicol. Chem.* 4:209-217.
- Boulanger, Y., C. M. Goodman, C. P. Forte, S. W. Fesik and I. M. Armitage. 1983. Model for mammalian metallothionein structure. *Proc. Natl. Acad. Sci. USA* 80:1501-1505.
- Brady, F. O., M. Panemangalore, and F. A. Day. 1979. In vivo and ex vivo induction of rat liver metallothionein. *Experientia* (Suppl.) 34:261-271.
- Bremner, I. and N. T. Davies. 1974. Studies on the appearance of a copper-binding protein in rat liver. *Biochem. Soc. Trans.* 2:425-427.
- Brenes-Pomales, A., G. Lindegren, and C. C. Lindegren. 1955. Gene control of copper-sensitivity in Saccharomyces. *Nature* 176:841-842.
- Brown, P. C., S. M. Beverly, and R. T. Schimke. 1981. Relationship of amplified DHFR genes to (double minute) chromosomes in unstably resistant mouse fibroblast cell lines. *Mol. Cell Biol.* 1:1077-1083.
- Brown, T. A. and D. G. Smith. 1976. The effects of silver nitrate on the growth and ultrastructure of the yeast Cryptococcus albidus. *Microbios. Lett.* 3:155-162.
- Butt, T. R., E. J. Sternberg, J. A. Gorman, P. Clark, D. Hamer, M. Rosenberg, and S. T. Crooke. 1984. Copper metallothionein of yeast, structure of the gene, and regulation of expression. *Proc. Natl. Acad. Sci. USA* 81:3332-3336.
- Cameron, E. and C. L. Foster. 1963. Observations on the histological effects of sub-lethal doses of cadmium chloride in the rabbit. I. The effect on the testis. *J. Anat.* 97:269-280.
- Carter, A. D., B. K. Felber, M. J. Walling, M. F. Jubier, C. J. Schmitt, and D. H. Havrec. 1984. Duplicated heavy metal control sequences of the mouse metallothionein-I gene. *Proc. Natl. Acad. Sci. USA* 81:7392-7396.
- Chappell, T. G., W. J. Welch, D. M. Schlossman, K. B. Palter, M. J. Schlesinger, and J. E. Rothman. 1986. Uncoating ATPase is a member of the 70 kilodalton family of stress proteins. *Cell* 45:3-13.

Cove, D. J. 1976a. Chlorate toxicity in Aspergillus nidulans: The selection and characterization of chlorate-resistant mutants. Heredity 36:191-203.

Cove, D. J. 1976b. Chlorate toxicity in Aspergillus nidulans. Studies of mutants altered in nitrate assimilation. Mol. Gen. Genet. 146:147-159.

Cyr, R. J. and R. L. Bernstein. 1984. Morphological changes and depressed phagocytic efficiency in Dictyostelium amoebae treated with toxic concentrations of cadmium. Environ. Res. 35:66-78.

Danielson, K. G., S. Ohi, and P. C. Huang. 1982. Immunochemical detection of metallothionein in specific epithelial cells of rat organs. Proc. Natl. Acad. Sci. USA 79:2301-2304.

Debec, A., R. Mokdad, and M. Wegnez. 1985. Metallothioneins and resistance to cadmium poisoning in Drosophila cells. Biochem. Biophys. Res. Commun. 127:143-152.

De Ruiter, N., V. Mailander, and H. Kappus. 1985. Effect of heavy metals on cellular growth, metabolism, and integrity of cultured Chinese hamster kidney cells. Xenobiotica 15:665-671.

Donn, G., E. Tischer, J. A. Smith, and H. M. Goodman. 1984. Herbicide-resistant alfalfa cells: an example of gene amplification in plants. J. Mol. Appl. Genet. 2:621-635.

Eaton, D. L. and B. F. Toal. 1982. Evaluation of the Cd/hemoglobin affinity assay for the rapid determination of metallothionein in biological tissues. Toxicol. Appl. Pharmacol. 66:134-142.

Endersen, L., A. Bakka, A. Glennas, K. M. Treit, and H. E. Rugstad. 1983. Stability of cadmium resistance and metallothionein levels in vitro and in vivo. Toxicol. Appl. Pharmacol. 67:274-283.

Fischer, A. B. 1985. Factors influencing cadmium uptake and cytotoxicity in cultured cells. Xenobiotica 15:751-757.

Fogel, S. and J. W. Welch. 1982. Tandem gene amplification mediates copper resistance in yeast. Proc. Natl. Acad. Sci. USA 79:5342-5346.

Foster, C. L. and E. Cameron. 1963. Observations on the histological effects of sub-lethal doses of cadmium chloride in the rabbit. II. The effect on the kidney cortex. *J. Anat.* 97:281-288.

Friberg, L., M. Piscator, G. F. Nordberg, and T. Kjellstrom. 1974. In: Cadmium in the Environment. 2nd ed. Cleveland, OH: CRC Press.

Gadd, G. M. and J. L. Mowll. 1983. The relationship between cadmium uptake, potassium release and viability in Saccharomyces cerevesiae. *FEMS Microbiol. Lett.* 16:45-48.

Gadd, G. M., A. Stewart, C. White, and J. L. Mowll. 1984. Copper uptake by whole cells and protoplasts of a wild-type and copper-resistant strain of Saccharomyces cerevesiae. *FEMS Microbiol. Lett.* 24:231-234.

Gick, G. G. and K. S. McCarty, Sr. 1981. Amplification of the metallothionein- I gene in cadmium- and zinc-resistant Chinese Hamster Ovary cells. *J. Biol. Chem.* 257:9049-9053.

Grindle, M. 1984. Isolation and characterization of vinclozolin resistant mutants of Neurospora crassa. *Trans. Br. Mycol. Soc.* 82:635-643.

Grindle, M. and W. Temple. 1982. Fungicide-resistance of os mutants of Neurospora crassa. *Neurospora Newsletter* 29:16.

Grindle, M. and W. Temple. 1983. Fungicide resistance of smco mutants of Neurospora crassa. *Neurospora Newsletter* 30:7-8.

Hager, L. J. and R. D. Palmtier. 1981. Transcriptional regulation of mouse liver MT-I gene by glucocorticoids. *Nature* 291:340-342.

Hamer, D. H., D. J. Thiele, and J. E. Lemontt. 1985. Function and autoregulation of yeast copperthionein. *Science* 228:685-690.

Higham, D. P., P. J. Sadler, and M. D. Scawen. 1984. Cadmium-resistant Pseudomonas putida synthesizes novel cadmium proteins. *Science* 225:1043-1046.

Hildebrand, C. E., J. K. Griffith, R. A. Tobey, R. A. Walkers, and M. D. Enger. 1982. Molecular mechanisms of cadmium detoxification in cadmium-resistant cultured cells: role of metallothionein and other inducible factors. *Dev. Toxicol. Environ. Sci.* 9:279-303

Hildebrand, C. E. and M. D. Enger. 1980. Regulation of Cd^{2+}/Zn^{2+} -stimulated metallothionein synthesis during induction, deinduction, and superinduction. *Biochemistry* 19:5850:5857.

Hildebrand, C. E., R. A. Tobey, E. W. Campbell, and M. D. Enger. 1979. A cadmium-resistant variant of the Chinese hamster (CHO) cell with increased metallothionein induction capacity. *Exp. Cell Res.* 124:237-246.

Hoffman, D. J. and S. K. Niyogi. 1977. Metal mutagens and carcinogens affect RNA synthesis rates in a distinct manner. *Science* 198:313-314.

Howe, Jr., H.B. and C. E. Terry, Jr. 1962. Genetic studies of resistance to chemical agents in Neurospora crassa. *Can. J. Genet. Cytol.* 4:447-452.

Hsu, K. S. 1965. Acriflavin resistance controlled by chromosomal genes in Neurospora. *Neurospora Newsletter* 8:4-6.

Hsu, K. S. 1963. The genetic basis of actidione resistance in Neurospora. *J. Gen. Microbiol.* 32:341-347.

Hsu, K. S. 1962. Genetic basis of drug-resistant mutants in Neurospora. *Rec. Genet. Soc. Am.* 31:92.

Ibrahim, M. A. K. and A. Coddington. 1976. Genetic studies on cycloheximide-resistant strains of Schizosaccharomyces pombe. *Heredity* 37:179-191.

Jacobsen, K. B. and J. E. Turner. 1980. The interaction of cadmium and certain other metal ions with proteins and nucleic acids. *Toxicology* 16:1-37.

Jacobsen, K. B., L. Opresko, R. K. Owenby, and N. T. Christie. 1981. Effects of cadmium on Drosophila: Toxicity, proteins, and transfer RNAs. *Toxicol. Appl. Pharmacol.* 60:368-378.

Kagi, J. H. R., S. R. Himmelhoch, P. D. Whanger, J. L. Bethune, and B. L. Vallee. 1974. Equine hepatic and renal metallothionein. Purification, molecular weight, amino acid composition, and metal content. *J. Biol. Chem.* 249:3537-3542.

Karin, M., A. Haslinger, H. Holtgreve, R. I. Richards, P. Krautner, H. M. Westphal, and M. Beato. 1984a. Characterization of DNA sequences through which cadmium and glucocorticoid hormones induce human MT-IIA gene. *Nature* 308:513-519.

Karin, M., R. Najarian, A. Haslinger, P. Valenzuela, J. Welch, and S. Fogel. 1984b. Primary structure and transcription of an amplified genetic locus: The CUP1 locus of yeast. *Proc. Natl. Acad. Sci. USA* 81:337-341.

Kaufman, R. J., P. C. Brown, and R. T. Schimke. 1981. Loss and stabilization of amplified DHFR genes in mouse sarcoma S-180 cell lines. *Mol. Cell Biol.* 1:1084-1093.

Kelley, P. M. and M. J. Schlesinger. 1982. Antibodies to two major chicken heat shock proteins in widely divergent species. *Mol. Cell Biol.* 2:267-274.

Kito, H., Y. Ose, V. Mizuhira, T. Sato, T. Ishikawa, and T. Tazawa. 1982a. Separation and purification of (Cd, Cu, Zn)-metallothionein in carp hepato-pancreas. *Comp. Biochem. Physiol. (C)* 73:121-127.

Kito, H., T. Tazawa, Y. Ose, T. Sato, and T. Ishikawa. 1982b. Protection by metallothionein against cadmium toxicity. *Comp. Biochem. Physiol. (C)* 73:135-139.

Kondo, N., M. Isobe, K. Imai, and T. Goto. 1985. Synthesis of metallothionein-like peptides cadystin A and B occurring in a fission yeast, and their isomers. *Agric. Biol. Chem.* 49:71-83.

Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685.

Lerch, K. 1980. Copper metallothionein, a copper-binding protein from Neurospora crassa. *Nature* 284:368-370.

Les, A. and R. W. Walker. 1984. Toxicity and binding of copper, zinc, and cadmium by the blue-green alga, Chroococcus parisi. *Water Air Soil Pollut.* 23:129-139.

- Levinson, W., H. Opperman, and J. Jackson. 1980. Transition series metals and sulfhydryl reagents induce the synthesis of four proteins in eukaryotic cells. *Biochem. Biophys. Acta.* 606:170-180.
- Lewis, D. 1963. Structural gene for the methionine-activating enzyme and its mutation as a cause of resistance to ethionine. *Nature* 200:151-154.
- Livingston, L. R. 1969. Locus-specific changes in cell wall composition characteristic of osmotic mutants of Neurospora crassa. *J. Bacteriol.* 99:85-90.
- Margoshes, M. and B. L. Vallee. 1957. A cadmium protein from equine kidney cortex. *J. Am. Chem. Soc.* 79:4813-4814.
- Marzluf, G. A. 1970. Genetic and metabolic controls for sulfate metabolism in Neurospora crassa: isolation and study of chromate-resistant and sulfate-transport-negative mutants. *J. Bacteriol.* 102:716-721.
- Mays, L. L. 1969. Isolation, characterization, and genetic analysis of osmotic mutants of Neurospora crassa. *Genetics* 63:781-794.
- Minagawa, T., N. Yanagishima, Y. Arakatsu, S. Nagasaki, and J. Ashida. 1951. Studies on the adaptation of yeast to copper II. The yeast nucleic acid fraction counteracting the inhibition of yeast growth by copper. *Botanical Magazine Tokyo* 64:753-754.
- Morselt, A. F. W., J. H. J. Copius Peereboom-Stegeman, E. Puvion, and V. J. Maarschalkerweerd. 1983. Investigation for the mechanism of cadmium toxicity at cellular level II. An electron microscopical study. *Arch. Toxicol.* 52:99-108.
- Münger, K., U. A. Germann, and K. Lerch. 1985. Isolation and structural organization of the Neurospora crassa copper metallothionein gene. *EMBO J.* 4:2665-2668.
- Munro, S. and H. Pelham. 1985. What turns on heat shock genes? *Nature* 317:477-478.
- Murasugi, A., C. Wada, and Y. Hayashi. 1981. Purification and unique properties in UV and CD spectra of Cd-binding peptide 1 from Schizosaccharomyces pombe. *Biochem. Biophys. Res. Commun.* 103:1021-1028.

Nagano, T., M. Miwa, Y. Suketa, and S. Okada. 1984. Isolation, physicochemical properties, and amino acid composition of a cadmium-binding protein from cadmium-treated Chlorella ellipsoidea. J. Inorg. Biochem. 21:61-71.

Naiki, N. and S. Yamagata. 1976. Isolation and some properties of copper-binding proteins found in a copper-resistant strain of yeast. Plant Cell Physiol. 17:1281-1295.

Nakamura, Y., S. Katayama, Y. Okada, F. Suzuki and Y. Nagata. 1981. The isolation and characterization of a cadmium and zinc-binding protein from Tetrahymena pyriformis. Agric. Biol. Chem. 45:1167-1172.

Neuhaüser, A., W. Klingmüller, and F. Kaudewitz. 1970. Selektion actidion-resistenter mutanten bei Neurospora crassa sowie ihre genetische und biochemische analyse. Mol. Gen. Genet. 106:180-194.

NIOSH, Division of Standards Development and Technology Transfer. 1984. Cadmium (Cd). (NIOSH Current Intelligence Bulletin No. 42). Atlanta: National Institute for Occupational Safety and Health (NIOSH). (DHHS Publication No. 84-116).

Nordberg, M. and Y. Kojima. 1979. Metallothionein and other low molecular weight metal-binding proteins. Experientia (Suppl.) 34:41-116.

Norris, P. R. and D. P. Kelly. 1977. Accumulation of cadmium and cobalt by Saccharomyces cerevesiae. J. Gen. Microbiol. 99:317-324.

Norris, P., W. K. Man, M. N. Hughes, and D. P. Kelly. 1976. Toxicity and accumulation of thallium in bacteria and yeast. Arch. Microbiol. 110:279-286.

O'Farrell, P. H. 1975. High resolution two-dimensional electrophoresis of proteins. J. Biol. Chem. 250:4007-4021.

Olafson, R. W., R. G. Sim, and K. G. Bolo. 1979. Isolation and chemical characterization of the heavy metal-binding protein metallothionein from marine invertebrates. Comp. Biochem. Physiol.(B) 62:407-416.

Overnell, J. 1982. A method for the isolation of metallothionein from the hepatopancreas of the crab Cancer pagurus that minimizes the effect of the tissue protease. Comp. Biochem. Physiol.(B) 73:547-553.

Plesofsky-Vig, N. and R. Brambl. 1985. Heat shock response of Neurospora crassa: Protein synthesis and induced thermotolerance. J. Bacteriol. 162:1083-1091.

Pongratz, M. and W. Klingmuller. 1973. Role of ribosomes in cycloheximide resistance of Neurospora mutants. Mol. Gen. Genet. 124:359-363.

Porter, H. 1974. The particulate half-cystine-rich copper protein of newborn liver. Relationship to metallothionein and subcellular localization in non-mitochondrial particles possibly representing heavy lysosomes. Biochem. Biophys. Res. Commun. 56:661-668.

Premakumar, R., D. R. Winge, and K. V. Rajagopalan. 1974. Copper-induced synthesis of copper-binding protein in rat liver and yeast. Fed. Proc. 33:1385.

Premakumar, R., D. R. Winge, R. D. Wiley, and K. V. Rajagopalan. 1975. Copper-induced synthesis of copper-chelatin in rat liver. Arch. Biochem. Biophys. 170:267-277.

Prinz, R. and V. Weser. 1975. A naturally occurring Cu-thionein in S. cerevesiae. Hoppe Seylers Z. Physiol. Chem. 356:767-776.

Rausser, W. 1984b. Partial purification and characterization of copper-binding protein from roots of Agrostis gigantea Roth. J. Plant Physiol. 115:143-152.

Rausser, W. E. 1984a. Isolation and partial purification of cadmium-binding protein from roots of the grass Agrostis gigantea. Plant Physiol. 74:1025-1029.

Rausser, W. E. and J. Glover. 1984. Cadmium-binding protein in roots of maize. Can. J. Bot. 62:1645-1650.

Ridlington, J. W. and B. A. Fowler. 1979. Isolation and partial characterization of a cadmium-binding protein from the American oyster (Crassostrea virginica). Chem. Biol. Interact. 25:127-138.

Roper, J. A. and E. Käfer. 1957. Acriflavine-resistant mutants of Aspergillus nidulans. J. Gen. Microbiol. 16:660-667.

- Rugstad, H. E. and T. Norseth. 1978. Cadmium resistance and the content of cadmium-binding protein in two enzyme-deficient mutants of mouse fibroblasts (L-cells). *Biochem. Pharmacol.* 27:647-650.
- Ryan, F. J., G. W. Beadle, and E. L. Tatum. 1943. The tube method of measuring the growth rate of Neurospora. *Am. J. Bot.* 30:784-799.
- Santone, K. S., D. Acota, and J. V. Bruckner. 1982. Cadmium toxicity in primary cultures of rat hepatocytes. *J. Toxicol. Environ. Health* 10:169-177.
- Savino, W., P. C. Huang, A. Corrigan, S. Berrihi and M. Dardenne. 1984. Thymic hormone-containing cells V. Immunohistological detection of metallothionein within the cells bearing thymulin (a zinc-containing hormone) in human and mouse thymuses. *J. Histochem. Cytochem.* 32:942-946.
- Séguin, C., B. K. Felber, A. D. Carter, and D. H. Hamer. 1984. Competition for cellular factors that activate metallothionein gene transcription. *Nature* 312:781-785.
- Shaikh, Z. A. 1982. Metallothionein as a storage protein for cadmium: Its toxicological implications. *Dev. Toxicol. Environ. Sci.* 9:69-76.
- Shapiro, A. L., E. Vinuela, and J. V. Maizel, Jr. 1967. Molecular weight estimation of polypeptide chains by electrophoresis in SDS-polyacrylamide gels. *Biochem. Biophys. Res. Commun.* 28:815-820.
- Shelton, K. R., J. M. Todd and P. M. Egle. 1986. The induction of stress-related proteins by lead. *J. Biol. Chem.* 261:1935-1940.
- Stuart, G. W., P. F. Searle, H. Y. Chen, K. L. Brinster and R. D. Palmiter. 1984. A 12-base-pair DNA motif that is repeated several times in metallothionein gene promotes confers metal regulation to a heterologous gene. *Proc. Natl. Acad. Sci. USA* 81:7318-7322.
- Summers, A. O. and S. Silver. 1978. Microbial transformations of metals. *Ann. Rev. Microbiol.* 32:637-672.
- Tatum, E. L. and G. W. Beadle. 1942. Genetic control of biochemical reactions in Neurospora: An "aminobenzoicless" mutant. *Proc. Natl. Acad. Sci. USA* 28:234-243.

Tomsett, A. B. and D. J. Cove. 1979. Deletion mapping of the *niiA niaD* gene region of Asperigillus nidulans. Genet. Res. 34:19-32.

Tomsett, A. B. and R. H. Garrett. 1980. The isolation and characterization of mutants defective in nitrate assimilation in Neurospora crassa. Genetics 95:649-660.

Trevithick, J. R. and R. L. Metzenberg. 1966. Genetic alteration and pore size and other properties of the Neurospora cell wall. J. Bacteriol. 92:1016-1020.

Tukendorf, A., S. Lyszczyk, and T. Baszynski. 1984. Copper-binding proteins in spinach tolerant to excess copper. J. Plant. Physiol. 115:351-360.

Tynecka, Z., Z. Gos, and J. Zajac. 1981. Reduced cadmium transport determined by a resistance plasmid in Staphylococcus aureus. J. Bacteriol. 147:305-312.

Vomvoyanni, V. 1974. Multigenic control of ribosomal properties associated with cycloheximide sensitivity in Neurospora crassa. Nature 248:505-510.

Waalkes, M. P., S. B. Chernoff, and C. D. Klaassen. 1984a. Cadmium-binding proteins of rat testes. Characterization of a low-molecular-mass protein that lacks identity with metallothionein. Biochem. J. 220:811-818.

Waalkes, M. P., S. B. Chernoff, and C. D. Klaassen. 1984b. Cadmium-binding proteins of rat testes. Apparent source of the protein of low molecular mass. Biochem. J. 220:819-824.

Waalkes, M. P., A. M. Poisner, G. W. Wood and C. D. Klaassen. 1984c. Metallothionein-like proteins in human placenta and fetal membranes. Toxicol. Appl. Pharmacol. 74:179-184.

Wahl, G. M., L. Vitto, and J. Bubnitz. 1983. Co-amplification of rRNA genes with CAD genes in N-(phosphonacetyl)-L-Aspartate-resistant syrian hamster cells. Mol. Cell Biol. 3:2066-2075.

Warr, J. R. and J. A. Roper. 1965. Resistance to various inhibitors in Aspergillus nidulans. J. Gen. Microbiol. 40:273-281.

- Welch, J. W., S. Fogel, G. Cathala, and M. Karin. 1983. Industrial yeasts display tandem gene iteration at the CUP1 region. *Mol. Cell Biol.* 3:1353-1361.
- Westergaard, M. and H. K. Mitchell. 1947. Neurospora V. A synthetic medium favoring sexual reproduction. *Am. J. Bot.* 34:573-577.
- Winge, D. R., R. Premakumar, and K. V. Rajagopalan. 1975. Metal-induced formation of metallothionein in rat liver. *Arch. Biochem. Biophys.* 170:242-252.
- Winge, D. R., R. Premakumar, R. D. Wiley, and K. V. Rajagopalan. 1975. Copper-chelatin: Purification and properties of a copper-binding protein from rat liver. *Arch. Biochem. Biophys.* 170:253-266.
- Wong, K. L. and C. D. Klaassen. 1979. Isolation and characterization of metallothionein which is highly concentrated in newborn rat liver. *J. Biol. Chem.* 254:12,399-12,403.
- Yoshikawa, H. and H. Ohta. 1982. Interaction of metals and metallothionein. *Dev. Toxicol. Environ. Sci.* 9:11-23.