INFORMATION TO USERS

The most advanced technology has been used to photograph and reproduce this manuscript from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps. Each original is also photographed in one exposure and is included in reduced form at the back of the book.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

U·M·I

University Microfilms International
A Bell & Howell Information Company
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
313/761-4700 800/521-0600

Order Number 9105163

**Association of cellular thiol redox status with mitogen-induced
calcium mobilization and cell cycle progression in human
fibroblasts**

Mallery, Susan Regina, Ph.D.

The Ohio State University, 1990

U·M·I
300 N. Zeeb Rd.
Ann Arbor, MI 48106

ASSOCIATION OF CELLULAR THIOL REDOX
STATUS WITH MITOGEN-INDUCED CALCIUM MOBILIZATION
AND CELL CYCLE PROGRESSION IN HUMAN FIBROBLASTS

DISSERTATION

Presented in Partial Fulfillment of the Requirements for
the Degree Doctor of Philosophy in the Graduate
School of the Ohio State University

By

Susan Regina Mallery B.S., M.A., D.D.S.

* * * * *

The Ohio State University

1990

Dissertation Committee

R.E. Stephens

G.P. Brierley

C.M. Allen

Approved by

A handwritten signature in dark ink, appearing to read 'R.E. Stephens', is written over a horizontal line. The signature is fluid and cursive.

Advisor

Department of Pathology

Maripat and Sandy

Je me souviens

ACKNOWLEDGEMENTS

Because of the nature of my project, I was able to interact with three very fine and diverse groups of people. Support from the College of Dentistry was superb. Dean Wallace's and Assistant Dean Melfi's efforts to facilitate the project administration and provide me with encouragement were greatly appreciated. Dr. Carl Allen supervised, quite expertly and also patiently, my clinical training. Ms. Marilyn Travis provided expert secretarial and emotional support through the seemingly endless revisions of the manuscripts.

Drs. Gerald Brierley and Dennis Jung, from the Department of Physiological Chemistry, afforded me the opportunity to learn and benefit from their scientific expertise. It was a delight to interact with individuals that are both so truly interested and competent in science.

Finally, I wish to thank my advisor, Dr. Ralph Stephens. Throughout the arduous process of trying to attain a Ph.D., Dr. Stephens remained very supportive and encouraging. He has the genuine interest and ability to be concerned for his students as individuals, and he did much to help with my personal and scientific growth.

VITA

June 16, 1954.Born - Parkersburg, W. Va.
1976.B.S., Zoology, Ohio State University
Columbus, Oh
1978.M.A., Physiology, Ohio State University
Columbus, OH
1981.D.D.S., Ohio State University
Columbus, Oh
1981-1983.Private practice general dentistry,
Cincinnati, Oh

PUBLICATIONS

"Gender-related variations in and the interaction of
cyclooxygenase metabolism of arachidonate and the oxidative
burst products of human neutrophils." Mallery, S.R.,
Zeligs, B.Z., Ramwell, P.W., Bellanti, J.A. J. Leukocyte
Biology, 40:133-146, 1986.

FIELDS OF STUDY

Major Field: Pathology, Dr. Ralph Stephens
Studies In: Oral Pathology, Dr. Carl M. Allen
Physiological Chemistry, Dr. Gerald P.
Brierley

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
VITA.	iv
LIST OF TABLES.	vi
LIST OF FIGURES.	vii
CHAPTER	
I. INTRODUCTION AND LITERATURE REVIEW.	1
Eucaryotic Cell Cycle	2
Importance of Go and G1 Phases of the Cell Cycle. .	6
Mitogenic Signal Transduction	12
Cellular Thiol Redox Status; Glutathione.	20
GSH and the Cell Cycle.	24
Association of GSH with Macromolecular Synthesis. .	26
Intracellular Ca ²⁺ Homeostasis and Ca ²⁺ Mobilization	31
Cellular Thiol Redox Status and Ca ²⁺ Sequestration.	36
II. MATERIALS AND METHODS	43
III. RESULTS	60
IV. DISCUSSION.	98
LIST OF REFERENCES.	109
APPENDIX	120

LIST OF TABLES

TABLE	PAGE
1. Preliminary GSH evaluation results.	64
2. Correlation of GSH Content with cell cycle distribution.	65
3. Correlation of GSH content with cell cycle progression following culture synchronization and GSH depletion protocols.	73
4. Distribution of nicotinamide nucleotides.	77
5. Increase in $[Ca^{2+}]_i$ response to B FGF	94
6. Raw data from Ca^{2+} mobilization experiments	120
7. Statistical analyses for Ca^{2+} mobilization data . .	122

LIST OF FIGURES

FIGURES		PAGE
1.	Eyer GSH Kinetic assay standard curve	62
2.	Linear regression line generated from comparison GSH (nmol)/mg protein vs. DNA index	66
3.	Representative histograms of DNA related fluor- escence of propidium iodide stained fibroblast nuclei	68
4.	Photomicrographs taken at a 50 x magnification of human gingival fibroblasts.	70
5.	Fura-2 penta potassium salt Ca^{2+} EGTA buffer mix standard curve.	81
6.	Free Ca^{2+} ratio recordings for the Fura-2 pentapotassium salt Ca^{2+} EGTA buffer mixes. . . .	83
7.	Demonstration of functional esterase activity in fura loaded fibroblasts	85
8.	Tritiated thymidine incorporation (61 Ci/mM) vs. time in culture	88
9.	Bar histogram of cell cycle distribution in G1 at harvest	92
10.	Resumption of cell cycle progression following recovery of cellular GSH levels	93
11.	Relationship between x levels of reduced nicotin- amide nucleotides and B FGF stimulated percent increase in $[\text{Ca}^{2+}]_i$	97
12.	Schematic diagram showing proposed mechanism of thiol redox association with cell cycle progression	107

CHAPTER I

The question as to what makes a cell divide appears superficially simple. Obviously, to be capable of growth and division, cells must be supplied with adequate nutrients, growth factors and mitogens. During most non-anabolic states, the extracellular milieu found in vivo is remarkably constant. However, during normal cell cycle progression, cells are only mitogen-responsive during the G1 phase of the cell cycle. Intracellular parameters are apparently the more variable of the components that determine mitogenic responsiveness. Cells in G1 must therefore have some unique cellular composition that allows mitogenic signal transduction only during this cell cycle state.

Current cell cycle theory maintains that, in the absence of outside intervention, once a cell progresses beyond G1, it is committed to the cell cycle. It would be futile, from both cytoprotective and energy conserving standpoints, for cells to commit to the cell cycle if the chances for successful cell cycle completion were not good. Cells have evolved mechanisms for environmental assessment, and only enter the active cell cycle when external

conditions are favorable. Apparently, cells' possess equally stringent, but less well defined, means to evaluate intracellular growth potential.

The hypothesis that this study tested was whether or not cellular thiol redox status is an intracellular parameter that affects cellular mitogenic responsiveness.

Several reasons make the cellular thiol redox status an attractive component to partake in cell cycle regulation. During late G1 and S, there are marked increases in activity of two thiol dependent enzymes, DNA polymerase and ribonucleotide reductase. Cellular increases in glutathione (GSH) levels coincide with these increased enzymatic activities. Although GSH's extraribosomal synthesis is energetically costly, it appears that GSH's contributions offset the additional energy expenditure necessary for its synthesis. Further, the cellular redox status reflects the cellular bioenergetic state. Prior to engaging in an energy demanding process such as cell cycle progression, it would be appropriate to determine if adequate energy stores existed for successful completion of the cell cycle.

Literature Review

I. Eucaryotic Cell Cycle

A. Overview

A recent cell biology review article by Murray and Kirschner succinctly defined the cell cycle as "the set of

events that is responsible for the duplication of the cell." (124)

Two diverse approaches have led to the current level of understanding of the eucaryotic cell cycle. The advances made in prokaryotic genetics during the 1950's and 1960's provided the basis for the genetic approach to clarifying the cell cycle. By analyzing mutations that resulted in cell cycle arrest, the genetic approach developed a "Domino Theory" - the cell cycle is comprised of a series of interdependent biochemical reactions. (29,41,147,111,37) Addressing the question from another perspective, physiologists and embryologists evaluated cells that were specialized for rapid cell division - eggs and oocytes. Studies based upon the oocyte led to the formulation of the "Biochemical Clock" approach - the cell cycle oscillates between two states - mitosis and interphase. (37,42,104)

Despite numerous attempts to devise a unified theory regarding the cell cycle, numerous factions and theories are still prevalent. (104,124) Points of contention regarding cell cycle theory are diverse, and include such aspects as the recruitability of noncycling cells back into the cycle, and does the cell cycle function as a continuum. (104,124)

Although some aspects of the cell cycle still remain debatable, there is a consensus of opinion regarding many features of the cell cycle. Actually, the cell cycle

entails the coordination between two specific, but not necessarily interrelated cellular activities - that of cellular growth and DNA synthesis. (1,104) During each cell cycle, the cell must double its mass (inclusive of all its organelles) and also duplicate its DNA. (1,104) The regulation of this process is remarkable from several aspects. Although chromosomal duplication can usually be accomplished more rapidly than cell growth, cytokinesis does not occur until both processes are completed.

(1,104,124) Further, over many cell divisions, daughter cells retain a remarkably constant size. (1,104,124) Apparently, the presence of feedback controls in the somatic cell cycle, ensures that processes occurring during the transition from cell cycle states are completed prior to progression to the next state. (1,104,124) In addition, these controls create dependency relations in the cell cycle, and permit modulation by external factors.

(135)

The cell cycle has been divided into 4 successive phases: G1, S, G2, M. (1) (G1=gap 1, S=DNA synthesis, G2=gap 2, M=mitosis) Originally, G1 was defined as the time interval (gap) that was noted between the identifiable events of mitosis and DNA synthesis. (1,135) The interphase period of the cell cycle is composed of successive G1, S and G2 phases, with the M phase of the cycle beginning with mitosis and ending with cytokinesis.

(1,135) Evidence for the selective presence of a factor active only in mitotic cells was provided in 1970 by the cell fusion work conducted by Rao and Johnson. (145) Their results demonstrated that fusion of cells in mitosis with cells in any other state induced some sort of mitotic response in the interphase nucleus. (145) Further, when cell fusion was conducted between any 2 interphase cells, the advanced nucleus "waited" for the lagging nucleus to attain the comparable cell cycle stage prior to resuming cell cycle progression. (145)

The identification of the same key regulator of the cell cycle by Masui and Markert in 1971, and Reynhout and Smith in 1974 provided experimental evidence regarding the existence of cell cycle regulating factors. (110,149) Both investigative teams demonstrated that unfertilized eggs contained an activity in their cytoplasm (separate from progesterone) that prompted meitotic maturation in immature oocytes. (110,149) This cytoplasmic activity was appropriately named maturation promoting factor (MPF). (110,149)

Subsequent studies have revealed the broad scope of the distribution of MPF. Active MPF is found only in mitotic or meiotic cells. It is not seen during interphase, and is present from simple eucaryotes (yeast extracts) to mammalian cells. (168,167,125) MPF, which has recently been purified from frog and starfish eggs, is a protein

kinase that phosphorylates histone H1, and is most probably identical to the growth-related H1 kinase, whose appearance during mitosis has been shown for years in many organisms. (96) The catalytic portion of MPF is felt to be p34^{cdc2}, which is the homolog of the product of the yeast cell cycle regulating gene, cdc2. (96,39)

A current cell cycle theory maintains that MPF, in coordination with a specific class of ubiquitously distributed (throughout the animal kingdom) proteins, called cyclins, promotes cell cycle entry into M. (39,62) In accordance with this theory, the level of p34^{cdc2} remains constant throughout the cell cycle in cycling cells, however, cyclin must accumulate during interphase. Cyclin and p34^{cdc2} then associate to form a complex that undergoes post translational modifications (probably dephosphorylation of p34^{cdc2} and phosphorylation of cyclin) generating active MPF. (123,124) Active MPF concurrently promotes entry into M and the degradation of cyclin, which then results in the return to interphase. (123,124)

B. Importance of the G0 and G1 Phases of the Cell Cycle

The cell cycle has been further refined, to include a G0 phase, which is characterized by cells that are metabolically quiescent, and yet are readily recruitable back into the cell cycle. (4,31,34) This inducible group of cells is physiologically important, as the rate of a cell population increase is primarily determined by the

fraction of cells that are currently cycling. (4,31,34) Further, the capacity to enter this quiescent, but healthy state, provides for a cytoprotective response in the event that the cell detects suboptimal conditions, such as a lack of nutrients, or the presence of cellular contact inhibition. (4,31,34)

A state of decreased metabolism characterizes G0 cells. (4,31,34) Because of protein and RNA degradation, G0 cells decrease in size. (4,31,34) During G0, both enzymatic and transmembrane transport activities are decreased. (4,31,34) Macromolecular synthesis in G0 is approximately 1/3 as rapid as during G1, and G0 ribosomes are monosomal rather than polysomal. (4,31,34) Upon return of more favorable growth conditions, G0 cells can reenter the cell cycle. (4,31,34) However, because of their need for additional metabolism, G0 cells require more time to reach S than cycling cells, and reenter the cell cycle at the G1 phase. (4,31,34)

G1 represents the most time variable phase of the cell cycle, with variations in the duration of G1 being correlated with variations in cell cycle times. (135) In early embryos, G1 can have a negligible duration, demonstrating that the G1 events required for the initiation of DNA synthesis can begin during the prior cell cycle. (140) Therefore, the observed kinetic gap between

mitosis and DNA synthesis is dependent upon the extent of progress from the previous cell cycle. (1)

For reasons in addition to its functional role (preparation for S) G1 represents an integral cell cycle phase. (4,31,34) The cellular movements in and out of G1 are the primary determinants of the post embryonic cellular proliferation rate. (4,31,34) Further, this "recruitment" process into G1 and entrance into active cell cycling is felt to be defective in cancer (CA). (134) In the absence of outside intervention, once a cell progresses beyond G1, it is committed to completing the cell cycle, and cell cycle events become largely independent of extracellular factors once cells enter into S. (134)

The cell cycle phase G1 has been divided into subphases, which are listed in sequence (from early G1 to late G1): competency, entry, progression, and assembly. (153) The discreet categories provided by this G1 classification don't entirely reflect the abundance of conflicting information regarding G1 progression. (124) As mentioned previously, it is during the G1 phase that the agonist receptor coupling (growth factors, hormones, neurotransmitters interacting with the cell and which function to promote cell cycle progression) occurs. (4,31,34)

Very early events that are observed in competent cells (first subphase G1) include alterations in chromatin

structure, enhanced transmembrane transport, and the generation of novel mRNA's. (153) However, a point of contention lies in determination of which factors are required to induce cellular competence. (154) Current experimental results imply that the capacity to ellicit a competence response (C) from a given factor, or sequence of factors, may vary depending upon the cell's origin e.g. bombesin may be very active in promoting competence in some cells, but not in others. (154) The time requirement for competent cells to reach S is the same as G0 cells, which is approximately 12 hours for 3T3 cells. (135)

Chen and Rabinovitch recently evaluated the effects of platelet-derived growth factor (PDGF), epidermal growth factor (EGF) and insulin-like growth factor I (IGF-I) on cell cycle parameters in human diploid fibroblasts. (25) These investigators found that PDGF and EGF regulated the cycling fraction of cells, while IGF-I affected the rate of exit from G1 to S. (25) Their findings were in good agreement with previous work conducted by Campisi and Pardee in which they described a sequence of several growth control points. (20) Campisi and Pardee stated that they felt EGF and PDGF exerted their effects in early G1, while insulin/IGF-I functioned around the G1/S interphase. (20) Of interest, Chen and Rabinovitch suggested that platelet release of PDGF in vivo may be a consequence of connective

tissue injury, and the released PDGF may subsequently be a stimulus for wound repair. (25)

Two of the genes activated during early G1 are c-myc and c-fos, which are associated with cellular proliferation and cellular differentiation, respectively. (143) Chen and Rabinovitch demonstrated that both PDGF and EGF stimulated the expression of c-myc and c-fos. (25) In a related study, Kaibuchi et al. reported that PDGF and fibroblast growth factor (FGF) elevated c-myc in RNA levels in Swiss 3T3 fibroblasts. (88) Further, these authors concluded that the growth factors induction of the c-myc gene activation was attributable to calcium (Ca^{2+}) mobilization and protein kinase C activation. (88)

If competent cells are incubated in a medium containing plasma, but lacking essential amino acids, they will progress toward S to a point in the entry subphase of G1, named V. (4,31,34) Following provision of the amino acids, cells at V require approximately 6 hours to reach S. (4,31,34,134) This 6 hour interval is close to the usual duration of G1 (M to S) in cycling cells. (4,31,34) Although the cellular biochemistry of the entry subphase is not well characterized, it is known that during this interval (C to V) that the increased turnover of protein and nucleic acids permits new macromolecular synthesis. (135) During entry, there is also an increase in polysomes and glycolytic enzymes. (135)

Distinct cellular and extracellular requirements are manifested as cells enter the progression subphase of G1. Unlike the previous G1 subphases, during progression the growth factor requirement has decreased. The exclusive growth factor required for 3T3 cells to progress from V to S is IGF-I. (180) Because the progression subphase requires rapid net protein synthesis, cells at this point show both a requirement for amino acids, and a marked sensitivity to protein synthesis inhibitors. (4,31,34) Proteins vital for cell cycle progression are synthesized during this time, and include enzymes required for DNA synthesis e.g. ribonucleotide reductase (RNR), thymidine kinase (TK), and DNA pol , in addition to the important regulatory protein, cyclin. (4,31,34)

Recent work by Sherley and Kelly provides insight into how cells regulate the activity of these S phase specific enzymes that are synthesized during the progression subphase of G1. (159) According to Sherley and Kelly, there are 2 subclasses of S phase specific enzymes. (159) The cellular regulation of enzymatic synthesis and the induction of the level of enzymatic activity provided the basis for classification. (159) Enzymes such as DNA pol and topoisomerase I show a small (2 to 4 fold) amplitude of induction during S. (171) In contrast, the other subgroup was characterized by large (10 to 20 fold) amplitudes of

induction and included such enzymes as thymidine kinase (TK), RNR, and dihydrofolate reductase (DHFR). (82)

Sherley and Kelly identified two post-transcriptional regulatory mechanisms for TK. (159) The increased rate of cellular synthesis of TK was due to an increase in efficiency of TK mRNA, and not due to increased mRNA production. (159) Further, because the stability of TK rapidly decreases after cell division, Sherley and Kelly proposed that this finding suggested a stage specific degradation mechanism. (159) They speculated that the G1 specific destruction of thymidine kinase may be mediated by a G1 phase, highly specific, protease. (159) Or, alternatively, TK may render itself sensitive to proteolysis by a G1 specific structural change. (159)

Once cells progress to the assembly subphase of G1, serum and rapid protein synthesis are no longer required for the remainder of the cell cycle. (19) Although intracellular events conducted at the end of G1 are not well defined, it is during assembly that the S phase specific enzymes migrate into the nucleus and organize into the replisome. (68)

C. Mitogenic Signal Transduction

In vivo, cells are bathed in serum and its associated mitogens and growth factors; however a cellular mitogenic response is only obtained during certain specific phases of the cell cycle. (135,4,31,124) Apparently, mitogenic

signal transduction from the extracellular milieu to the cell's nucleus is a carefully regulated event.

(135,4,31,124) Intracellular conditions apparently modulate cellular mitogenic responsiveness.

A group of small proteins, the purified growth factors, are the most frequently studied agents that stimulate cell proliferation. Because of the large volume of growth factor studies that have been conducted, there is an abundance of experimental findings. (135,4,31,124) Understandably, these data are not always in agreement, and therefore this affects the ability to make generalized conclusions. (135,4,31,124) Well defined responses that are noted in a wide variety of cells after mitogen addition include ionic signalling, the activation of specific genes, and a general metabolic stimulation. Areas that remain more nebulous, concern which specific signalling pathways are activated by which growth factors, and are the effects of combined growth factors synergistic or merely additive. (71,151,133,55) Some of the literature dealing with the more controversial aspects of growth factor responses will be discussed. (71,151,133,55)

Hesketh et al. conducted an extensive study to evaluate single cell (3T3 fibroblasts) responses to the sequential additions of mitogens. (71) Upon mitogenic stimulation they noted a common cellular response - which was to increase both the intracellular $[Ca^{2+}]$ ($[Ca^{2+}]_i$) and

intracellular pH (pH_i). (71) Further, the cellular Ca^{2+} response was either "all or none." (71) If a cell was refractory to one stimulus, it remained refractory to all other mitogens. (71) Only EGF caused a $[\text{Ca}^{2+}]_i$ response that was dependent upon Ca^{2+} in the extracellular medium. (71) Hesketh et al.'s data showed that there is no simple correlation between early cellular responses (Ca^{2+} and pH) and subsequent DNA synthesis. (71) The authors concluded that ionic signals are not sufficient to commit cells to S phase. (71)

A study by Ross and Ballard reported findings that refute the need for interplay between competence (e.g. FGF) and progression (e.g. IGF-I) factors to initiate DNA synthesis. (151) These authors determined that in BHK-21 cells, FGF alone was sufficient to promote ^3H incorporation at levels that were 70% of a serum induced ^3H thymidine incorporation. (151) Ross and Ballard concluded that in order to attain a maximal growth rate, more than one factor may be necessary. (151) Their results did show that the presence of serum produced the most sustained anabolic response. (151)

Pandiella et al. studied the plasma membrane hyperpolarizations and $[\text{Ca}^{2+}]_i$ increases in NIH-3T3 fibroblasts in response to various mitogens. (133) It was determined that the mitogen-induced increases in $[\text{Ca}^{2+}]$ were paralleled by a plasma membrane hyperpolarization.

(133) As a K^+ efflux was primarily responsible for this hyperpolarization, these investigators found that by progressively increasing the extracellular $[K^+]$, they were able to reverse the membrane hyperpolarizing effects of PDGF and FGF. (133) The cellular responses noted with PDGF and FGF were much slower than those induced by bradykinin, extracellular ATP, or EGF. (133) The authors concluded that the similarities between PDGF and FGF induced early signals (time course and insensitivity to phorbol esters) suggested similar transmembrane signalling. (133) Further, they noted that the variation in time course responses between EGF (very rapid, 16-20 sec) and PDGF/FGF (30 sec to 3 minutes) implied that there was a difference in coupling mechanisms at the corresponding receptors, that may involve tyrosine kinase phosphorylation sites. (133)

Gonzalez et al. conducted a study to evaluate the effects of EGF, serum, and nucleotides on cytosolic free Ca^{2+} in single A431 cells. (55) In agreement with Hesketh et al., these investigators noted that EGF elicited increases in $[Ca^{2+}]_i$ were dependent upon extracellular Ca^{2+} . (55) Unlike Hesketh or Pandiella, they also found that the EGF response was heterogeneous, both in regard to time to respond (10-30 sec) and cellular extent of response (did not see on all or none phenomenon). (55) No lag or heterogeneity was noted in cellular

responses to fetal calf serum or ATP or UTP. (55)
Further, when cells were incubated in Ca^{2+} free medium,
there was no second response to any mitogen. (55) The
authors attributed this finding to a Ca^{2+} pool depletion.
(135,4,31,124)

As can be appreciated from both the cited and noncited
literature, cellular mitogenic signalling pathways are
fascinating, but are by no means, clarified.
(135,4,31,124,71,133,55)

Numerous authors have speculated as to the routes used
intracellularly to transmit mitogenic signals. (135) In
their signal transduction, growth factors combine with
their specific receptors, which are proteins that span the
plasma membrane. (13) From here, mitogenic signalling
pathways become divergent, but converge at a common end
point which is the activation of a cascade of kinases.
(181) Growth factors such as EGF cause their receptor
molecules to dimerize upon binding. (181) The
dimerization approximates the receptor's intracellular
domains; with their subsequent proximity then resulting in
autophosphorylation by the receptor's tyrosine kinase.
(181) Tyrosines, serines and threonines on other proteins
may then be phosphorylated, thereby further transmitting
the signal. (154)

Other growth factors trigger the kinase cascade by a
less direct mechanisms. (154) Upon binding, FGF activates

phospholipase C, resulting in the hydrolysis of phospholipids, generating diacylglycerol (DAG) and inositol phosphates. (154) In the presence of increased $[Ca^{2+}]_i$, DAG activates protein kinase C, while inositol phospholipids promote Ca^{2+} mobilization from both intracellular and extracellular stores. (78,85)

There is currently some debate regarding which inositol products are most physiologically active, and what functions these products perform. (83,84,85) The most thoroughly studied is the 1, 4, 5 inositol triphosphate (1, 4, 5 Ins P_3), which has been shown to be active in mobilizing Ca^{2+} from both intracellular and extracellular sites. (83,84,85) Joseph and Williamson recently published a review in which they discussed inositol phospholipids and Ca^{2+} release. (85) They stated that in most cells, 1, 4, 5 Ins P_3 is capable of releasing only a fraction (20-40%) of non-mitochondrial Ca^{2+} stores. The sites of Ca^{2+} release were related to the endoplasmic reticulum and the plasma membrane. Volpe et al. proposed that the responsive Ca^{2+} pool may be present in discreet subcellular structures, termed "calciosomes". (174) These authors also discussed factors limiting the efficacy of 1, 4, 5 Ins P_3 , which included the restricted diffusion of such a highly charged molecule, and the presence of soluble and membrane bound phosphatases. A major conclusion made by Joseph and Williamson was that 1, 3, 4,

5 Ins P_4 has a variable ability to mobilize Ca^{2+} , but is much less potent than 1, 4, 5 Ins P_3 in this regard. (85) This conclusion by Joseph and Williamson regarding inositol product efficacy is interesting in light of the recent publication by Hill et al. (72) Hill et al. determined that 1, 3, 4, 5 Ins P_4 stimulated the initiation of DNA synthesis in Ca^{2+} deprived rat liver cells. (72) Further, because 1, 3, 4, 5 Ins P_4 has been shown to promote Ca^{2+} sequestration, Hill et al. proposed that 1, 3, 4, 5 Ins P_4 may function as a means of inactivating the Ca^{2+} mobilization due to 1, 4, 5 Ins P_3 . (72)

A common theme prevailing during the different mitogenic signalling pathways is Ca^{2+} mobilization, coinciding with the activation of the kinase cascade. (154,135,181) There is extensive experimental evidence showing that Ca^{2+} is vital for cell proliferation. (115) Hesketh et al. conducted early studies in which they demonstrated that the Ca^{2+} ionophore A23187 stimulated DNA synthesis in human and pig lymphocytes. (69) Boynton et al. have described that there are two extracellular Ca^{2+} dependent stages of G1. (12) The first extracellular Ca^{2+} dependent stage was found to occur during the initial 30 minutes of mitogen stimulation, with the 2nd stage determined to occur 8 to 10 hours later at a point immediately prior to chromosome replication. (12) These investigators found that the addition of 1.25 mM

extracellular Ca^{2+} to the medium during either of these Ca^{2+} dependent stages stimulated low Ca^{2+} /blocked cells to progress through G1. (12) Of interest, many tumor cell lines have a lower requirement for extracellular Ca^{2+} , and will progress through the cell cycle at extracellular $[\text{Ca}^{2+}]$ from 1 - 10 u M. (12)

One of the cellular systems activated by Ca^{2+} mobilization is the Na^+/H^+ antiporter. (129) In a well designed study, Ober and Pardie evaluated the routes of activation of the Na^+/H^+ antiporter in Chinese hamster embryo fibroblasts. (129) They noted that a rapid stimulation of an amiloride sensitive Na^+/H^+ antiporter was a common response of quiescent cells to growth factors. (129) They proposed that the cells were using two methods to increase the pH_i , but both methods entailed an influx of Na^+ and an efflux of H^+ . The first route involved protein kinase C and DAG, while the second mechanism was apparently Ca^{2+} mediated. Their results demonstrated that functional protein kinase C is necessary for thrombin induced increases in pH_i . In contrast, EGF employed a Ca^{2+} dependent route for antiporter activation. The authors concluded that in vivo EGF and thrombin might each act by both protein kinase C and Ca^{2+} /calmodulin dependent pathways; with thrombin relying primarily on protein kinase C, and EGF more dependent on Ca^{2+} . (129)

Calcium's diverse role in cellular proliferation can be further appreciated through a study published by Chin et al. (26) These investigators proposed that Ca^{2+} regulates translation in eucaryotic cells through the modulation of the rate of initiation rather than through polypeptide chain elongation or termination. (26) Previous reports have shown that the major support processes for protein synthesis were not perturbed by Ca^{2+} depletion e.g. maintenance ATP or GTP concentrations, or protein catabolism. (26) Chin et al. found that the reintroduction of Ca^{2+} promoted the accumulation of the 43 S ribosomal preinitiation complex, and allowed its subsequent combination with a 60 S subunit to form the 80 S monosome. Further, they noted a rapid formation of a methionylated complex in response to Ca^{2+} induced changes originating in the endoplasmic reticulum. These Ca^{2+} induced alterations are ultimately responsible for stimulating the initiation of protein synthesis. (26) The plausibility of the mechanism is supported by the requirement for rapid protein synthesis during G1, which is a time in the cell cycle where mitogen induced Ca^{2+} mobilization would have occurred. (135)

II. Cellular Thiol Redox Status

A. Glutathione (GSH)

GSH (glutamylcysteinylglycine) was isolated and named by the British biochemist Frederick Gowland Hopkins in

1921. (76) The chemical analysis to determine GSH's structure was completed in 1935 by Harington and Mead.

(113) GSH represents the primary intracellular free thiol, is usually found intracellularly in high (0.1-10mM) levels, and often comprises over 90% of the total cellular nonprotein sulfur. (113) GSH is ubiquitous in its distribution, from both an intracellular (located throughout the cell) and biological (found in animals, plants and microorganisms) contexts. (94)

The GSH status of cells is defined by the total cellular concentration of GSH in all of its possible forms. (94) Intracellular forms for GSH include: GSH itself, the disulfide, GSSG, mixed disulfides (mostly GSS-protein), thiol esters, and GSH derivatives linked through non-sulfur bonds. (94) In their review, Kosower and Kosower described the cellular GSH status not as a constant, fixed value, but as a dynamic system that was capable of altering GSH distributions in response to perturbations in the cell. (94)

GSH is synthesized extraribosomally, by the consecutive actions of γ -glutamylcysteine synthetase and GSH synthetase. (113) Both enzymatic steps require ATP. The 1st enzyme, γ -glutamylcysteine synthetase, is feedback inhibited by GSH. (113) Because extraribosomal peptide synthesis is rare in eucaryotes, GSH's cellular contributions apparently offset the additional energy

expenditure required for its synthesis. (113) Two structural features of GSH, the glutamyl linkage and the -SH are responsible for its structural stability and are also closely associated with GSH's functions. (113) Throughout evolution, GSH has been adapted to perform many diverse cellular roles that include: detoxification of xenobiotics, inactivation of reactive oxygen species (ROS), maintenance of protein thiol status by disulfide interchange, functioning as an enzymatic cofactor, and participation in reactions involving the synthesis of proteins and nucleic acids. (113,92,93,94)

Four specific enzyme systems of which GSH is a component are the GSH peroxidases, GSH transhydrogenases (thiol transferases) GSH S-transferases and GSSG reductase. Both the GSH peroxidases and GSH-S transferases function in cytoprotective roles, while the GSH reductase restores cellular levels of GSH. (113)

During cellular metabolism, the potentially damaging oxidants, superoxide and H_2O_2 are generated. (24) The GSH peroxidases use GSH to reduce these ROS, thereby protecting cellular proteins and membranes from oxidation.

(94,95,113) In addition, the GSH peroxidases function in the arachidonate cascade, partaking in leukotriene generation and the regulation of prostacyclin synthesis. (113,52) There are both selenium dependent and selenium independent GSH peroxidases. (113,15) In the selenium

containing enzymes, the active site (felt to be at selenocysteine) interacts with the substrate, which is followed by reaction of the selenocysteine-substrate product with GSH. (113) The GSH transhydrogenases function in those metabolic processes that require thiol disulfide exchange, which include: protein synthesis and degradation, regulation of enzymatic activity, reduction of cystine, and synthesis of the deoxyribose intermediates for synthesis of DNA. (113)

Various exogenous compounds with an electrophilic center can conjugate with GSH. (94,113) Introduction of an electrophilic center may be the result of a previous reaction e.g. formation of an epoxide by a microsomal oxygenase. (94,113) GSH-electrophilic compound reactions may occur spontaneously, or be catalyzed by GSH S-transferases. (81) The majority of GSH conjugates are converted to mercapturic acids, which are then ultimately excreted from the cell. (81,113)

GSH reductase is a broadly distributed flavoprotein that catalyzes the reaction:

GSSG reductase

Because this reaction is essentially irreversible, the ratio of GSH:GSSG is very high in cells that have the NADPH

reducing equivalents available. (113) Other GSSG reductase substrates include mixed disulfides generated between GSH and glutamylcysteine or coenzyme A. (21) Of significance, GSSG reductase closely links cellular GSH status with the cellular redox state. (94,113)

B. GSH and the Cell Cycle

Rapkine, in 1931, used sea urchin eggs to evaluate the role of thiols in cell growth. (146) He determined that changes in the trichloroacetic acid soluble thiol content correlated with the cell division cycle. Lowest levels of TCA soluble thiols were found post fertilization, with peaks determined to occur just prior to, and at, cleavage. (145) Subsequent studies: Sakai and Dan, 1959. Dan, 1966; Neufeld and Mazia, 1957; and Mano, 1971. confirmed that there were cyclic variations in the TCA (and KCl) soluble thiol fractions. (155,32,126,107) Harris and Patt, 1969; Harris and Tang, 1973; Krammer, 1975; determined that the nonprotein thiol content increases during cell cycle progression from G1 to S in mammalian cells. (66,67,95) In addition, Cernock and Weinbergova, 1969; showed parallel increases in cellular GSH levels with mitotic activity in regenerating rat liver. (22)

More recently, Principe et al. reported a study in which they evaluated protein thiols during the cell cycle. (142) This group reported, that in tumorigenic rat cell lines, neither the amount and distribution of protein

thiols, nor cell cycle progression was affected by buthionine sulfoximine (BSO) treatment. The lack of discernable alteration in protein thiols is not surprising in light of the generalized nature of the o-phthalaldehyde reaction and the tremendous cellular protein thiol reserve. (150,156) Further, by limiting their GSH depletion to BSO, these investigators only inhibited the de novo synthesis of GSH, and did not alter existing GSH levels. (60)

Numerous recent studies have been conducted to further clarify the association between GSH and cellular proliferation. (139,99,157,51) In 1983, Post et al. reported that cellular levels of GSH were greatly affected by culture conditions. (139) Cellular GSH levels were most dependent upon the presence of serum; cells cultured with serum showed a sharp increase in GSH levels up to 24 hours after passage. Cellular GSH levels then decreased after 24 hours in non-passed, serum containing cultures. (139)

Lee et al. reported in 1988 that GSH levels in human and rodent tumor cells increased in proportion to cell volume during cell cycle progression. (99) However, these investigators found that the apparent cell cycle related changes in GSH levels dissipated when GSH levels were normalized in accordance with cell volume. (99) Of importance, it has been shown that in neoplastic cells,

both cellular thiol content and patterns of cell cycle progression deviate from nontransformed cells. (135)

In 1986, Shaw and Chou reported an association between 3T3 fibroblast GSH content and mitogenic stimulation. Fibroblast GSH levels were diminished by a combined treatment using both BSO, which inhibited synthesis, and 2 CHX-1-one, which depleted existing GSH levels. Results of this study showed that GSH depletion decreased by 22% the fibroblasts' capability to incorporate ^3H -thymidine. (157)

Fidelus et al., 1987; conducted a study in murine splenic lymphocytes in which they augmented GSH levels with 2-mercaptoethanol (2 ME) and 2-L-oxothiazolidine-4-carboxylate (OTC), and also decreased cellular GSH with BSO. (51) It was determined that lymphocytes responded to the 2 ME and OTC protocols with enhanced GSH levels. Further the "GSH augmented" cultures showed increased levels of proliferation and activation. (51) BSO treated cells had decreased levels of GSH, and showed an inhibition of cellular proliferation, but normal levels of activation. These investigators also reported a positive linear relationship between lymphocyte GSH levels and ^3H -thymidine incorporation. (51)

C. Association of GSH with Macromolecular Synthesis

Cellular protein synthesis has been shown to be affected by cellular GSH status. (92,93,94) The early protein synthesis inhibition studies reported by Kosower

and Kosower in 1972 and 1974 were conducted on diamide treated reticulocytes. (92,93) In the majority of cells, the GSH:GSSG ratio greatly favors the reduced, GSH form. (94) However, after exposure to diamide, most of the GSH pool is oxidized to GSSG. (93) In diamide treated reticulocytes, both the initiation and elongation steps of protein synthesis were inhibited. (92,93) Polypeptide chain elongation was found to resume after partial recovery of GSH. The initiation stage was the more sensitive towards altered GSH status. (93) Only upon essentially complete regeneration of GSH did the initiation of polypeptide chains resume. (92,93) Balkow et al., 1975; and Levin et al., 1976; speculated that GSSG may be affecting protein synthesis by activating a protein kinase that phosphorylates, and thereby inactivates, an initiation factor necessary for the initiation complex. (3,103) In 1978 and 1979 Ernst et al. identified the heme regulated eIF-2 kinase (HRI) that phosphorylates the 38K Da - subunit of the reticulocyte initiation factor, eIF-2. (45,46) Ernst et al. also demonstrated that the HRI induced inhibition was reversed by the addition of high concentrations of nonphosphorylated eIF-2, or 8mM concentrations of cAMP. (46) Ernst et al., 1978; Lenz et al., 1978; West et al., 1979; Michelson et al., 1984; Jackson et al., 1983; demonstrated that in gel filtered lysates, the HRI induced inhibition of initiation of

protein synthesis was prevented or reversed by physiological levels of glucose -6-phosphate, or 2-deoxyglucose-6-phosphate (0.02-1mM). (45,102,176,117,79) The sugar phosphate participates in two independent functions in polypeptide chain initiation: NADPH generation, and as a cofactor at an undetermined step in initiation. To prevent GSSG induced HRI activation, both of these sugar phosphate functions are necessary. (45,46,79)

In 1988, Kan et al. reported a study in which they monitored the activity of reversing factor (RF), which is a multipolypeptide that participates in the recycling of eIF-2. (90) It was determined that RF was inactivated in GSSG treated reticulocyte lysates. (90) Kan et al. speculated that RF recovery and the resumption of protein initiation may be dependent on the restoration of the thiol groups on RF and the dephosphorylation of the 15 S RF subunit. (90) These authors concluded that the recovery of protein synthesis in GSSG treated lysates was dependent upon, and proportional to the extent of RF recovery. (90) The findings by Kan et al. were in good agreement with a study published in 1986 by Dholakia et al. (36) In their paper, Dholakia et al. reported that RF contains bound NADPH, and that increases in levels of NAD(P)^+ inhibited RF activity. (36) These findings helped substantiate that there is an association between cellular thiol redox status and

cellular processes integral for cell cycle progression.

(36)

In their 1986 study, Shaw & Chou reported biphasic increases, during late G1 and S, in log growth fibroblast GSH levels. (157) These cell cycle times, late G1 and S, coincide with a marked enhanced activity of two thiol dependent DNA synthesizing enzymes, DNA pol (pol) and RNR. (43,122,132)

Pol is a component enzyme of the DNA replicase complex. (28,53) Because pol requires an intact thiol group for activity, in vitro DNA synthesis assays include mM levels of dithiothreitol, to ensure thiol reduction. (28,53) In 1984, Ottinger and Hubscher provided further evidence to support the importance of intact thiols in DNA synthesis. (132) They showed that all four segments of the DNA replicase holoenzyme were extremely sensitive to the thiol alkylator, N-ethylmaleimide. (132)

In 1950, Hammerston and Reichard reported that deoxyribonucleotides are generated by direct reduction of ribonucleotides. (63) These investigators also isolated some of the DNA synthesizing enzymes: RNR, thioredoxin and thioredoxin reductase. (63) RNR catalyzes the first unique step in DNA synthesis. (75) Because deoxyribonucleotides are generated by RNR's reduction of the 2'OH of the ribose sugar, RNR is dependent upon a source of reducing equivalents. (75) Holmgren, 1979; and

1981; characterized how GSH facilitates the activity of RNR. (73,74,75) Both thioredoxin and glutaredoxin can provide RNR with its necessary reducing equivalents.

(73,74,75) GSH can help maintain the function of thioredoxin (as it is thiol dependent), and as glutaredoxin, GSH is the $H^\bullet$ donor. (73,74,75) Of importance, by the activity of GSSG reductase, the cellular GSH status is interrelated with the cellular redox state. (94,113)

Erickson et al., 1984; and Engstrom et al., 1985; reported that mammalian RNR is composed of two subunits, M1 and M2. (44,43) The M1 RNR subunit contains the nucleotide binding site, while the M2 subunit contains non heme Fe, and is responsible for the generation of a tyrosyl free radical. (43,44) EPR spectroscopy has shown that this tyrosyl free radical has a half life of approximately 10 minutes. (43,44) Further, the presence of non heme Fe introduces the possibility for Fenton chemistry. (30) Therefore, very reactive e.g. $OH^\bullet$ potentially mutagenic to cellular DNA, compounds could be generated. (30,23)

However, the findings by Mbemba et al., 1985; and Shaw and Chou, 1986; keep open the possibility for a second role for GSH in DNA synthesis - that of cytoprotection. (112,157) Mbemba et al. reported on the ubiquitous cellular distribution of GSH; both free GSH and GSH/glutaredoxin were located in cellular nuclei. (112)

Further, the timing of the Shaw and Chou reported increases in cellular GSH levels (late G1 and S) corresponds to the cell cycle times of maximal RNR activity. (43,44)

Therefore, GSH, in association with the GSH peroxidases, may also act to decrease the potential for mutagenesis during DNA synthesis. (94,113)

III. Association of Cellular Thiol Redox Status with Ca^{2+} Mobilization

A. Intracellular Ca^{2+} Homeostasis and Ca^{2+} Mobilization

The intracellular free Ca^{2+} is maintained in the range of 10^{-7}M . (78) The "ultimate control" over $[\text{Ca}^{2+}]_i$ is exerted by the plasma membrane, because only the plasma membrane maintains this 10^4 gradient, and has access to the extracellular Ca^{2+} reserves. (78) The total intracellular Ca^{2+} is much greater than 10^{-7}M ; however this Ca^{2+} is bound to proteins, membranes, or intracellular organelles e.g. mitochondria, endoplasmic reticulum (e.r.), golgi apparatus, or nuclei. (78) The Ca^{2+} dependent enzymes are regulated by the intracellular free Ca^{2+} pool, and are responsive to the transient increases in $[\text{Ca}^{2+}]_i$. (78)

In 1980, Stolze and Schulz, and in 1981, Tan and Tashjian, reported that for short response assays, intracellular organelles are adequate to regulate $[\text{Ca}^{2+}]_i$. (166,169) However, to induce a second, or more extended Ca^{2+} response the cell was dependent on the presence of extracellular Ca^{2+} . (166,169) Both of these studies

noted that when the agonist responsive pool was being refilled, the $[Ca^{2+}]_i$ did not rise. (166,169) This finding implied that the Ca^{2+} responsive pool communicates with the extracellular fluid. (166,169,179)

During the 1970's and early 1980's, a debate ensued regarding which intracellular organelle provided the source of Ca^{2+} during mobilization. (17) Early studies had focused on the mitochondria; results of this work had shown that mitochondria can sequester Ca^{2+} in an ATP dependent process. (158) It was also determined that in some tissues e.g. liver, the mitochondria did contain the major Ca^{2+} stores. (158)

However, experiments conducted during the 1980's supported the e.r. as the intracellular Ca^{2+} regulator. (158,6,14,165,162) Somylo et al. observed that in hepatocytes previously loaded with Ca^{2+} , the mitochondria contained the primary Ca^{2+} store. (162) But, when the hepatocytes were suspended in physiological levels of Ca^{2+} , the e.r. possessed most of the stored Ca^{2+} . (162) Evidence that the mitochondria were responding to, and not initiating the increases in $[Ca^{2+}]_i$ was provided by Shears et al. in 1984. (158) Using tissues that had been challenged to release Ca^{2+} , then rapidly fractionated, Shears et al. found that the mitochondrial Ca^{2+} levels had increased. (158) Studies were then conducted by Streb and Schulz on permeabilized cell preparations that were treated

with selective inhibition of either mitochondrial or e.r. Ca^{2+} uptake. (164) These investigators found that the e.r. is the primary organelle used in intracellular Ca^{2+} buffering. (164) They also noted that the mitochondrial affinity for Ca^{2+} is lower than that of the e.r.; mitochondria will load Ca^{2+} when the $[\text{Ca}^{2+}]_i$ rises to the μM range. (164) Other evidence in favor of the e.r. is that it is this organelle that is the primary target of 1, 4, 5 Ins P_3 . In 1986, Spat et al. reported specific binding of radiolabelled 1, 4, 5 Ins P_3 to m icrosomal membranes. (163)

Although no longer considered as the primary Ca^{2+} regulators, mitochondria are essential in maintaining Ca^{2+} homeostasis. (78) Because of their "insatiable appetite" for Ca^{2+} , the mitochondria readily respond to increases in $[\text{Ca}^{2+}]_i$. (78) Denton and McCormack proposed that an underlying reason for the mitochondrial Ca^{2+} uptake is that specific intra-mitochondrial enzymes e.g. pyruvate dehydrogenase, and NAD linked isocitrate dehydrogenase, are physiologically regulated by Ca^{2+} . (35)

In 1975, Mitchell noted that specific agonists that caused inositide degradation also promoted an increase in $[\text{Ca}^{2+}]_i$. (118) Inositide turnover is initiated following activation of a phosphodiesterase upon receptor binding. (78) The phosphatidyl inositol (4,5) bisphosphate is then

hydrolyzed into diacylglycerol (DAG) and 1, 4, 5 Ins P_3 .

(78)

Investigations conducted by Berridge in 1983, and Berridge et al. in 1984, showed that 1, 4, 5 Ins P_3 represented a primary inositide degradatory product and was not generated secondarily to Ca^{2+} mobilization. (8,11) Berridge proposed that 1, 4, 5 Ins P_3 was the "missing link" that connected the receptor mediated inositide turnover with intracellular Ca^{2+} stores. (8,9,10,11) A series of experiments conducted by Streb et al., using permeabilized cells and subcellular fractions, confirmed that the e.r. is the target for 1, 4, 5 Ins P_3 . (165) The specific binding of the radiolabelled 1, 4, 5 Ins P_3 to microsomal membranes reported by Spat et al. confirmed the presence of a specific receptor. (163) In 1985, Volpe et al. and Prentki et al. reported findings that showed that not all of the e.r. Ca^{2+} pool is released upon stimulation. (173,141) The responsive, trigger pool of Ca^{2+} was proposed to be contained in discrete structures within the e.r., which Volpe et al. called "calciosomes". (174) The work by Volpe et al. and Prentki et al. suggested that there is a specific portion of the e.r. that contains the 1, 4, 5 Ins P_3 receptors, and that these receptors were probably close to the membrane. (173,141) The receptor proximity to the membrane is in good agreement with the conclusions of Joseph and Williamson regarding the

difficulty of a highly charged molecule, like 1, 4, 5 Ins P_3 , travelling a long distance. (85)

Streb et al. and Muallam et al. conducted studies that further clarified the mechanism of action of the 1, 4, 5 Ins P_3 initiated Ca^{2+} release. (165,121) Streb et al reported that the e.r. Ca^{2+} release triggered by 1, 4, 5 Ins P_3 is both rapid and catalytic. (165) Muallam et al. determined that the 1, 4, 5 Ins P_3 initiated release opens a Ca^{2+} conductance which releases Ca^{2+} in exchange for another cation, felt most probably to be K^+ . (121) Smith et al. concluded that because of its temperature dependence, that this Ca^{2+} transport is more likely a pore as opposed to a carrier. (161) Muallam et al. also found that the uptake of Ca^{2+} into the e.r. required ATP hydrolysis and the co-transport of an anion, but the Ca^{2+} release from the e.r. was not energy dependent. (121)

In 1986, Dawson et al. reported that guanosine triphosphate (GTP), in the presence of polyethylene glycol, could release Ca^{2+} and augment the Ca^{2+} mobilization of 1, 4, 5 Ins P_3 in liver microsomes. (33) Dawson et al. concluded that for the 1, 4, 5 Ins P_3 - Ca^{2+} system to be active, a GTP mediated protein phosphorylation needed to occur. (33)

Joseph et al. recently conducted studies to further characterize the GTP - mediated microsomal Ca^{2+} release. (84) It was found that in the presence of polyethylene

glycol, GTP reversibly increased the permeability of a transmembrane "pore" in microsomal membranes. (84) The authors suggested that the GTP responsive "pore" may link 1, 4, 5 Ins P_3 sensitive sites to other Ca^{2+} containing compartments. (84)

The catabolism of 1, 4, 5 Ins P_3 is conducted by a 1, 4, 5 Ins P_3 phosphatase that was discovered and characterized by Downes et al. (38) These authors noted similarities between another 2nd messenger, cAMP, and 1, 4, 5 Ins P_3 . (38) Both of these second messengers have specific and potent deactivation mechanisms. (38) And both cAMP phosphodiesterase and 1, 4, 5 Ins P_3 phosphatase are located within the plasma membrane when they are membrane bound. (38)

Investigation of cellular Ca^{2+} regulation and mechanisms of Ca^{2+} mobilization has continued to be a prime research area.

B. Cellular Thiol Redox Status and Ca^{2+} Sequestration

Because mitochondria were known to be active in respiration-coupled uptake of Ca^{2+} , they were used in many of the early studies that evaluated Ca^{2+} regulation. (152,100,16) In 1964, Rossi and Lehninger reported that the addition of Mg^{2+} and ATP prevented the uncoupling effect of high extra-mitochondrial Ca^{2+} on respiring mitochondria. (152) Subsequent studies continued to pursue the mechanism responsible for the mitochondrial

damage induced by high Ca^{2+} . (40,77,5) Drahota et al, 1965, reported that incorporation of ATP and increased levels of protein delayed the Ca^{2+} release of Ca^{2+} loaded mitochondria. (152) The Drahota et al. results inferred that the mitochondrial damage was not detectable until the Ca^{2+} release began. (40) In 1976, Hunter et al. reported that during the Ca^{2+} damaged state, mitochondria show a non-specific increase in permeability; e.g. sucrose was shown to enter the matrix. (77) Further, in 1977 Baum and Al-Shaikhaly showed that the mitochondrial swelling associated with high Ca^{2+} was of colloidal osmotic origin and was preventable by the addition of colloid to the medium. (5)

In 1978, Lehninger et al. first reported an association between the steady state redox poise of mitochondrial pyridine nucleotides and their retention and release of Ca^{2+} . (101) This study used mitochondria that were isolated from normal rat heart and liver and provided succinate as the energy source. (101) Rotenone was added to alleviate net electron flow from the endogenous pyridine nucleotides to O_2 . As long as the pyridine nucleotides remained reduced, the mitochondria showed rapid uptake and retention of Ca^{2+} . Exposure to oxidants e.g. acetoacetate, caused a mitochondrial Ca^{2+} release, while addition of reductants, e.g. isocitrate, promoted a re-uptake of the released Ca^{2+} . (101) Alteration of the mitochondrial

pyridine nucleotides to more oxidized or reduced states caused successive cycles of Ca^{2+} release and re-uptake. Replacement of succinate with ascorbate or ATP provided similar results. Lehninger et al. proposed that the pyridine nucleotide redox state is a component of a feedback regulation of mitochondrial enzymes. When mitochondrial pyridine nucleotides are in a reduced state, Ca^{2+} is retained in the mitochondria and is available to stimulate the Ca^{2+} dependent enzymes. (101)

Following the 1978 report by Lehninger et al., subsequent studies were conducted to further define the redox state and Ca^{2+} retention association. Lotscher et al., 1979; investigated the effects of hydroperoxides on the redox state of rat liver mitochondrial pyridine nucleotides. (105) Their experimental design applied the finding that feeding of a selenium deficient diet markedly decreases rat liver mitochondrial GSH peroxidase (GPO) activity. After exposure to H_2O_2 or t-butyl hydroperoxide rat liver mitochondria with functional GPO concomitantly lost Ca^{2+} and oxidized NAD(P)H. Mitochondria isolated from selenium deficient animals did not oxidize NADPH or release Ca^{2+} . Lotscher et al. hypothesized that the redox poise of mitochondrial pyridine nucleotides was at least partially regulated by the activities of GSH reductase and GPO. These findings introduced cellular thiol status into the area of Ca^{2+} regulation. (105)

The contribution of cellular thiols on rat heart mitochondrial Ca^{2+} retention was further explored by Harris et al. (65) Thiol alkylating agents, e.g. N-ethyl-ethylmaleimide, stimulated Ca^{2+} efflux; whereas thiol reductants e.g. dithiothreitol, slowed the Ca^{2+} release. This study also noted that the addition of ADP and Mg^{2+} decreased mitochondrial Ca^{2+} efflux. (65) Harris et al. concluded that mitochondrial membrane permeability is at least partially controlled by the binding of ADP and Mg^{2+} on the inner membrane. (65) Reduction of certain thiols, by NADPH and an ATP dependent transhydrogenase, was felt to be a factor in the ADP Mg^{2+} binding to the inner membrane. (65)

Glucagon, which had been shown to augment mitochondrial Ca^{2+} retention, was used in the study conducted by Prpic and Bygrave. (144) Glucagon exposed rat liver mitochondria showed both a slower rate of NADPH oxidation and Ca^{2+} efflux than the saline treated controls when exposed to oxalacetate. (144) The authors proposed that glucagon, by stimulation of the ATP linked transhydrogenase, kept the mitochondrial pyridine nucleotides in a reduced, therefore more Ca^{2+} retentive, state. (144)

Jung and Brierley, 1981; and 1982; reported that the redox state of mitochondrial pyridine nucleotides affected the efflux of another cation, K^+ . (86,87) Results from

the study published in 1981 showed that the mechanism for K^+ efflux from uncoupled mitochondria differs from Ca^{2+} efflux. (86) Mitochondrial Ca^{2+} efflux had been shown to occur following an uncoupler induced loss of the protonmotive force. (86) For K^+ efflux from uncoupled mitochondria to occur, oxidation of mitochondrial NADPH needed to be accompanied by either: an elevation of matrix phosphate prior to the introduction of an uncoupler, or ethylene glycol bis (B-amino-ethyl ether) - N, N, N', N' - tetracetic acid (EGTA), or ruthenium red added in addition to the uncoupler. The authors concluded that neither of these K^+ releasing conditions appeared to use the K^+/H^+ exchanger that is present in mitochondria that have maintained their membrane potential. (86) In 1982, Jung and Brierley reported findings that clarified the association between K^+ efflux and the redox state of pyridine nucleotides in uncoupled mitochondria. (87) Their results implied that mitochondria have a latent pathway for K^+ permeability. The closure and opening of this pathway was proposed to be reversible, and was associated with the mitochondrial pyridine nucleotides redox poise, or a group in close communication, e.g. a membrane bound thiol. (87)

Despite many investigations, the mechanism of mitochondrial Ca^{2+} efflux is still controversial. In 1988, Vercesi et al. used deenergized mitochondria with oxidized

pyridine nucleotides to study Ca^{2+} induced mitochondrial damage. (172) De-energized mitochondrial Ca^{2+} induced swelling was increased by ionophores and decreased by Mg^{2+} and ATP, and ruthenium red. (172) The authors concluded that Ca^{2+} cycling is not an essential feature for Ca^{2+} induced damage. Also, for the mitochondrial damage to occur, the Ca^{2+} must be contained within the mitochondria. (172)

Orrenius and colleagues have applied a slightly different approach to the study of Ca^{2+} sequestration by using intact cells and microsomes in addition to mitochondria. (7,170) In 1982, Bellomo et al. reported that rat hepatocytes contained two pools, a mitochondrial and an extramitochondrial, (e.r.) for Ca^{2+} compartmentation. (7) Addition of t-butylhydroperoxide caused Ca^{2+} efflux from both Ca^{2+} pools, along with oxidation of NAD(P)H and GSH. Inhibition of GSH reductase reduced the mitochondrial Ca^{2+} efflux while it increased the e.r. Ca^{2+} release. The authors suggested that different regulatory mechanisms may modulate the two Ca^{2+} pools. The mitochondrial Ca^{2+} pool was proposed to be more dependent on NAD(P)H, while the e.r. pool regulation was more thiol dependent. Because of the GSH reductase dependency upon NAD(P)H, these two parameters are interdependent. (7) In 1985, Thor et al. reported results from a study which investigated the role of

microsomal thiol groups in Ca^{2+} sequestration. (170)

Specifically, the study evaluated the ATP dependent Ca^{2+} sequestration by rat liver microsomes. Depletion of microsomal ATP by the addition of glucose and hexokinase resulted in a rapid Ca^{2+} uptake, then Ca^{2+} release. Further, microsomal Ca^{2+} sequestration was inhibited by alkalating agents e.g. p-chloromercuribenzone, or agents that oxidized e.g. diamide, thiol groups. Microsomal pre-treatment with cystamine, which promotes the formation of protein mixed disulfides, also caused inhibition of Ca^{2+} sequestration. The authors concluded that microsomal Ca^{2+} sequestration is dependent on protein sulfhydryl groups.

(170)

CHAPTER II

MATERIALS AND METHODS

For clarification, and to facilitate following the progression of the experimental procedures used, this section has been divided into three categories:

- I. DEVELOPMENT OF THE MODEL
- II. BIOCHEMICAL CHARACTERIZATION
- III. MITOGENIC RESPONSE EVALUATION.

I. DEVELOPMENT OF THE MODEL

Cell Culture

As part of an ongoing study to assess the protective role of GSH in the oral cavity, human gingival normal fibroblast explants were obtained via biopsy, from six periodontally and systematically healthy volunteers, who had remained medication free for at least 1 week prior to biopsy. Following palatal block anesthesia (to alleviate any tissue disruption from local anesthetic infiltration), the gingiva was harvested from a specified site i.e. interdental papilla between the maxillary 1st molar and 2nd premolar.

Gingival fibroblast cultures were grown in a modification of Eagle Minimum Essential Medium (EMEM) (GIBCO, Grand Island, N.Y.), formula #78-5048. This growth

medium ("B") consisted of EMEM with the addition of 1.5X essential amino acids, 1.5X vitamins, 2X non-essential amino acids, and approximately 1X l-glutamine. Fetal bovine serum (FBS) (GIBCO) was added to the medium for a final concentration of 10% ("B-10"), except during the "synchronization" protocol, in which the serum concentration was decreased to 0.3%. Following the establishment of primary cultures, the fibroblasts were grown in antibiotic free B-10 medium.

At passage, and at cellular harvesting, the cultures were first rinsed with phosphate buffered saline (PBS), then detached from the substrate with equal volumes of 0.02% EDTA-PBS and 0.01% trypsin (prepared from lyophilized trypsin, Worthington Biochemicals, Freehold, N.J., 10 mg/100 ml PBS). All experiments were conducted on cultures that had undergone no greater than 20 population doubling levels (PDL).

Ethionine Mediated Cell Synchronization

Fibroblast cultures were synchronized via a modification of the protocol of Wilke et al. (177,178)

Log phase fibroblast cultures were split 1:2, mono-dispersed, plated and allowed to adhere for 3 hours in B-10 (to alleviate the colonization noted with plating in serum deficient medium). Following attachment, the cells were gently rinsed with room temperature PBS, then synchronized by a 2 cycle blocking procedure. Cells were incubated for

2 interspersed 48 hour periods in B medium with 0.3% FBS (B-0.3), containing a final concentration of 4mM ethionine. (L-Ethionine, #E1260, Sigma Chem. Co., St. Louis, Mo.) Between the ethionine incubations, the cell block was released via a 48 hour incubation in B-10; cultures then underwent a 2nd 2:3 split prior to the 2nd cell block.

Fibroblasts grown with this protocol showed a readily reversible (i.e. <24 hr) G1 phase cellular synchronization; with >80% of the cell population arrested in G1.

GSH Depletion Protocol

Cultures were synchronized via the ethionine protocol. One hour prior to the introduction of B-10, existing levels of GSH were decreased by treatment with 200 uM 2-cyclohexene-1-one (2-CHX-1-one). (Aldrich Chemical Co., Milwaukee, Wis.) The flasks were then rinsed with PBS, and these cultures were incubated for 2 hours in B-10 containing 1×10^{-3} M buthionine sulfoximine (BSO) (Sigma), then fresh B-10 was added. This protocol employs a dual means of decreasing cellular GSH levels. 2-CHX-1-one depletes the cellular levels of existing GSH in a specific fashion, as it is a substrate for GSH-S-transferases. (175) In addition, BSO inhibits glutamylcysteine synthetase, the initial enzymatic step in GSH synthesis. (60) Control cultures were subjected to comparable rinses and medium changes. Cells were incubated at 37°C, 5% CO₂ for 24 or 40 hours for concurrent GSH assessment and flow cytometry.

Culture groups evaluated were:

- L = Log growth controls,
- S = Synchronized controls, harvested immediately after the 6 day low serum/ethionine synchronization,
- L-D 0= Log growth cultures harvested immediately after the GSH depletion protocol,
- S-D 0= Synchronized cultures, harvested immediately after the GSH depletion protocol.
- S 24= Synchronized controls, harvested after an additional 24 hours in B-10,
- S-D 24 = Experimental, synchronized, GSH depleted, harvested after an additional 24 hours in B-10,
- S-D 40 = Experimental, synchronized, GSH depleted harvested after an additional 40 hours in B-10.

II. BIOCHEMICAL CHARACTERIZATION

Determination of Total Cellular GSH

After harvesting and centrifugation, the cell pellet was resuspended to a volume of 0.5 ml in 4mM EDTA-PBS for enumeration and viability (trypan blue exclusion) quantification. Proteins were precipitated via addition of an equal volume of 2M HClO₄ (perchloric acid). The samples were stored at -20° C. Twelve hours or less prior to assay, the supernatants were neutralized to pH 6.02 (using 2M KOH + 0.4M morpholineethanesulfonic acid (MES), (Sigma Chemical Co., St. Louis, Mo.) and frozen overnight.

Cell pellets from samples used in GSSG determination were resuspended to 1 ml in ice cold 1M HClO_4 , 1mM EDTA, 0.02 M NEM (N-ethylmaleimide, Sigma Chemical Co., St. Louis, Mo.). Samples were stored at -20°C . Twelve hours or less prior to the assay, the samples were thawed, 1 ml of 1.3 M K_2HPO_4 was added, samples were stirred for 30 minutes, and then centrifuged to pellet the KClO_4 salt. The NEM contained in the GSSG samples (which were kept on ice during the extraction) was removed by 5 extractions, using a 2 fold excess of ice cold, H_2O saturated, ethyl acetate. A light stream of N_2 gas was used to remove any residual ethyl acetate, and the samples stored at -20°C overnight. For control purposes, the GSSG standards underwent the same ethyl acetate extraction as the GSSG samples.

Cellular levels of total GSH (i.e. GSH, and the disulfide, GSSG) were determined in accordance with the methodology of Eyer and Podhradsky. (4) NADPH, GSH, GSSG, glutathione reductase (#G4759, Type IV) (GR), DTNB, were obtained from Sigma Chemical Co., St. Louis, Mo. The enzymatic kinetics in the Eyer assay have been modified so that the GR step is no longer rate limiting. Therefore, this method is much less susceptible to any endogenous, i.e. intrasample, GR inhibitors. (131)

Rapid reaction kinetics were followed on a SLM-AMINCO DW 2C dual wavelength spectrophotometer, with the following instrument settings: 412 nm excitation, 550 nm emission,

3nm bandwidth, 0.05Abs, 50 sec/in. Sample concentrations (nmol/ml) were obtained by comparison from a 10 point standard curve conducted concurrently. The GSH standard curve ranged from 0.2 nmol/ml to 2 nmol/ml; the GSSG standards ranged from 0.125 nmol/ml to 1.5 nmol/ml. Both the GSH and the GSSG standard curves were linear over their entire range, and results were expressed as GSH or GSSG/mg protein.

Protein Determination

Cellular proteins were determined in accordance with the methodology of Lowry et al. (106). Protein standards were treated in the identical fashion as cell samples i.e. standard proteins were precipitated by addition of equal volumes of 2M HClO₄ and 4mM EDTA. Further, bovine gamma globulins (Sigma Chem. Co., Bovine Cohn Fraction II) were selected as the protein standard due to their intermediate reactivity in the Lowry assay. (138)

Determination of Total and Reduced Nicotinamide Nucleotides

After harvesting the samples that were used to determine the reduced and total (NAD(P)H+NAD(P)⁺) nicotinamide nucleotides were resuspended to 500 ul in ice-cold PBS 0.02% EDTA, to which 2.5 ul of 2M KOH was added, adjusting the pH to ≥ 10.50 . In contrast, the oxidized sample, which served as an internal standard control, was resuspended to 500 ul in a 0.4M MES buffer, pH=6.02, that contained 6×10^{-4} M oxaloacetate. The redox samples were

then transferred to vials designed for cryostorage (Corning Glass Co., Corning, N.Y.).

To induce cellular decompartmentalization, the samples were immersed in liquid N_2 , then rapidly thawed and vortexed vigorously. Preliminary work had demonstrated that this procedure disrupts cellular, cytoplasmic, and nuclear membranes.

Next the reduced and total samples were resuspended to a final volume of 1 ml in the "redox buffer" and the "modified redox buffer", respectively. The redox buffer contained 10 mM Nicotinamide, 20 mM $NaHCO_3$, and 5% Triton X-100 (Calbiochem, La Jolla, Ca), pH=10.62. The modified redox buffer also contained: 20 mM B OH butyrate, 5uM rotenone, 20 mM ETOH, and 0.2 mM aminooxyacetate.

The redox samples were either assayed immediately, or wrapped in foil, and stored overnight in liquid N_2 .

Redox Sample Preparation

Prior to being assayed, the redox samples were heated for 20 minutes in a 60 degree C water bath to degrade the oxidized nicotinamide nucleotides. Preliminary experiments had confirmed the stability (no change in optical density) of the NADPH standards following this heating protocol.

Next, denatured protein was extracted with 2 M $HClO_4$, and the samples centrifuged to pellet the acid precipitable protein. Post centrifugation, the pH of the samples was immediately adjusted to approximately 10.60 with 2M KOH.

The nicotinamide nucleotide redox state assay was conducted on a SLM-AMINCO DW 2 C dual wavelength spectrophotometer using 340 minus 370 nm, 3nm bandwidth. The NAD(P)H levels contained in the samples were determined via comparison against the optical densities obtained from a concurrently conducted 12 point standard curve. The NADPH (#N 7505, Sigma) standards were prepared fresh, in the redox buffer, and ranged from 2.5 nmol/ml-100 nmol/ml. The assay was conducted in the linear portion of the curve ($r \geq 0.999$).

The oxidized sample functioned as an internal standard control, to account for any NAD(P)H spectral changes that were attributable to the presence of cytoplasmic components. Equivalent amounts of the oxidized sample (with the pH adjusted to 10.60) were mixed 1:2 with a specific standard, e.g. #10, thereby diluting the final concentration of the standard by 1/2 (equal to the amount of NADPH in our standard #6). The O.D. obtained from the 1:2 oxidized sample:standard 10 was then compared against the O.D. obtained for standard 6, any increase in O.D. above standard 6 was attributable to cellular-related NADPH spectral changes and this value was then subtracted from all the samples.

III. MITOGEN RESPONSE EVALUATION

Ca²⁺ EGTA Buffer Standard Calibration

Ca^{2+} EGTA calibration buffers were prepared in accordance with the specifications of the computer program designed by Fabiato. (48,49,50) This program incorporates the specified experimental parameters e.g. ionic strength of buffer, temperature, pH, into the calibration calculations. The Ca^{2+} EGTA buffer mixes were prepared in a "calibration buffer" that approximated the intracellular milieu, and contained 20 mM pH 7.2 K-HEPES (Sigma) 120 mM KCl, 1 mM MgSO_4 , and 1 mM EGTA. Atomic absorption was used to confirm the molarity of the CaCl_2 stock. Eight graded Ca^{2+} EGTA buffer mixes, that ranged from 56.2 nM to 1000 nM free Ca^{2+} , were prepared. The free Ca^{2+} concentrations of the Ca^{2+} EGTA standards were determined via a Ca^{2+} electrode, using reference buffers. (61)

Two fura dissociation constants (K_d) were determined. First, the K_d for fura-2 (pentapotassium salt, Molecular Probes) was determined in the Ca^{2+} -EGTA buffer mixes. Because cellular components can affect fura binding, a second K_d for fura-2 AM was determined in the cellular in situ calibration system. (27) For the determination of this K_d , the Ca^{2+} -EGTA buffer mixes contained cells ($3 \times 10^6/\text{ml}$) with collapsed gradients. A correlation coefficient was calculated by determination of the line generated from the fluorimetry obtained free Ca^{2+} ratios (345/380 vs 510nm.) vs the known Ca^{2+} concentrations of the Ca^{2+} -EGTA buffer mixes. ($r=0.999$ for both the cell free

and cell containing systems). The correlation coefficient was then divided by the slope of the line generated to find the K_d . The K_d for the cell free system was determined to be 167; for the in situ system, the K_d was 128.

In situ Calibration and Collapse of Intracellular Gradients

Fibroblasts were harvested, centrifuged, and the cell pellets were resuspended in B medium containing 0.1% bovine serum albumin (BSA), (Sigma), at approximately 3×10^6 cells/ml. The fura-2 AM (Molecular Probes, Inc.) was dissolved in anhydrous dimethyl sulfoxide (#27, 685-5, Aldrich Chemical Co., Milwaukee, Wi.) on the days of the experiments. The cells were then loaded with dye by an incubation with 1 μ M fura-2 AM for 45 minutes at 37°C, 5% CO₂. During, and following the fura loading, the samples were protected from light.

Post incubation, the fura-loaded cells were washed twice in Ca²⁺ and chelator free PBS, and then resuspended in the graded Ca²⁺ EGTA buffer mixes. Selective ionophores and other metabolic inhibitors were added to each of the fura-loaded cell samples that were suspended in the Ca²⁺ EGTA calibration standards to abolish the intracellular gradients. To the "in situ" standards, which were prepared in the "calibration buffer" (selected as an electrolyte solution approximating the intracellular milieu) the following were added (final concentration) 20 mM NaN₃ (Sigma), 10 mM 2-deoxyglucose (Grade III, Sigma),

0.5 ug/ml nigericin (#481990, Calbiochem, La Jolla, Ca.), 2 uM ionomycin (Calbiochem), and 2 uM carbonyl cyanide M-chloropropenyl hydrazone (CCCP) (Calbiochem). The reagents employed to collapse the intracellular gradients had been shown to not affect the fluorescence of a related fluorochrome, indo-1. (27)

The in situ standards were then equilibrated by a 180 minute incubation in an Orbit Microprocessor Shaker Bath (Lab Line Instruments, Melrose Park, Il.) set at 37° C, with the speed at 100 rpm. Post equilibration, the pH of the in situ standards was determined, and if necessary, re-adjusted back to a pH of 7.2.

Fluorimetry

Readings of the in situ cellular standards were conducted on a Perkin Elmer LS-5B luminescence spectrophotometer (Perkin Elmer, Oak Brook, Il.) with the excitation and emission band width at 10nm. Excitation was conducted at 345 nm and 380 nm, emission at 510 nm. The standards and cuvette holders were maintained at 37° C by a Haake D3 circulating water bath (Haake, Berlin, W. Germany). Further the samples were read during constant stirring, by the use of a tapered top cuvette stirring system (Instech Labs, Plymouth Meeting, Pa.).

The in situ calibration curve was conducted several times to ensure reproducibility. The free $[Ca^{2+}]_i$ of resting fibroblasts was determined to be 98, 93, and 101 nM

on three separate evaluations; values which closely approximate reported resting $[Ca]_i$ levels of 100 nM. (78) Ca^{2+} mobilization experiments were then run at identical instrument settings. Cellular samples were evaluated to determine the daily R_{min} and R_{max} in conjunction with every Ca^{2+} mobilization assay. Free Ca^{2+} was determined using the ratio method of Grynkiewicz et al. (61)

Ca^{2+} Mobilization Assay

Cell samples were obtained from all the culture growth regimines i.e. L, S, S 24, S-D 24 to assess cellular responsiveness to the mitogen, B FGF, by monitoring $[Ca^{2+}]_i$.

Cell loading with fura was conducted as in the in situ protocol. Two separate parameters of the Ca^{2+} mobilization response were evaluated. To determine cellular capacity to respond with intrinsic Ca^{2+} stores, fura loaded cells were resuspended in Ca^{2+} free PBS. Ratio readings were first taken to determine sample prechallenged levels of $[Ca^{2+}]_i$. Then, the samples were challenged with B FGF (#1123149, human recombinant B FGF, Boehringer Mannheim, Indianapolis, In.) at a dose of 50 ng/ml. Ratio readings were recorded immediately post addition of B FGF, and every 30 seconds thereafter for 7 minutes.

Because part of the Ca^{2+} mitogenic response is felt to involve cellular ability to mobilize Ca^{2+} from both intracellular and extracellular sites, fura loaded cells

were also suspended in a " Ca^{2+} mobilization buffer". (78) This buffer contained 10 mM K-HEPES (pH7.2), 145 mM NaCl_2 , 5 mM KCl, 1mM Mg SO_4 , 0.5 mM Na_2HPO_4 , 5 mM glucose, and physiologic extracellular concentrations (1mM) of CaCl_2 . Cell samples were incubated at 37°C for 45 minutes in this buffer, unstimulated Ca^{2+} levels recorded, and then these cells were challenged with B FGF and the free Ca^{2+} ratios recorded.

Human B FGF was selected as the mitogenic stimulus for several reasons. B FGF has been shown to induce Ca^{2+} mobilization from numerous cultured cells, including fibroblasts (56,57). Further, as these cultures are human cells, the species compatibility should be reflected in appropriate receptor expression.

Cell Preparation for Flow Cytometry and DNA Staining Procedures

Cell nuclear preparations and DNA staining were conducted in accordance with the methods of Larsen et al. (97) Cells were harvested, suspended in PBS, and centrifuged at $500 \times g$, 4°C , for 10 minutes. The pellet was resuspended to 2 ml in a phosphate (6.5 mM Na_2HPO_4 , 2.7 mM $\text{K H}_2\text{PO}_4$, 137 mM NaCl , 0.5 nM EDTA, pH7.2) - 0.1% Triton buffer and placed in an ice bath for five minutes. The cells were then lysed via tube inversion. Fixation of the nuclear preparation was done by addition of 0.7 ml of a 4%

formaldehyde solution. Samples were stored at 4°C for < 12 hour prior to flow cytometry.

To remove RNA, the nuclei were incubated on the day of staining for 20 minutes in a 37°C H₂O bath with a final concentration of 150 U/ml of RNAase (RNAase A, Worthington Biochemicals, Freehold, N.J.). Prior to centrifugation, the nuclei were vortexed (to remove cytoplasmic tags) and passed through a 53u nylon mesh filter, (Spectra/Mesh, Spectrum Medical Industries, Inc., Los Angeles, Ca), to remove cellular debris.

Finally, the nuclei were suspended in 50 ug/ml propidium iodide (PI) (Sigma), and incubated overnight at 4°C. Flow cytometry was conducted the following morning.

Flow Cytometry

The DNA content of the PI stained nuclei was analyzed on an Ortho System 50H Cytofluorograf (Bectin Dickinson & Co.) at a flow rate of less than 300 particles per second. The nuclei were suspended in a phosphate-0.1% Triton buffer at a concentration of approximately 10⁶/ml. The carrier sheath fluid was phosphate buffered normal saline. Excitation output was 200 milliwatts from the 488 nanometer line of a Spectra Physics 164-05 argon ion laser in light mode. The PI emission was collected through a 630 long-pass filter, and analyzed on an Ortho 2150 Data Acquisition System (Data General Nova 4 CPU, running MPOS). Prior to each run of samples, instrument linearity and calibration

were verified with fluorescent microspheres (PolySciences, Inc., Warrington, Pa.). Repeatability over time was confirmed with P.I. microbead standards (Flow Cytometry Standards Corp., Research Park, N. Carolina) as an experimental internal standard. PI stained fixed calf thymocyte nuclei confirmed a 2:1 (G2/M:G1G0) DNA index. The distribution of nuclei in specific cell cycle phases (G0/G1, S, G2/M) was derived from the area of the red fluorescence histograms, assuming a gaussian distribution of the G-G1 and G2/M about their maxima, and attributing the inclusive residual DNA histogram segment to S phase nuclei. Debris, aggregates, and doublet nuclei (which falsely increase the G2/M population) were excluded from the DNA histogram by first plotting fluorescence pulse peak versus pulse area as a bitmap, two parameter, cytogram and using a logic gate to exclude nonconforming pulses.

³H Thymidine DNA Synthesis

The DNA synthesis assay was conducted in triplicate, using 24 well plates, with a 1 hour pulse label of 1 u Ci of ³H-thymidine (61 C/mM) per well. Three culture groups, S+B0.3, S+B-10 and S-D+B-10 were evaluated. Cells were cultured in T150 flasks until the final 48 hours of the synchronization protocol. At this time, the cells were transferred, at a density of approximately 40,000 cells/well, to the 24 well plates. This precaution was taken to avoid any cellular manipulation prior to the

addition of the ^3H -thymidine label. Sample collection time points and numbers of samples taken varied depending upon the experimental group.

Because of the reports by Wilke et al., in which irreversible growth arrest occurred in cultures maintained for >72 hours in ethionine containing medium, the synchronized +B0.3/ethionine (S+B0.3) cultures had samples taken at -23, -18, -12, -6, 0, 12, and 24 hours. (177, 178) Therefore, the time pulls at 0, 12 and 24 hours showed the ^3H thymidine incorporation of a culture that had completed the 6 day synchronization protocol.

The synchronized +B-10 (S+B-10) and synchronized, GSH depleted (S-D+B-10) cultures had samples taken at 13, 19, 22, 25, 28, 31, 34, 37, 40, 43, 49, 55 and 61 hours, the S+B-10 group had an additional sample taken at 0 hours.

Cellular harvesting was conducted at 4°C , using ice-cold reagents. The cells were washed twice with PBS, and then washed once with PBS-0.02% EDTA. Cells were harvested using 0.2 ml trypsin per cell well. Post trypsinization, 0.8 ml of B-10 was added to each well, and the cell suspensions were transferred to labelled sample tubes containing 1 ml of 10% TCA and vortexed vigorously. Cell samples were stored at 4°C until transfer of the sample to filter papers (Whatman GF/A glass microfibre paper, Whatman Ltd. Maidstone, England) which were then added directly to the scintillation cocktail.

Sister wells (unlabelled cells from the same culture groups) were harvested concurrently with each labelled sample to provide cell counts.

CHAPTER III

RESULTS

Preliminary studies were conducted to determine levels of GSH in the fibroblast cultures. A representative standard curve for the GSH kinetic assay is shown in Fig 1. Fibroblast GSH levels (nmol/mg protein) ranged from 13.29 to 51.62, $\bar{x} = 32.10$, $n = 25$. (Table 1) Since levels of GSSG were below detection in these cultures ($<10^{-8}M$), the GSH was present in its reduced state.

Subsequent studies were then conducted to determine if the range in culture GSH levels was associated with population cell cycle distribution. Concurrent determination of culture GSH levels and DNA content, by flow cytometry, showed a GSH-cell cycle association. (Table 2, Fig. 2) Cultures with the lowest GSH levels (nmol/mg protein), $\bar{X} = 20.08 \pm 0.37$, showed the highest population distribution in G1, $73.55 \pm 2.23\%$. Intermediate GSH levels, 35.00 ± 2.61 , were detected in cultures that contained an even dispersion of their cellular populations in G1 and beyond G1, $51.43 \pm 4.04\%$ of cells in G1. Highest GSH levels, 44.43 ± 1.84 , were found in cultures with the majority of their cellular populations beyond G1 ($29.13 \pm 15.08\%$ in G1). Further, there was a strong positive

correlation ($r=0.999$) between the cultures' DNA index and GSH levels. (Fig. 2) Further work was then conducted to clarify the GSH-cell cycle association. These studies specifically addressed whether GSH is necessary for cell cycle progression beyond G1. To be able to answer this question required development of both a G1, stage specific population baseline and an effective and reversible GSH depletion protocol.

Initial attempts at culture synchronization employed a 72 hour incubation in low serum (B 0.3) medium. Results obtained from flow cytometric DNA analyses revealed that serum deprivation did not provide G1 synchronization. Over 50% of the cellular populations were not in G1. The serum-deprived cells appeared to not have arrested in G1; instead they were "leisurely" progressing in the cell cycle and were distributed in S, G2 and M.

Ethionine was then selected, to be used in conjunction with low serum medium, to provide G1 growth arrest. Ethionine was selected as the synchronization agent for several reasons. Wilke et al. had reported using ethionine to obtain a reversible, G1 growth arrest in normal human keratinocytes. Further, ethionine usage was appealing because the contribution of the cellular redox poise was a parameter of interest. The putative mechanism of action of ethionine entails a decrease in availability of cellular adenine nucleotides. (108)

Figure 1

Representative standard curve obtained during GSH determination assay. GSH standards ranged from 0.2 nmol/ml to 2 nmol/ml, with 0.2 nmol/ml increments. The correlation coefficient was routinely $r \geq 0.999$. This assay was conducted on a SLM-AMINCO DW 2C dual wavelength spectrophotometer, with the following instrument settings: 412 nm excitation, 550 nm emission, 3 nm bandwidth, 0.05Abs, 50 sec/in.

Figure 1

Table 1

Preliminary GSH evaluation results

<u>Group</u>	<u>(nmol/mg protein)</u>	<u>x Protein (mg)</u>	<u>Cell Counts</u>
	<u>x GSH</u>	<u>(mg)</u>	
I n=8	17.58 $\pm$ 3.07 (SD)	0.236 $\pm$ 0.075 (SD)	1.67x10 ⁶ $\pm$ 0.47 (SD)
II n=3	28.80 $\pm$ 0.66	0.333 $\pm$ 0.034	1.72x10 ⁶ $\pm$ 0.31
III n=8	35.46 $\pm$ 2.38	0.429 $\pm$ 0.058	1.93x10 ⁶ $\pm$ 0.47
IV n=6	44.59 $\pm$ 4.12	0.434 $\pm$ 0.080	1.62x10 ⁶ $\pm$ 0.96

Table 2

Correlation Of GSH Content With Cell Cycle Distribution*

Cell Cycle		
<u>Group</u>	<u>x[GSH](nmol/mg protein)</u>	<u>Distribution (% cells)</u>
I n=3	20.08±0.37	G1=73.55±2.23 S=18.77±3.56 G2/M= 7.67±5.81
II n=8	35.00±2.61	G1=51.43± 4.04 S=32.30±10.65 G2/M=16.26±10.48
III n=4	44.43±1.84	G1=29.13±15.08 S=57.90±27.24 G2/M=12.97±12.19

*=All data expressed as mean ± standard deviation, n=number separate cultures in each assay.

Figure 2

Linear regression line ($r=0.999$) that is generated from data in Table 2 by comparison of the mean levels of GSH (nmol/mg protein) vs the group DNA Index. The DNA Index was determined by summation of the groups' cell cycle distributions $[(x\%G1 \times 2) + (x\%S \times 3) + (x\%G2/M \times 4)]$, and then converting the percentages to integers.

Figure 2

Figure 3

Representative histograms of DNA related fluorescence of propidium iodide stained human gingival fibroblast nuclei that were harvested following: L:Log phase growth; S:6 day ethionine synchronization protocol; S 24:24 hr incubation in rich medium after synchronization; S-D 24:24 hr incubation in rich medium after synchronization and glutathione (GSH) depletion; S-D 40:40 hr incubation in rich medium after synchronization and GSH depletion.

Figure 3

Figure 4
Photomicrographs taken at a 50 X magnification of
fibroblast cultures harvested:

- L = during log growth
- S = following 6 day ethionine synchronization
protocol,
- S 24 = following ethionine synchronization, then a 24
hour incubation in B-10,
- S-D 24 = following ethionine synchronization and GSH
depletion, followed by a 24 hour incubation in B-
10.

Figure 4

The modified Wilke ethionine synchronization protocol provided a G1 stage specific population baseline. Culture synchronization efficacy was evaluated by 2 methods, flow cytometric DNA analyses and ^3H thymidine incorporation. DNA analyses showed that following the ethionine protocol, $\geq 80\%$ of the synchronized fibroblasts were in G1. (Fig. 3) ^3H thymidine incorporation was negligible in the synchronized cultures. (Data will be presented in an upcoming section of results.)

Studies were also conducted to determine the efficiency of the GSH depletion protocol and to evaluate if this procedure affected cellular integrity. Viability, as determined by trypan blue exclusion, was routinely $\geq 93\%$ for all cultures at harvest. (data not shown) Further, the GSH depletion protocol was shown to be both effective and reversible. There was a significant difference in S-D 24 GSH levels ($p < 0.001$ t test and Mann Whitney U) in comparison to GSH levels contained in groups L, S, S 24, and S-D 40. (Table 3) GSH depletion treated cultures (S-D) were restored to log growth GSH levels after 40 hours in B-10. (Table 3) These findings confirmed that this protocol provided significant GSH depletion, while concurrently maintaining cellular viability inclusive of the capacity of GSH recovery.

Qualitative cellular changes were apparent in the synchronized (S) and synchronized, GSH depleted (S-D)

Table 3

Correlation of GSH Content With
Cell Cycle Progression Following Culture
Synchronization and GSH Depletion Protocols *

<u>LABEL</u>	<u>PROTEIN (mg)</u>	<u>[GSH]</u>	<u>CELL CYCLE</u>
		<u>(nmol/mg protein)</u>	<u>(x% POPULATION IN G1)</u>
L	0.445±0.147 n=8	39.75±8.53 n=8	49.48±15.67 n=6
S	0.229±0.078 n=9	48.05±12.63 n=9	79.83±3.34 n=5
S 24	0.426±0.079 n=10	44.62±8.99 n=10	37.28±17.63 n=8
S-D 24	0.386±0.070 n=10	3.93±2.81 n=10	74.23±6.27 n=7
S-D 40	0.281±0.011 n=2	35.04±0.20 n=2	43.96±1.33 n=2

- L = Log growth controls
- S = Synchronized controls, harvested immediately after the 6 day ethionine synchronization
- S 24 = Synchronized controls, harvested after an additional 24 hours in B-10
- S-D 24 = Experimental, synchronized, GSH depleted, harvested after an additional 24 hours in B-10
- S-D 40 = Experimental, synchronized, GSH depleted cultures that were harvested after an additional 40 hrs. in B-10.

*=All data expressed as mean ± standard deviation, n=number of separate cultures in each assay.

cultures. (Fig. 4) In both the S and the S-D cultures, there was a marked absence of mitotic figures. This finding demonstrated the lack of ongoing cell cycle progression. Synchronized cells appeared elongated and spindly in comparison to log growth fibroblasts. Some of the S-D fibroblasts showed marked cytoskeletal changes. These cytoskeletal alterations were most apparent immediately after GSH depletion. However, a few S-D cells still retained a rounded morphology 24 hours after GSH depletion.

Because of the importance of the GSH/GSSG ratio as an indicator of the cellular redox state, and despite nondetectable levels of GSSG in our preliminary studies, cellular GSSG levels continued to be monitored throughout the synchronization and GSH depletion protocols. Results remained consistent with our earlier findings. The oxidized, GSSG form, remained nondetectable in culture groups L, S, S 24, S-D 24, S-D 40. Further, despite the low levels of GSH in the S-D 24 cultures, the existing GSH was present in its reduced state.

To evaluate the immediate cellular response to GSH depletion, GSH & GSSG samples were also taken immediately following the GSH depletion protocol from both L & S cultures. These studies asked two specific questions. First, would the response to GSH depletion vary between log growth and synchronized cultures? Second, would a

transient intracellular oxidation occur following cellular exposure to 2-CHX-1-one and BSO? The results demonstrated that log growth and synchronized cells respond similarly to GSH depletion. GSH levels were nondetectable in both groups. Further, as GSSG levels were also nondetectable, it was determined that intracellular retention of GSSG did not occur, even as a temporary event, following GSH depletion.

These findings addressed the oxidant stress issue. Lack of intracellular retention of GSSG ruled out the formation of protein mixed disulfides as a consequence of the GSH depletion protocol and clarified the targeted nature of our depletion regimine.

Experiments were then conducted to determine if the ethionine synchronization protocol affected the cellular redox state.

As would be anticipated based upon ethionine's proposed mechanism of action, nicotinamide nucleotide assay results demonstrated that the ethionine synchronization protocol perturbed the cellular redox state. (Table 4) The altered redox poise (relative to log growth control cultures) observed in our synchronized cultures was both transient and reversible. Ethionine synchronized cultures, incubated for an additional 24 hours in B-10 (Group S 24) showed a rebound of culture redox poise to a more reduced state ($43.90 \pm 13.43\%$ oxidized to $15.69 \pm 3.65\%$ oxidized), and a

decrease in total nicotinamide nucleotide levels (12.77 ± 0.70 nmol/mg protein to 10.46 ± 1.21 nmol/mg protein), to distributions approximating those seen in log growth cultures (Group L) (10.00 ± 1.35 nmol/mg protein). In contrast, the redox state and total nicotinamide nucleotide levels of the synchronized, GSH depleted cultures (Group S-D 24) did not deviate from the levels obtained from the synchronized control cultures (Group S), despite an additional 24 hour incubation of the D cultures in B-10.

Total nicotinamide nucleotide levels were elevated in both the synchronized control (Group S) (12.77 ± 0.70 nmol/mg protein) and the synchronized, GSH depleted (Group S-D 24) (12.07 ± 1.03 nmol/mg protein), cultures relative to levels obtained from log growth cultures (Group L). (10.00 ± 1.35 nmol/mg protein). Comparison of total nicotinamide nucleotide levels, Group L vs Group S is significant via the test, $p < 0.05$.

Relative to log growth (L) cultures, ($9.93 \pm 5.05\%$ oxidized), oxidation of the cellular redox state was noted in both the S and S-D cultures. Further, these differences were statistically significant; the percent redox poise oxidation L vs S was significant at the $p < 0.01$ level, L vs S-D 24, $p < 0.05$, students t test.

The two time course GSH depletion experiments provided information regarding culture redox poise response to the

Table 4

Distribution of Nicotinamide Nucleotides(nmol nicotinamide nucleotides/mg protein $\pm$ S.D.)

NAD(P)H+NAD(P) ⁺		NAD(P)H	NAD(P) ⁺	
<u>Group</u>	<u>Total</u>	<u>Reduced</u>	<u>Oxidized</u>	<u>%Oxidized</u>
L				
n=4	10.00 $\pm$ 1.35*	9.01 $\pm$ 1.56	0.98 $\pm$ 0.41	9.93 $\pm$ 5.05*+
S				
n=3	12.77 $\pm$ 0.70*	7.21 $\pm$ 2.05	5.56 $\pm$ 1.56	43.90 $\pm$ 13.43+
S 24				
n=3	10.46 $\pm$ 1.21	8.79 $\pm$ 0.68	1.49 $\pm$ 0.29	15.69 $\pm$ 3.65
S-D 24				
n=3	12.07 $\pm$ 1.03	6.93 $\pm$ 1.27	5.13 $\pm$ 1.32	42.47 $\pm$ 9.46*
S-D 0				
n=1	13.62	9.56	4.06	29.79
S-D 40'				
n=1	9.00	6.38	2.62	29.11

L = Log growth controls,

S = Synchronized controls, harvested immediately after the 6-day low serum/ethionine synchronization,

S 24 = Synchronized controls, harvested after an additional 24 hours in B-10,

S-D 24 = Synchronized, GSH depleted, harvested after 24 hrs in B-10

S-D 0** = Experimental, synchronized, harvested immediately after the GSH depletion,

S-D 40 = Experimental, synchronized, harvested 40 hrs after GSH depletion.

* = Significant $p < 0.05$ + = Significant $p < 0.01$

GSH depletion protocol. S-D 0 (cultures harvested immediately post GSH depletion) showed a percent oxidation lower than that seen in S-D 24 cultures which were harvested at 24 hours; (29.79% oxidation vs 42.47%).

Therefore, the cellular redox poise oxidation increased during the 24 hours following the GSH depletion. S-D 40 cultures (harvested 40 hours post GSH depletion) showed both a decrease in percent oxidation (29.11%) and total nicotinamide nucleotide levels (9.00 nmol/mg) relative to the S-D 24 cultures, which were harvested at 24 hours.

The preliminary results confirmed that a satisfactory model had been developed, through which one could study the two specific research questions. First, is GSH a necessary intracellular component for cell cycle progression beyond G1? Second, is the cellular thiol redox status an intracellular factor that affects how cells in G1 and mitogens interact?

Two aspects of the mitogen-cellular response were evaluated in subsequent studies. The sustained mitogenic responsiveness, to evaluate cell cycle progression, was determined by ^3H thymidine incorporation and flow cytometric DNA analyses. Immediate mitogenic responsiveness was determined by measuring increases in $[\text{Ca}^{2+}]_i$ in response to B FGF.

Additional background experiments were required to develop the Ca^{2+} mobilization assay. To confirm the free

Ca^{2+} levels in the Ca^{2+} EGTA buffer mixes, a fura-2 pentapotassium salt standard curve was conducted. The standard curve obtained is shown in Fig. 5, the 345/380 nm vs 510 nm ratios are shown in Fig. 6.

As discussed (in detail) in the Methods section, two fura K_d were determined. The K_d for the cell free, fura-2 pentapotassium salt was determined to be 167; for the cellular in situ system, the K_d was 128.

Experiments were conducted to determine fibroblast resting $[\text{Ca}^{2+}]_i$. Fura loaded cells were suspended in the Ca^{2+} EGTA buffer mix that had a free $[\text{Ca}^{2+}]_i$ of 10^{-7}M , and the intracellular free Ca^{2+} ratios recorded. The free $[\text{Ca}^{2+}]_i$ of resting fibroblasts was determined to be 98, 93 and 101 nM on three separate evaluations.

Studies were then performed to determine whether the fibroblasts possessed the esterase necessary to hydrolyze the fura -2 AM.

Resting, fura loaded cells showed an excitation peak at 350 nm. Following cell lysis with 0.1% Triton X-100, the excitation spectra in the presence and absence of 5mM EGTA were consistent with the fura -2 anion, indicative of de-esterification of the AM, (Fig. 7) As a final preliminary control study, 15 ul aliquots of a 1 mg BSA/1 ml PBS solution (the B FGF solvent) were added to fura loaded cells during constant stirring, and the free $[\text{Ca}^{2+}]_i$ ratios

were recorded. No change in $[Ca^{2+}]_i$ levels occurred after the addition of the BSA in PBS solution.

Results obtained by both 3H -thymidine incorporation and flow cytometry showed an association between cellular GSH levels and cellular capacity to progress beyond G1. (Table 3, Fig. 8) Importantly, the ethionine synchronization protocol did not affect cellular GSH levels. Synchronized cultures contained comparable levels of GSH as log growth fibroblasts. By 24 hours after the reintroduction of rich medium, these formerly synchronized cultures responded with cell cycle progression beyond G1. In contrast, the synchronized, GSH depleted cultures remained in a G1 state at the 24 hour harvest. Comparison of the DNA distributions, as determined by flow cytometry, showed that a statistically significant difference ($p < 0.001$, t test, Mann Whitney U) existed between the synchronized S 24), and synchronized, GSH depleted (S-D 24) cultures with regard to cell cycle progression beyond G1 at the 24 hour harvest.

Results obtained from the 3H -thymidine incorporation assay were consistent with our flow cytometry data. (Fig 8) Tritiated thymidine incorporation was negligible in the

Figure 5

Fura-2 pentapotassium salt Ca^{2+} EGTA buffer mix standard curve. Levels of free Ca^{2+} in the Ca^{2+} EGTA buffer mixes ranged from 56.2 nM to 1000 nM. Also evaluated were a minimum (0 nM Ca^{2+}) and a maximum (821 mM free Ca^{2+}). Readings were conducted on a Perkin Elmer LS-5B luminescence spectrophotometer, with the excitation and emission band width at 10 nm, excitation conducted at 345 and 380 nm, emission at 510 nm.

Figure 5

Figure 6

Free Ca^{2+} ratio recordings for the fura-2 pentapotassium salt Ca^{2+} EGTA buffer mixes standard curve. Levels of free Ca^{2+} in the EGTA buffer mixes ranged from 56.2 nM to 1000 nM; 0 mM and 821 mM free Ca^{2+} samples were conducted for the minimum and maximum, respectively.

Figure 6

Figure 7

Demonstration of functional esterase activity in fura loaded fibroblasts. The curve with a peak at 350 nm is obtained from intact, fura-loaded fibroblasts. Curves that peak at 345 nm and 360 nm are obtained after cell lysis with Triton x-100 in the absence, and presence of 5 mM EGTA, respectively.

Figure 7

control, synchronized cultures that were maintained in the low serum/ethionine media. Recovery from the G1 growth arrested state was prolonged in the S-D cultures.

Synchronized cultures showed initiation of DNA synthesis between 13 - 19 hours following reintroduction of B-10; S-D cultures entered S between 25 - 28 hours. Further, there was a 6-hour difference in ^3H -thymidine incorporation peaks; S cultures peaked at 28 hours, S-D cultures at 34 hours. A 2nd peak of ^3H -thymidine incorporation occurred in the S cultures between 43 - 49 hours. The timing (24 hours post earlier initiation of DNA synthesis) and appearance of this peak is most consistent with a second round of DNA synthesis occurring in some of the cells as population synchronization is lost.

After 40 hours in rich medium, the formerly GSH depleted cultures had concurrently restored their GSH levels and showed cell cycle progression.

Results obtained from the flow cytometric DNA analyses showed an association between cellular sustained mitogenic responsiveness and a specific cellular thiol redox state. When cultures progressed beyond G1 they contained abundant reducing equivalents in the form of reduced nicotinamide nucleotides and increased GSH levels. Synchronized cultures had recovered from the ethionine induced redox poise oxidation after 24 hours in B-10, and progressed beyond G1. In contrast, the S-D 24 cultures, which

Figure 8

Tritiated thymidine incorporation (1 hour pulse label, of 1 μ Ci 3 H-thymidine (61 Ci/mM) vs. time in culture. The synchronization protocol was completed at time 0. Human gingival fibroblasts were pulse labelled during the following cell conditions:

Completion of the final 24 hours of the synchronization protocol, and then culture maintenance for the next 24 hours in B0.3+ethionine.

Introduction of ethionine-free, B-10 at time 0 to G1 synchronized cultures.

Introduction of ethionine-free, B-10 at time 0 to G1 synchronized, GSH depleted, cultures.

Figure 8

retained an oxidized redox poise and depleted GSH levels after 24 hours in B-10, did not show cell cycle progression.

Groups S and S-D 24 contained comparable population distributions in the G1, growth arrested state despite the fact that the S-D 24 cultures had an additional 24 hours in B-10 after synchronization. (Fig. 9) In comparison to either the S or S-D 24 cultures, the S 24 cultures showed a significant difference ($p < 0.001$ Mann Whitney U) in cell cycle progression. Formerly G1, growth arrested cultures resumed cell cycle progression following cellular GSH and redox poise recovery. (Fig. 10)

Results of the B FGF assays (immediate mitogen responsiveness) revealed that there existed statistically significant differences between our culture groups in cellular capacities to respond to a mitogenic stimulus by increasing $[Ca^{2+}]_i$. (Table 5) (Appendix, Tables 6 and 7) Prior to the addition of B FGF, there were no significant differences noted in resting $[Ca^{2+}]_i$, demonstrating that $[Ca^{2+}]_i$ were not affected by thiol redox perturbation. All culture groups showed a response lag period (between 0-30 sec) following addition of B FGF. Maximal increases in $[Ca^{2+}]_i$ were recorded 1 - 3 minutes after FGF challenge, after which $[Ca^{2+}]_i$ slowly declined and then plateaued at levels 1.5 to 2 fold above resting $[Ca^{2+}]_i$. Groups L and S 24 showed the highest B FGF responses; with maximal levels

of $[Ca^{2+}]_i$ recorded when these cells were suspended in the "Ca²⁺ Mobilization Buffer." (Maximal stimulated $[Ca^{2+}]_i$ 406.69±49.50nM±S.D. and 401.00±44.38nM,±S.D., n=4 for each group, groups L and S 24, respectively.) The Kruskal-Wallis One Way Analysis of Variance test showed a $p < 0.05$ for both the intrinsic and challenge parameters in comparison of Groups L, S, S 24 and S-D 24. Next, a Mann Whitney U test was conducted to determine between which specific groups (L, Fig. 10S, S 24, or S-D 24) the differences existed. The same levels of significance were found (Intrinsic $p < 0.008$, Challenge $p < 0.014$) in comparison of the log growth (L) cultures with either the synchronized (S) or the synchronized, GSH depleted cultures (S-D 24) in cellular capacities to mobilize Ca²⁺. Comparison of the synchronized, 24 hour B-10 (S 24) culture Ca²⁺ results vs either the S or S-D 24 group results showed that the S 24 cultures responded with significantly higher $[Ca^{2+}]_i$ $p < 0.029$ for both the intrinsic and challenge parameters. No difference in Ca²⁺ mobilization capacity was found in comparison of the L vs S 24 culture groups.

Comparison of our redox poise, Ca²⁺ mobilization, and cell cycle progression results demonstrated the association of cellular thiol redox state with mitogen induced responses. After 24 hours in B-10, the redox poise of the S 24 cultures became more reduced, and then both intracellular GSH levels and redox state approximated

Figure 9

Bar histogram depicting cell cycle distribution (as $\bar{X}$ fraction $\pm$ SD of cell population in G1) of human gingival fibroblasts that were harvested for flow cytometric analysis after: Group S: Ethionine synchronization protocol; Group S-D 24: 24 hr incubation in rich medium after synchronization and GSH depletion protocols; Group S 24: 24 hr incubation in rich medium after synchronization protocol.

Figure 10

Normalized data (based upon establishment of peak G1 distribution of group D cultures to 100), obtained during concurrently conducted experiments, of DNA related fluorescence of propidium iodide stained human gingival fibroblast nuclei that were harvested following: (_____) S-D 24: 24 hour incubation in rich medium post synchronization and GSH depletion; (-----) S-D 40: 40 hour incubation in rich medium post synchronization and GSH depletion.

Table 5

Increase in $[Ca^{2+}]_i$ in Response to B FGF*

Group	<u>Intrinsic</u>		<u>Challenge</u>	
	$[Ca^{2+}]_i$ (nM)		$[Ca^{2+}]_i$ (nM)	
	<u>pre B FGF</u>		<u>pre B FGF</u>	
L:	213 $\pm$ 39	17.8 $\pm$ 7.5 n=5	360 $\pm$ 44	13.0 $\pm$ 1.4 n=4
S:	209 $\pm$ 26	8.6 $\pm$ 1.4 n=4	346 $\pm$ 40	5.1 $\pm$ 3.3 n=4
S 24:	208 $\pm$ 37	17.9 $\pm$ 8.2 n=4	344 $\pm$ 38	17.0 $\pm$ 7.5 n=4
S-D 24:	196 $\pm$ 30	7.4 $\pm$ 3.1 n=4	350 $\pm$ 30	4.7 $\pm$ 3.0 n=4

* All data expressed as mean % increase above nonchallenged $[Ca^{2+}]_i \pm$ S.D. n = number of separate cultures in each assay

** Intrinsic: Fura-loaded cells resuspended in PBS

** Challenge: Fura-loaded cells resuspended in Ca^{2+} Mobilization Buffer

those in L cultures. The increases in $[Ca^{2+}]_i$ in response to B FGF was also similar in these two groups. In addition, the S 24 cultures showed a sustained mitogenic responsiveness by demonstrating significant cell cycle progression beyond their G1 growth arrested state after 24 hours in rich medium. In contrast, the redox state of both the S and S-D 24 cultures remained significantly oxidized at harvest. These culture groups were also less capable of responding to mitogenic stimuli by Ca^{2+} mobilization or cell cycle progression. There exists a strong positive correlation in comparison of culture group x levels of reduced nicotinamide nucleotides with culture capacity for mitogen induced Ca^{2+} mobilization. (Fig. 11) ($r=0.995$, Intrinsic vs x levels reduced nicotinamide nucleotides, $r=0.938$ Challenge vs x levels reduced nicotinamide nucleotides.)

Figure 11

Linear regression relationship between x levels of reduced nicotinamide nucleotides and B FGF stimulated percent increase in $[Ca^{2+}]_i$ in human gingival fibroblasts from the following experimental groups:

L = Log growth; S = Synchronized, harvested after the 6 day ethionine synchronization; S 24 = Synchronized, harvested after an additional 24 hr incubation in B-10; S-D 24 = Synchronized, GSH depleted, harvested after an additional 24 hr incubation in B-10. Intrinsic = fura loaded cells suspended in PBS. Challenge = fura loaded cells suspended in " Ca^{2+} Mobilization Buffer" (see Methods)

Figure 11

CHAPTER IV

DISCUSSION

The regulation of cell cycle progression is complex and entails the interaction between extracellular factors and intracellular responses. (135,124,4,31) Evidence suggests that intracellular factors are key in determining how cells respond to mitogens. (135,124,4,31) In vivo, cells are bathed in mitogen-rich serum; yet are only responsive to these mitogens during the G1 phase of the cell cycle. (135,124,4,31) Previous studies have reported G1 related changes in cellular GSH and thiol levels. (146,67,157)

A common finding among previous reports evaluating cellular GSH levels and this study is the association of increased cellular GSH content with ongoing cell cycle progression. (112,157) This investigation expands upon these previous studies by reporting the resumption of cell cycle progression upon intracellular GSH recovery. This study is also unique from prior cell cycle studies in that assays were conducted to determine both total GSH and GSSG. Evaluation for GSSG is vital to determine if intracellular oxidation has occurred. These results are in good agreement with earlier reports with regard to both cellular

GSH levels, and the magnitude of cellular GSH increases as cells progress beyond G1. (112,157) Further, there is consistency between this data and a recent report by Fidelus et al. (51) In both the Fidelus and this investigation, cellular depletion of GSH significantly affected cell cycle progression beyond G1. (51) Also, in both the Fidelus and this study, a linear relationship was noted between GSH and a parameter of cell cycle progression; Fidelus et al. reported a linear relationship between GSH levels and ^3H -thymidine incorporation, this study found a correlation between cellular GSH levels and the DNA index. (51)

The purpose of this investigation was to evaluate if the cellular thiol redox status is a determinant in how cells in G1 respond to mitogens. The results imply that cellular thiol redox status is a critical intracellular parameter that affects how mitogens and cells in G1 interact.

In separate studies, investigators have reported the increase in thiol dependent processes e.g. activities of DNA pol and RNR, during G1 through S. (28,53,98,132) As this cell cycle time frame coincides with the cellular increase in GSH levels, it is unlikely that this finding is coincidental. Instead, it is proposed that there exists an association between these events; as GSH is integral in the

provision of the reducing equivalents necessary to maintain these thiol dependent processes.

This study employed a modification of the ethionine synchronization protocol monitoring cellular GSSG levels.

The results showed that cellular levels of GSSG remained nondetectable in all experimental groups throughout the study. The significance of this finding is twofold. First, the lack of increase in GSSG revealed that cellular GSH-S-transferase activity was intact and that the cells responded to 2 CHX-1-one via the appropriate, enzymatic clearance. (94,113,175) Further, the lack of retention of GSSG eliminated any nonspecific protein mixed disulfide formation as a consequence of our GSH depletion protocol; thereby ruling out a generalized protein thiol oxidation as the cause for delayed growth arrest recovery in GSH depleted cultures.

This investigation reports a delayed growth arrest recovery in GSH depleted fibroblasts, in conjunction with the resumption of cell cycle progression upon restoration of cellular GSH levels. Although these findings suggest an association between cellular GSH status and ongoing cell cycle progression, it is proposed that GSH is not the sole contributor in this association. Cellular GSH status is interdependent with the cellular redox state. (94,113) Via glutaredoxin and thioredoxin, both GSH and NADPH contribute the necessary reducing equivalents used by RNR

during DNA synthesis. (73,74,75) Further, the activity of GSH reductase is dependent upon NADPH to reduce GSSG.

(94,113) Given the magnitude of increase in thiol dependent processes as cells progress from G1 through S, it is appropriate that both the cellular redox state and GSH status would be involved in maintaining these enzymatic activities.

Depletion of cellular GSH levels effectively removed an important source of reducing equivalents. Therefore, a delayed entrance into S phase would be anticipated as activities of both DNA pol and RNR would be restricted.

(18,53,98,132) However, following cellular GSH recovery, both glutaredoxin and GSH would be available for provision of the reducing equivalents necessary for function of RNR and DNA pol .

These findings support an association between cellular GSH levels and cell cycle progression beyond G1. It is suggested that a prominent factor in this association is the cellular thiol redox status, via its maintenance of the activities of 2 enzymes integral during late G1 through S, DNA pol and RNR.

The cell cycle distribution and redox poise results showed that ethionine performed two experimental functions. First, the ethionine synchronization provided a G1, stage specific population baseline. Second, exposure to ethionine perturbed the cellular redox poise, which was one

of the intracellular parameters that was to be investigated. Ethionine caused an oxidation of the cellular redox state and an increase in cellular total nicotinamide nucleotide levels. These redox poise alterations were noted in cultures harvested immediately after the ethionine synchronization, and in synchronized, GSH depleted (S-D 24) cultures. Both of these redox poise changes may be attributable to a bioenergetic perturbation. The intracellular availability of adenine nucleotides would be affected by the presence of ethionine. (108) The increased intracellular oxidation should stimulate reducing equivalent generating enzymes e.g. isocitrate dehydrogenase, malic enzyme, and the first two enzymes of the hexose monophosphate shunt, therefore accounting for the increases in total nicotinamide nucleotides. (114) Because the depletion of cellular GSH removes an important source of reducing equivalents, the S-D cultures faced an additional obstacle enroute to thiol redox recovery. (94,113) This was reflected by the delayed recovery from growth arrest in the S-D cultures.

The redox poise findings compared favorably with reported values. (89) Results obtained from the S and S-D 24 cultures agreed well in both total and reduced/oxidized distributions with other "stressed" cells - those harvested from fasted animals. (89) The L and S 24 total

nicotinamide nucleotide levels compared well to results obtained from cells harvested during the fed state. (89)

The $[Ca^{2+}]_i$ levels obtained during the resting and B FGF challenged assays are in good agreement with previously published reports. (18,27,133) Significant intergroup differences, which were related to the cellular redox state, were detected in the B FGF Ca^{2+} mobilization assays. The decreased Ca^{2+} mobilization observed in the S and S-D 24 groups may reflect a perturbation of the interrelated cellular redox poise and bioenergetic status. During mitogenic signal transduction, many growth factors promote Ca^{2+} mobilization in conjunction with activation of the Na^+/H^+ antiport. (119,120,129) The ATP may not be available to stimulate the monovalent ion fluxes in these bioenergetically perturbed cultures.

Variability in cellular growth factor expression has been cited as a potential cause for differences in $[Ca^{2+}]_i$ responses. (18) Because cellular harvesting and mitogen response assays were conducted in an identical fashion for all groups, any variations in receptor numbers or expression would be due to intrinsic cellular differences. The differences noted in mitogenic responsiveness may indeed reflect a growth factor receptor down regulation in those cultures that were less mitogen-responsive-groups S and S-D 24.

The B FGF stimulated Ca^{2+} results showed intragroup variability, as depicted by the standard deviations within each group. This study was conducted on human cell strains, not established cell lines. Therefore, variability among individual cell strains would be expected. It is well accepted that cellular populations are comprised of various subpopulations, which include cells that are heterogeneous in their responses to Ca^{2+} mobilization. (27,55) Via the fura-fluorometric assay, significant differences were detected among the biochemical averages of the experimental groups. The flow cytometric DNA analyses complimented the fura data by establishing that cellular subpopulations existed in regard to mitogenic responses.

Specific segments of the endoplasmic reticulum, some of which communicate with the plasma membrane, are proposed to provide the Ca^{2+} source for the transient, mitogen-induced, increases in $[\text{Ca}^{2+}]_i$. (78) Previous publications have demonstrated an association between the regulation of Ca^{2+} release from mitochondria and the redox state of nicotinamide nucleotides. (162,144,101) Oxidation of the redox state promoted a mitochondrial Ca^{2+} efflux. (101,144) In addition, investigators that reported other cation membrane transport studies, proposed that NAD(P)H, or a group in close redox communication, e.g. a membrane bound thiol, functions to restrict cation flow through

membrane channels. (86,87) Further, microsomal Ca^{2+} sequestration has been shown to be perturbed by reagents that promote oxidation, or diminish ATP levels. (170) Both of these cellular alterations, depletion of ATP and redox poise oxidation, could result from ethionine exposure. (109)

In this study, resting $[\text{Ca}^{2+}]_i$ levels were comparable; while differences were noted in the "mitogen-responsive" Ca^{2+} pool. Volpe et al. recently proposed that discrete "calciosomes" comprise the responsive Ca^{2+} pool in the endoplasmic reticulum. (174) Because Ca^{2+} efflux in mitochondria was associated with an oxidized redox state, we propose two plausible explanations for the Ca^{2+} results. First, prior to mitogen interaction, an inhibition of Ca^{2+} sequestration occurred in the thiol redox perturbed cultures. The released Ca^{2+} would then be readily sequestered by mitochondria, thereby maintaining normal $[\text{Ca}^{2+}]_i$. Alternatively, the Ca^{2+} loading into the responsive pool may not have occurred in those cells with an oxidized redox poise if it is an ATP dependent process. In either situation, a mitogen responsive pool of Ca^{2+} would not exist in the thiol redox perturbed cultures.

This study evaluated two parameters of the mitogenic response. Ca^{2+} mobilization and cell cycle progression were the immediate and sustained parameters, respectively. These mitogenic responses were interrelated with the

cellular thiol redox state. Cultures with increased GSH and reduced nicotinamide nucleotide levels showed Ca^{2+} mobilization also showed cell cycle progression; implying a complete mitogenic signal transduction.

Because cells employ an enzymatic means of rapid response, mitogenic signal transduction is a rapidly occurring event. (135) A cascade of kinases, which serve to transmit and amplify the initial message, are activated during the mitogenic response. (135) Because many of these kinases are Ca^{2+} dependent, kinase cascade stimulation is dependent upon an increase in $[\text{Ca}^{2+}]_i$. (135) To transduce the mitogenic signal, a cell must transiently increase $[\text{Ca}^{2+}]_i$ to augment the activation of the kinase cascade. (135) (Fig. 12)

In conclusion, I propose that the cellular response to those mitogenic stimuli that function via inositol phosphates is interdependent upon cellular thiol redox status. Cells that have a cellular thiol redox status below a threshold response point, may have compromised Ca^{2+} mobilization, and therefore be incapable of triggering the mitogen induced cascade.

Figure 12

Schematic diagram showing proposed mechanism of the association of cellular thiol redox status with mitogenic signal transduction.

Figure 12

LIST OF REFERENCES

1. Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., Watson, J.D.: In: Molecular Biology of the Cell. Garland Publishing Inc., N.Y. pp. 640-645, 1983.
2. Balk, S.D., Whitfield, J.F., Yondale, T., Braun, A.C.: Proc. Natl. Acad. Sci U.S.A. 70:675-679, 1973.
3. Balkow, K., Hunt, T., and Jackson, R.J. Biochem. Biophys. Res. Commun. 67, 366-375, 1975.
4. Baserya, R. The Biology of Cell Reproduction. Harvard Univ. Press, Cambridge, MA., 1985.
5. Baum, H. and Al-Shaikhaly, M. Biochem. Soc. Trans. 5, 1094-1095, 1977.
6. Becker, G.L., Fiskum, G., Lehninger, A.L. J. Biol. Chem. 255:9009-9012, 1980.
7. Bellomo, G., Jewell, S.A., Thor, H. and Orrenius, S. Proc. Natl. Acad. Sci. USA 79, 6842-6846, 1982.
8. Berridge, M.J. Biochem. J. 212:849-858, 1983.
9. Berridge, M.J., Irvine, R.F.: Nature 312:315-321, 1984.
10. Berridge, M.J.: Biochem. J. 220:345-360, 1984.
11. Berridge, M.J., Buchan, P.B., Heslop, J.P. Mol. Cell. Endocrinol. 36:37-42, 1984.
12. Boynton, A.L. Cooney, R.V., Hill, T.D., Nilsson, T., Arkhammar, P. and Berggren, P.O. Exp. Cell Res. 181, 245-255, 1989.
13. Boyer, P.D., and Krebs, E.G., Eds. The Enzymes Academic Press, New York, ed 3, 1986. vol. 17.
14. Burgess, G.M., McKinney, J.S., Fabiato, A., Leslie, B.A., Putney, J.W. Jr. J. Biol. Chem. 258; 15336-15345, 1983.

15. Burk, R.F. Nishiki, K., Lawrence, R.A., Chance, B. J. Biol. Chem. 253, 43-46, 1978.
16. Bygrave, F.L. Curr Top. Bioenerg. 6, 259-318, 1977.
17. Bygrave, F.L. Biol. Rev. 53:43-79, 1978.
18. Byron, L.K. and Villereal, M.L.: J. Biol. Chem. 264:18234-18239, 1989.
19. Campisi, J., Morreo, G., Pardee, A.B. Exp. Cell RES. 152, 459, 1984.
20. Campisi, J. and Pardee, A.B. Mol. Cell. Biol. 4, 1807-1814, 1984.
21. Carlberg, I., Mannervik, B. Biochim. Biophys. Acta. 484, 268-274, 1977.
22. Cernoch, M. and Weinbergova, O. Physiol. Bohemoslov., 18, 161-165, 1969.
23. Cerutti, P. A. Science, 227, 375-381, 1985.
24. Chance, B., Sies, Sies, H., Boveris, A. Physiol. Rev. 59, 527-605, 1979.
25. Chen, Y. and Rabinovitch, P.S. J. Cell. Physiol. 140, 59-67, 1989.
26. Chin, K.V., Cade, C., Brostrom, C.O., Galuska, E.M., Brostrom, M.A., J. Biol. Chem. 34, 16509-16514, 1987.
27. Chused, T.M., Wilson, H.A., Greenblatt, D., Ishida, Y., Edison, L.J., Tsien, R.Y. and Finkleman, F.D.: Cytometry 8:396-404, 1987.
28. Collins, J.M. and Chu, A.K. Binding of the DNA polymerase - DNA Primase Complex to the Nuclear Matrix in HeLa Cells. Biochem, 26:5600-5607; 1987.
29. Colwill, R.W. and Sheinin, R. Proc. Natl. Acad. Sci. U.S.A. 80,4644-4648, 1983.
30. Cross, C.E. Ann. Intern. Med. 107, 526-545, 1987.
31. Cross, F. Weintraub, H., Roberts, J.: Annu. Rev. Cell Biol. 5:341, 1989.
32. Dan, K. In "Cell Synchrony" (I.L. Cameron and G.M. Padilla, eds.), pp 307-327, Academic Press, N.Y., 1966.

33. Dawson, A.P., Comerford, J.G., Fulton, D.V. *Biochem. J.* 234:311-315, 1986.
34. Denhart, D.T., Edwards, D.R., Parfett, C.L.J. *Biochim. Biophys. Acta* 865, 83, 1986.
35. Denton, R.M. and McCormack, J.G., Edgell, N.J. *Biochem. J.* 190:107-117, 1980.
36. Dholakia, J.N., Mueser, T.C., Woodley, C.L., Parkhurst, L.J. and Wahba, A.J. *Proc. Natl. Acad. Sci. U.S.A.* 83, 6746-6750, 1986.
37. Donachie, W.D. in *The Cell Cycle* (John. P.C. L., ed) Cambridge University Press, London.
38. Downes, C.P., Mussat, M. C., Michell, R.H. *Biochem. J.* 203:169-177, 1982.
39. Draetta, G. Brizuela, L., Moran, B., Beach, D. *Cold Spring Harbor Symposia*, 53, 195-201, 1988.
40. Drahota, Z., Carafoli, E., Rossi, C.S., Gamble, R.L., and Lehninger, A.L. *J. Biol. Chem.* 240, 2712-2720, 1965.
41. Eagon, R.G. *J. Bacteriol.* 83, 736-737, 1962.
42. Edmunds, L.N. *Cell Cycle Clocks*, Marcel Dekker, New York, 1984.
43. Engstrom, S., Graslund, A., Skog, S., Thelander, L., and Tribukait, B. *J. Biol. Chem.* 260:9114-9116; 1985.
44. Erikson, S., Graslund, A., Skog, S., Thelander, L., Tribukait, B. *J. Biol. Chem.* 259:11695-11700; 1984.
45. Ernst, V., Levin, D.H., and London, I.M. *J. Biol. Chem.* 253, 7163-7172, 1978.
46. Ernst, V., Levin, D.H. and London, I.M. *Proc. Natl. Acad. Sci. U.S.A.* 76, 2118-2122, 1979.
47. Eyer, P. and Podhradsky, D.: *Anal. Biochem.* 153:57-66, 1986.
48. Fabiato, A.: *J. Physiol. (paris)* 75:463-505, 1979.
49. Fabiato, A.: *J. Gen. Physiol.* 78:457-497, 1981.
50. Fabiato, A.: *J. Gen. Physiol.* 85:247-289, 1985.

51. Fidelus, R.K., Ginouves, P., Lawrence, D. and Isan, M.F.
52. Flohe, L. CIBA Found. Symp. 65, 95-122, 1979.
53. Foster, K.A. and Collins, J.M. J. Biol. Chem. 260:4229-4235; 1985.
54. Freifelder, D. Molecular Biology: A Comprehensive Introduction to Procaryotes and Eucaryotes. Ch. 13 Translation: The Machinery and Chemical Nature of Protein Synthesis. The Machinery and Chemical Nature of Protein Synthesis. Jones and Bartlett Publishers, Boston, Mass. pp 535; 1983.
55. Gonzalez, F.A., Gross, D.J. Heppel, L.A. and Webb, W.W.: J. Cell. Physiol. 135:269-276, 1988.
56. Gospodarowicz, D., Neufeld, G., Schweigerer, L.: Molecular and Cell. Endocrin. 46:187-204, 1986.
57. Gospodarowicz, D., Neufeld, G., Schweigerer, L.: J. Cell. Physio. Supplement 5:15-26, 1987.
58. Granner, D.K., From: Harper's Review of Biochemistry. Lange Medical Publications. Los Altos Ca. Ch. 36, 505-514, 1985.
59. Griffith, O.W., Meister, A. Proc. Natl. Acad. Sci. U.S.A. 77, 3384-3387, 1980.
60. Griffith, O.W. and Meister, A. J. Biol. Chem. 254; 7558-7560, 1979.
61. Gryniewicz, G., Poenie, M., and Tsien, R.: J. Biol. Chem. 260:3440-3450, 1985.
62. Hagan, I., Hayles, J., Nurse, P. J. Cell Biol. 91, 587, 1988.
63. Hammarston, E., and Reichard, P. J. Biol. Chem. 183, 105-109, 1950.
64. Harington, C.R. and Mead, T.H. Biochem. J. 29, 1602-1617, 1935.
65. Harris, E.J., Al-Shaikholy, M., Baum, H. Biochem. J. 182, 455-464, 1979.
66. Harris, J.W. and Patt, H.M. Exp. Cell. Res. 56:134-151; 1969.

67. Harris, J.W. and Teng, S.S.: J. Cell. Physiol. 81:91-95, 1973.
68. Harvey, G. and Pearson, C.K. J. Cell. Physiol. 134, 25, 1988.
69. Hesketh, T.R., Smith, G.A., Houslay, M.D., Warren, G.B., Metcalfe, J.C. Nature, 267, 490-494, 1977.
70. Hesketh, T.R., Moore, J.P., Morris, J.D.H.: Nature 313:481-484, 1985.
71. Hesketh, T.R., Morris, J.D.H., Moore, J.P., Metcalfe, J.C. J. Biol. Chem. 263, 11879-11886, 1988.
72. Hill, T.D., Zwiller, J., Boynton, A.L. J. Cell. Physiol. 140, 403-407, 1989.
73. Holmgren, A. J. Biol. Chem. 254:3664-3671; 1979.
74. Holmgren, A. J. Biol. Chem. 254:3672-3678; 1979.
75. Holmgren, A. Curr. Top. Cell. Reg. 19:47-76; 1981.
76. Hopkins, F.G. Biochem. J. 15, 286-305, 1921.
77. Hunter, D.R., Haworth, R.A., and Southard, J.H. J. Biol. Chem. 251, 5069-5077, 1976.
78. Irvine, R.F.: Br. Med. Bull. 42:369-374, 1986.
79. Jackson, R.J., Campbell, E.A., Herbert, P. and Hunt, T. Eur. J. Biochem. 131, 289-301, 1983.
80. Jakoby, W.B. ed. Enzymatic Basis of Detoxification, Vol. 2, New York: Academic, 1980.
81. Jakoby, W.B. ed. Methods Enzymol. 77, 1-476, 1981.
82. Jenh, C.H., Rao, L.G., and Johnson, L.F. J. Cell. Physiol. 122, 149-154, 1985.
83. Joseph, S.K., Thomas, A.P., Williams, R.J., Irvine, R.F., Williamson, J.R.: J. Biol. Chem. 259:3077-3081, 1984.
84. Joseph, S.K., Rice, H.L., and Nicchitta, C.V. Biochim. Biophys. Acta. 945:185-194, 1988.
85. Joseph, S.K. and Williamson, J.R. Arch. Biochem. Biophys. 273, 1-15, 1989.

86. Jung, D. and Brierley, G.P.: J. Biol. Chem. 256:10490-10496, 1981.
87. Jung, D., and Brierley, G.P.: Biochem. Biophys. Res. Comm. 106:1372-1377, 1982.
88. Kaibuchi, K. Tsuda, T., Kikuchi, A., Tanimoto, T., Yamashita, T. and Takai, Y. J. Biol. Chem. 261, 1187-1192, 1986.
89. Kalhorn, T.F., Thummel, K.E., Nelson, S.D., Slattery, J.T.: Anal. Biochem. 151:343-347, 1985.
90. Kan, B., London, I.M. and Levin, D.H. J. Biol. Chem. 263, 15652-15656, 1988.
91. Kelly, K., Cochran, B.H., Stiles, C.D., Leder, P.: Nature 313:481-484, 1985.
92. Kosower, N.S., Vanderhoff, G.A., Kosower, E.M. Biochim. Biophys Acta, 272, 623-637, 1972.
93. Kosower, N.S. and Kosower, E.M. "Glutathione" (L. Flohe' et al., eds.), pp 276-287, Thieme, Stuttgart, 1974.
94. Kosower, N.S., Kosower, E.M.: Int. Rev. Cytol. 54:109-160, 1978.
95. Krammer, G. Diplomarbeit, Eberhard-Karls-Univ., Tübingen, 1975.
96. Labbe, J.C., Lee, M.G., Nurse, P., Picard, A., Doree, M. Nature, 355, 251, 1988.
97. Larsen, J.K., Munch-Petersen, B., Christensen, J., Jorgensen, R.: Cytometry 7:54-63, 1986.
98. Laskey, R.A., Fairman, M.P., Blow, J.J. Science 246:609-613; 1989.
99. Lee, F.Y., Siemann, D.W., Allalunis-Turner, M.J., Kang, P.C. Cancer Res. 48, 3661-3665, 1988.
100. Lehninger, A.L., Carafoli, E., and Rossi, C.S. Adv. Enzymol. 29, 259-320, 1967.
101. Lehninger, A.L., Vercesi, A., Bababunmi, E.N.: Proc. Natl. Acad. Sci., USA, 75:1690-1694, 1978.
102. Lenz, J.R., Chatterjee, G.E. Maroney, P.E., and Baglioni, C. Biochemistry 17, 80-87, 1978.

103. Levin, D.H., Ranu, R.S., Ernst, V., and London, I.M.
Proc. Natl. Acad. Sci. U.S.A. 73, 3112-3116, 1976.
104. Lloyd, D. Biochem. J. 242, 313-321, 1987.
105. Lotscher, H.R., Winterhalter, K.H., Carafoli, E., and
Richter, C. Proc. Natl. Acad. Sci. USA, 76, 4340,
1979.
106. Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall,
R.J.: J. Biol. Chem. 193:265-275, 1951.
107. Mano, Y. Arch. Biochem. Biophys. 146, 237-248, 1971.
108. Mayes, P.A. From: Harper's Review of Biochemistry.
Lange Medical Publications Los Altos, Ca. Ch. 18.
Metabolism of Lipids, 244-45, 1985.
109. Mayes, P.A. From: Harper's Review of Biochemistry.
Lange Medical Publications Los Altos, Ca. Cha. 12
Biological Oxidation 128-146, 1985.
110. Masui, Y. and Markert, C.J. Exp. Zool. 177, 129,
1971.
111. Matsumoto, K., Uno, I. and Ishikawa, T. Exp. Cell
Res. 146, 151-161; 1983.
112. Mbemba, F., Houbion, A., Raes, M., Remacle, J.
Biochem. Biophys. Acta 838:211-220; 1985.
113. Meister, A. and Anderson, M.E.: Annu. Rev. Biochem.
52:711-760, 1983.
114. Metcalfe, J.C., Hesketh, T.R., Smith, G.A., Morris,
J.D. H., Corps, A.N., Moore, J.P.: J. Cell Sci
3:199-228, 1985.
115. Metcalfe, J.C., Moore, J.P., Smith, G.A., Hesketh,
T.R. Br. Med. Bull. 42, 405-412, 1986.
116. Meuller, G.C.: Fed. Proc. 28:1780-1789, 1969.
117. Michelson, A.M., Ernst, V., Levin, D.H., and London,
I.M. J. Biol. Chem. 259:8529-8533, 1984.
118. Mitchell, R.H. Biochim. Biophys. Acta. 415:81-147,
1975.
119. Moolenaar, W.H., Tertoolen, L.G.J., and de Laat,
S.W.: J. Biol. Chem. 259:8066-8069, 1984.

120. Moolenaar, W.H., Aerts, R.J., Tertoolen, L.G.J. and de Laat, S.W.: J. Biol. Chem. 261:279-284, 1986.
121. Muallem, S., Schoeffield, M., Pandol, S., Sachs, G. Proc. Natl. Acad. Sci. USA 82:4433-4437, 1985.
122. Murphree, S., Stubblefield, E., Moore, E.C. Exp. Cell Res. 58:118-124; 1969.
123. Murray, A.W. Nature 326, 542, 1987.
124. Murray, A.W. and Kirschner, M.W.: Science 246:614-621, 1989.
125. Nelkin, B., Nichols, C., Vogelstein, B. FEBS Let. 109, 233, 1980.
126. Neufield, E.F. and Mazia, D. Exp. Cell Res. 13, 622-624, 1957.
127. Nicholls, D.G.: Br. Med. Bull. 42:353-358, 1986.
128. Nowell, P.C.: Science 194:23-28, 1976.
129. Ober, S.S. and Pardee, A.B.: J. Cell. Physiol. 132:311-317, 1987.
130. Oehlert, W.: Cell Tiss. Kinet. 6:325-335, 1973.
131. Oshino, N. and Chance, B. Biochem J. 162:509-525, 1977.
132. Ottiger, H.P. and Hubocher, U. Proc. Natl. Acad. Sci., USA, 81:3993-3997; 1984.
133. Pandiella, A. Magni, M. and Meldolesi, J.: Biochem. Biophys. Res. Comm. 163:1325-1331, 1989.
134. Pardee, A.B. Cancer Res. 47, 1488, 1987.
135. Pardee, A.B.: Science 246:603-608, 1989.
136. Pitot, H.C. Fundamentals of Oncology 34d edition Marcel Dekker, Inc. N.Y., N.Y., 1986.
137. Poenie, M., Alderton, J., Tsien, R.Y., Steinhardt, R.A.: Nature 315:147-149, 1985.
138. Pollard, H., Menard, R., Brandt, H.A. Pazoles, C.J. Creutz, C.E., Ramu, A. Anal. Biochem. 86:761-763, 1978.

139. Post, G.B., Keller, D.A., Connor, K.A., Menzel, D.B.
Biochem. Biophys. Res. Comm. 114, 737-742, 1983.
140. Precott, D. Int. Rev. Cytol. 100, 93, 1987.
141. Prentki, M., Corkey, B.E., Matchinsky, F.M.: J.
Biol. Chem. 260:9185-1990, 1985.
142. Principe, P., Wilson, G.D., Riley, P.A., Slater, T.F.
Cytometry 10:750-761, 1989.
143. Pringle, J.R. Curr. Opin. Cell Biol. 1, 237, 1989.
144. Prpic, V. and Bygrave, F.L.: J. Biol. Chem.
255:6193-6199, 1980.
145. Rao, P.N. and Johnson, R.T. Nature, 225, 159, 1970.
146. Rapkine, L.: Ann. Physiol. Physiochim. Biol. 7:382-
417, 1931.
147. Reed, S. and Nasmyth, K.A. Proc. Natl. Acad. Sci. 77,
2119, 1981.
148. Reinhart, P.H., Van de Pol, E., Taylor, W.M.,
Bygrave, F.L. Biochem. J. 218:415-420, 1984.
149. Reynhout, J.K. and Smith, L.D. Dev. Biol. 38, 394,
1974.
150. Rodwell, V.W. In: Harper's Review of Biochemistry
20th edition Ch 3 Amino Acids and Peptides, Ch 4
Proteins, Lange Medical Publications. Los Altos, Ca.
99 21-24; 32-41, 1985.
151. Ross, M. and Ballard, F.J. Biochem. J. 249, 363-368,
1988.
152. Rossi, C.S. and Lehninger, A.L. J. Biol. Chem. 239,
3971-3980, 1964.
153. Rossow, P.W., Riddle, V.G.H., Pardee, A.B. Proc.
Natl. Acad. Sci. U.S.A. 76, 4446, 1979.
154. Rozengurt, E. Science, 234, 161, 1986.
155. Sakai, H. and Dan, K. Exp. Cell Res. 16, 24-41, 1959.
156. Sedlak, J. and Lindsay, R.H. Anal. Biochem. 25:192-
205, 1968.

157. Shaw, J.P. and Chou, I.N.: J. Cell. Physiol. 129:193-198, 1986.
158. Shears, S.B. and Kirk, C.J. Biochem. J. 220:417-421, 1984.
159. Sherley, J.L. and Kelly, T.J. J. Biol. Chem. 263, 8350-8358, 1988.
160. Simons, S.S. Jr., Johnson, D.F. Anal. Biochem. 90:705-725, 1978.
161. Smith, J.B., Smith, L., Higgins, B.L. J. Biol. Chem. 260:14413-14416, 1985.
162. Somylo, A.P., Bond, M., Sombyo, A.V.: Nature 314:622-625, 1985.
163. Spat, A., Fabiato, A., Rubin, R.P. Biochem. J. 233:929-932, 1986.
164. Streb, H. and Schulz, I. Am. J. Physiol. 245:G347-G357, 1983.
165. Streb, H., Irvine, R.F., Berridge, M.J., Schulz, I.: Nature 306:67-69, 1983.
166. Stolze, H. and Schulz, I. Am. J. Physiol. 238, G338-G348, 1980.
167. Sunkara, P.S., Wright, D.A., Rao, P.N. Proc. Natl. Acad. Sci. U.S.A. 76, 2799, 1979.
168. Tachibana, K., Yanagishima, N., Kishimoto, T. J. Cell Sci. 88, 273, 1987.
169. Tan, K-N., Tashjian, A.H. J. Biol. Chem. 256, 8994-9002, 1981.
170. Thor, H., Hartzell, P., Svensson, S.A., Orrenius, S. Biochem. Pharmacol. 34:3717-3723, 1985.
171. Tricoli, J.V., Sahai, B.M., McCormick, P.J., Jarlinski, S.J., Bertram, J.S., and Kowalski, D. Exp. Cell Res. 158, 1-14, 1985.
172. Vercesi, A.E., Ferraz, V.L., Macedo, D.V. and Fiskum, G. Biochem. Biophys. Res. Comm. 154, 934-941, 1988.
173. Volpe, P., Salviati, G., Di Virgilio, F., Pozzan, T. Nature 316:347-349, 1985.

174. Volpe, P., Kraus, K.H., Hashimoto, S., Zorzato, F., Pozzan, T., Meldelosi, J., and Lew, D.P.: Proc. Natl. Acad. Sci. 85:1091-1095, 1988.
175. Wedner, H.J., Simchowicz, L., Stenson, W.F., Fischman, C.M. J. Clin. Invest. 68:535-543, 1981.
176. West, D.K., Lenz, J.R. and Baglioni, C. Biochemistry, 18, 624-632, 1979.
177. Wilke, M.S., Hsu, B.M., Willie, J.J., Pittelkow, M.R., Scott, R.E.: Am. J. Path. 131:171-181, 1988.
178. Wilke, M.S., Hsu, B.M., Scott, R.E.: Lab. Invest. 58:660-666, 1988.
179. Williamson, J.R., Cooper, R.H., Hoek, J.B. Biochim. Biophys. Acta. 639:243-295, 1981.
180. Yang, H.C. and Pardee, A.B. J. Cell. Physiol. 127, 410, 1986.
181. Yarden, Y. and Ullrich, A. Annu. Rev. Biochem. 57, 443, 1988.
182. Ziegler, D.M.: Annu. Rev. Biochem. 54:305-329, 1985.

APPENDIX

Table 6

Raw Data from Ca^{2+} Mobilization Experiments

Experimental Group	pre B FGF [Ca^{2+}] _i (nM)	% Increase in [Ca^{2+}] _i	after B FGF [Ca^{2+}] _i (nM)
L Intrinsic	252 174 252 204 180 x=213±39	29.9 15.2 12.8 11.5 19.9 x=17.8±7.5 (S.D.)	327 200 284 227 216 x=251±53 (S.D.)
L Challenge	390 330 404 316 x=360±44 (S.D.)	14.1 11.3 14.1 12.3 x=13.0±1.4 (S.D.)	445 367 461 355 x=407±54 (S.D.)
S Intrinsic	190 183 235 228 x=209±26 (S.D.)	10.3 9.0 7.6 7.5 x=8.6±1.4 (S.D.)	210 199 253 245 x=227±26 (S.D.)
S Challenge	374 318 306 386 x=346±40 (S.D.)	5.6 9.2 4.0 1.4 5.1±3.3 (S.D.)	395 347 318 391 x=367±37 (S.D.)
S-24 Intrinsic	171 182 245 234 x=208±37 (S.D.)	10.2 17.7 14.6 29.3 x=17.9±8.2 (S.D.)	188 214 281 303 x=247±54 (S.D.)

Table 6 (continued)

Experimental Group	pre B FGF [Ca ²⁺] _i (nM)	% Increase in [Ca ²⁺] _i	after B FGF [Ca ²⁺] _i (nM)
S-24 Challenge	382	5.7	404
	306	18.4	362
	370	22.5	453
	318	20.1	382
	x=344±38 (S.D.)	x=17.0±7.5 (S.D.)	x=400±39 (S.D.)
S-D 24 Intrinsic	175	5.6	185
	166	11.0	184
	226	4.1	235
	217	8.7	236
	x=196±30 (S.D.)	x=7.4±3.1 (S.D.)	x=210±29 (S.D.)
S-D 24 Challenge	320	8.0	346
	371	0.9	374
	380	4.4	397
	329	5.6	347
	x=350±30 (S.D.)	x=4.7±3.0 (S.D.)	x=366±24 (S.D.)

Table 7

Statistical Analysis for Ca^{2+} Mobilization Data

For these analyses, I used two Non-Parametric tests, the Kruskal-Wallis One Way Analysis of Variance, and the Mann-Whitney U. The Kruskal-Wallis test was used first, to determine if there were any differences between the groups. The Mann Whitney U test was then conducted to determine between which groups the differences existed. Being Non-Parametric tests, these analyses are appropriate to run on an outbred population - e.g. cells obtained from human donors.

Experimental Groups: L, S, S 24, S-D 24.

Score = Percentage increase in $[\text{Ca}^{2+}]_i$ after addition of
B FGF

Scores are ranked low to high.

L		Intrinsic S		S 24		S-D 24	
Score	Rank	Score	Rank	Score	Rank	Score	Rank
29.9	17	10.3	8	10.2	7	5.6	2
15.2	13	9.0	6	17.7	14	11.0	9
12.8	11	7.6	4	14.6	12	4.1	1
11.5	10	7.5	3	29.3	16	8.7	5
19.9	15						

$$\begin{array}{llll}
 (R_L)^2 - N_L = & (R_S)^2 - N_S = & (R_{S\ 24})^2 - N_{S24} = & (R_{SD24})^2 - \\
 N_{SD24} = & & & \\
 871.2 & 110.3 & 600.3 & 72.3
 \end{array}$$

$N = \text{Total Nos. Scores} = 17$
 $\text{Sum of } R^2/N = 64.8$
 $64.8 - 54 = 10.8 \quad df=3$
 Significant $p < 0.05$

Table 7 (continued)

L		Challenge S		S 24		S-D 24	
Score	Rank	Score	Rank	Score	Rank	Score	Rank
14.1	12	5.6	5	5.7	7	8.0	8
11.3	10	9.2	9	18.4	14	0.9	1
14.1	13	4.0	3	22.5	16	4.4	4
12.3	11	1.4	2	20.1	15	5.6	6
$(R_L)^2 - N_L =$ $N_{S-D24} =$ 529		$(R_S)^2 - N_S =$ 90.3		$(R_{S24})^2 - N_{S24} =$ 676		$(R_{S-D24})^2 -$ 90.3	

$N = \text{Total Nos. Scores} = 16$
 $\text{Sum } R^2 / N = 1385.5 \times 12/16 (17)$
 $= 60.9 - 3 (17) = 9.9 \quad df = K-1=3$
 Significant $p < 0.05$ level

Mann Whitney U

Group L vs Group S
 Intrinsic $p < 0.08$
 Challenge $p < 0.014$

Group L vs Group S-D 24
 Intrinsic $p < 0.08$
 Challenge $p < 0.014$

Group S 24 vs Group S
 Intrinsic and Challenge $p < 0.029$

Group S 24 vs Group S-D 24
 Intrinsic and Challenge $p < 0.029$

(Mann Whitney U is a ranked score comparison conducted between the individual groups).